2/ V.2-A

aval Education and
aining Command

NAVEDTRA 10062
MAJOR REVISION
February 1989
0502-LP-215-3700

Training Manual
(TRAMAN)

Mathematics, Volume 2-A

ISTRIBUTION STATEMENT A: Approved for public release; distribution is unlim

onfederal government personnel wanting a copy of this document
ust use the purchasing instructions on the inside cover.

The terms training manual (TRAMAN) and nonresident training course (NRTC) are now the terms used to describe Navy nonresident training program materials. Specifically, a TRAMAN includes a rate training manual (RTM), officer text (OT), single subject training manual (SSTM), or modular single or multiple subject training manual (MODULE); and a NRTC includes nonresident career course (NRCC), officer correspondence course (OCC), enlisted correspondence course (ECC) or combination thereof.

Although the words "he," "him," and "his" are used sparingly in this manual to enhance communication, they are not intended to be gender driven nor to affront or discriminate against anyone reading this text.

DISTRIBUTION STATEMENT A: Approved for public release; distribution is unli

Nonfederal government personnel wanting a copy of this document must write to Superintendent of D Government Printing Office, Washington, DC 20402 OR Commanding Officer, Naval Publications and For

PREFACE

This training manual (TRAMAN), *Mathematics*, Volume 2-
NAVEDTRA 10062, and its nonresident training course (NRTC
NAVEDTRA 80062, form a self-study training package. T
purpose of this training package is to aid those personnel w
need an extension of the knowledge of mathematics. To serve t
wide variety of personnel needs, we have made the text gene
in nature; it is not directed toward any one specific specialit

The definitions and notations of logarithms followed
computations with logarithms occur early in the te
Trigonometric ratios and analysis and applications along with a
to computations occur next. Trigonometric identities a
equations are followed by vectors and forces.

To aid you in understanding the subject matter, we ha
presented numerous examples and practice problems throughc
the text; additional practice problems are provided at the end
each chapter.

The NRTC designed for use with this TRAMAN consists
individual assignments. Each assignment is a series of questic
based upon the textbook. You should study the TRAMAN pag
given at the beginning of each assignment before trying to answ
the questions in your NRTC.

Before attempting this course, you should already have
understanding of the fundamentals of algebra and trigonomet
A review of applicable chapters in *Mathematics*, Volume
NAVEDTRA 10069-D1, will be of great assistance to you
completing this course.

The TRAMAN is automatically packaged with the NRT
Ordering information is available in the *List of Training Manu
and Correspondence Courses*, NAVEDTRA 10061. However,
text alone may be ordered separately (to be used for train
sessions, etc.) from NPFC, Philadelphia.

This TRAMAN and associated NRTC were prepared by
Naval Education and Training Program Management Supp

Activity, Pensacola, Florida, for the Chief of Naval Educati
and Training. Technical review was provided by the Chief of Na
Technical Training, Millington, Tennessee; Commanding Offic
Naval Nuclear Power School (Enlisted Mathematics Divisio
Naval Training Center, Orlando, Florida; Commanding Offic
Service School Command, Great Lakes, Illinois; Commandi
Officer, Service School Command, San Diego, California; a
Superintendent, U.S. Naval Academy (Mathematics Departme
Annapolis, Maryland.

Your suggestions and comments concerning this TRAMA
and its NRTC are invited. Comment sheets have been inclu
with both the TRAMAN and NRTC.

1989 Edition

Stock Ordering N
0502-LP-215-37

Published by
NAVEDTRA EDUCATION AND TRAINING PROGRA
MANAGEMENT SUPPORT ACTIVITY

UNITED STATES
GOVERNMENT PRINTING OFFICE
WASHINGTON, D.C.: 1989

THE UNITED STATES NAVY

GUARDIAN OF OUR COUNTRY

The United States Navy is responsible for maintaining control of th sea and is a ready force on watch at home and overseas, capable o strong action to preserve the peace or of instant offensive action t win in war.

It is upon the maintenance of this control that our country's gloriou future depends; the United States Navy exists to make it so.

WE SERVE WITH HONOR

Tradition, valor, and victory are the Navy's heritage from the past. T these may be added dedication, discipline, and vigilance as th watchwords of the present and the future.

At home or on distant stations we serve with pride, confident in th respect of our country, our shipmates, and our families.

Our responsibilities sober us; our adversities strengthen us.

Service to God and Country is our special privilege. We serve wit honor.

THE FUTURE OF THE NAVY

The Navy will always employ new weapons, new techniques, an greater power to protect and defend the United States on the sea under the sea, and in the air.

Now and in the future, control of the sea gives the United States he greatest advantage for the maintenance of peace and for victory i war.

Mobility, surprise, dispersal, and offensive power are the keynotes o the new Navy. The roots of the Navy lie in a strong belief in th future, in continued dedication to our tasks, and in reflection on ou heritage from the past.

Never have our opportunities and our responsibilities been greater.

CONTENTS

CHAPTER P

1. Logarithms .....
2. Computations with Logarithms .....
3. Trigonometric Measurements .....
4. Trigonometric Analysis .....
5. Oblique Triangles .....
6. Trigonometric Identities and Equations ....
7. Vectors and Forces .....

APPENDIX

I. Common Logarithms of Numbers..... A
II. Natural Sines and Cosines..... A
III. Natural Tangents and Cotangents ..... AI

INDEX ..... INDE

CHAPTER 1

LOGARITHMS

LEARNING OBJECTIVES

Upon completion of this chapter, you should be able to do the following:

1. Define exponential form and logarithmic form.
2. Apply laws of multiplication, division, powers, and roots for calculating logarithms.
3. Determine the characteristic and mantissa of common logarithms.
4. Interpolate using logarithm tables.
5. Find common logarithms, antilogarithms, and natural logarithms using logarithm tables.

INTRODUCTION

The basic definitions and terminology associated with the study of logarithms were discussed in *Mathematics*, Volume 1, NAVEDTRA 10069-D1. Some of these basic topics are reviewed in the following paragraphs, followed by discussion of the use of logarithm tables and natural logarithms.

REVIEW OF DEFINITIONS

The most important definition to remember when dealing with logarithms is that *every logarithm is an exponent.* For example, since 3^2 is equal to 9, the logarithm of 9 to the base 3 is 2. Stating a logarithmic relationship requires that a base be stated or implied; the various exponents that designate powers of the base are logarithms to that base.

The usual method of expressing the basic definition of logarithms in symbols is as follows:

If b is a positive number other than 1 and x is a real number (positive, negative, or zero) in the equation $b^x = a$, *then* $x = \log_b a$. The two forms shown in the foregoing expression are defined as follows:

Exponential form: $b^x = a$

Logarithmic form: $x = \log_b a$

EXAMPLE: Change the expression $2^3 = 8$ to logarithmic form.

SOLUTION: If $b^x = a$, then $\log_b a = x$.

If

$$2^3 = 8$$

where

$$b = 2$$

$$x = 3$$

and

$$a = 8$$

then

$$\log_2 8 = 3$$

EXAMPLE: Change the expression $\log_{10} 100 = 2$ to exponential form.

SOLUTION: If $\log_b a = x$, then $b^x = a$.

If

$$\log_{10} 100 = 2$$

then

$$10^2 = 100$$

PRACTICE PROBLEMS:

1. Change $10^3 = 1{,}000$ to logarithmic form.
2. Change $e^x = N$ to logarithmic form.
3. Change $\log_2 4 = 2$ to exponential form.
4. Change $\log_{10} 3.16 = 1/2$ to exponential form.

ANSWERS:

1. $\log_{10} 1{,}000 = 3$
2. $\log_e N = x$
3. $2^2 = 4$
4. $10^{1/2} = 3.16$

LAWS FOR CALCULATION

The following two abilities are necessary for logarithmic calculation:

1. *Recognition of logarithms as exponents*
2. *Knowledge of the Laws for Logarithms*

The first of these abilities was discussed in the foregoing section. The second is the subject of the following paragraphs.

MULTIPLICATION

Law 1. *The logarithm of a product is equal to the sum of the logarithms of the factors.*

Suppose that we wish to multiply A and B, and we know the following:

$$A = 10^m$$

and

$$B = 10^n$$

Then the product AB is

$$AB = 10^m \times 10^n$$

$$= 10^{m+n}$$

Applying the basic definition of logarithms, we see that these equations would correspond to

$$\log_{10} A = m$$

$$\log_{10} B = n$$

$$\log_{10} AB = m + n$$

Therefore, $\log_{10} AB = \log_{10} A + \log_{10} B$

EXAMPLE: Multiply 100 times 1,000 using logarithms.

SOLUTION:

If $100 = 10^2$, then $\log_{10} 100 = 2$.

If $1{,}000 = 10^3$, then $\log_{10} 1{,}000 = 3$

So if

$$\log_{10}(100 \times 1{,}000) = \log_{10} 100 + \log_{10} 1{,}000$$

$$= 2 + 3$$

$$= 5$$

then the corresponding exponential form is

$$100 \times 1{,}000 = 10^5$$

$$= 100{,}000$$

DIVISION

Law 2. *The logarithm of a quotient is equal to the differenc*
of the logarithms of the dividend and the divisor

If A is to be divided by B, and

$$A = 10^m$$

and

$$B = 10^n$$

then

$$\frac{A}{B} = \frac{10^m}{10^n} = 10^{m-n}$$

In logarithmic form,

$$\log_{10} A = m$$

$$\log_{10} B = n$$

$$\log_{10} \frac{A}{B} = m - n$$

Therefore, $\log_{10} \frac{A}{B} = \log_{10} A - \log_{10} B$

EXAMPLE: Divide 1,000 by 100 using logarithms.

SOLUTION:

$$\log_{10} 1{,}000 = 3$$

$$\log_{10} 100 = 2$$

In logarithmic form,

$$\log_{10} \frac{1{,}000}{100} = \log_{10} 1{,}000 - \log_{10} 100$$

$$= 3 - 2$$

$$= 1$$

Therefore,

$$\frac{1{,}000}{100} = 10^1$$

$$= 10$$

POWERS

Law 3. *The logarithm of a number raised to a power is equal to the exponent times the logarithm of the number.*

Suppose

$$A = 10^m$$

and

$$A^n = (10^m)^n = 10^{mn}$$

Then

$$\log_{10} A = m$$

and

$$\log_{10} A^n = mn$$

Therefore, $\log_{10} A^n = (\log_{10} A)n = n \log_{10} A$

EXAMPLE: Find the value of 100^2 using logarithms.

SOLUTION: In logarithmic form,

$$\log_{10} 100^2 = 2 \log_{10} 100$$

$$= (2)(2)$$

$$= 4$$

Therefore,

$$100^2 = 10^4$$

$$= 10{,}000$$

ROOTS

Law 4. *The logarithm of the nth root of a number is equal to the logarithm of the number divided by n, the index of the root.*

Suppose

and

$$\sqrt[n]{A} = \sqrt[n]{10^m} = 10^{m/n}$$

Then

$$\log_{10} A = m$$

and

$$\log_{10}\sqrt[n]{A} = \frac{m}{n}$$

Therefore, $\log_{10}\sqrt[n]{A} = \dfrac{\log_{10} A}{n} = \dfrac{1}{n}\log_{10} A$

EXAMPLE: Find $\sqrt{100}$ using logarithms.

SOLUTION:

$$\sqrt{100} = 100^{1/2}$$

In logarithmic form,

$$\log_{10} 100^{1/2} = \frac{1}{2}\log_{10} 100$$

$$= \frac{1}{2}(2)$$

$$= 1$$

In exponential form,

$$100^{1/2} = 10^1$$

Therefore,

$$\sqrt{100} = 10$$

PRACTICE PROBLEMS:

Find the values of the following using the Laws for Logarithms:

1. $10{,}000 \times 1{,}000 \times 10$

2. 10,000/10

3. $1{,}000^3$

4. $\sqrt[3]{1{,}000}$

ANSWERS:

1. 100,000,000

2. 1,000

3. 1,000,000,000

4. 10

COMMON LOGARITHMS

We could construct tables of logarithms using any number as a base. For purposes of calculation, the most logical number for a base is 10, the base of the decimal number system. Logarithms to the base 10 are called *common logarithms*. Therefore, in the discussion which follows, no base designation is used. *The expression* log *A is understood to mean the base 10 logarithm of A or* $\log_{10} A$.

Most of the numbers encountered in various calculations are not integral (whole number) powers of 10.

EXAMPLE: Express 316 as a base 10 logarithm.

SOLUTION:

$$316 = 10^{2.4997}$$

Therefore, in logarithmic form

Every logarithm consists of an integral part, the *characteristic*, and a fractional part, the *mantissa*. The logarithm of 316 is

$$2.4997 = 2 + .4997$$

so the characteristic is 2 and the mantissa is .4997.

POSITIVE CHARACTERISTICS

The characteristic for the logarithm of an integer may be determined by inspection. For example, if the integer is between 1 and 10, it is equal to a power of 10 between 0 and 1. This concept is explained fully in *Mathematics*, Volume 1.

The numbers in the following list serve to illustrate how the characteristic is determined by the size of the number:

$$\log 3.6 = 0.5563$$

$$\log 36 = 1.5563$$

$$\log 360 = 2.5563$$

$$\log 3{,}600 = 3.5563$$

Since log 1 is 0 and log 10 is 1, we expect the logarithm of 3.6 to be a number between 0 and 1. Therefore, its characteristic is 0. On the other hand, 3,600 is greater than 1,000 and less than 10,000. Therefore, its logarithm is between log 1,000 and log 10,000, and its characteristic is 3.

Scientific notation provides a convenient method for determining the characteristic. For example, 3,600 is written as 3.6×10^3 in scientific notation. Thus, we have

$$\begin{aligned} \log 3{,}600 &= \log (3.6 \times 10^3) \\ &= \log 3.6 + \log 10^3 \\ &= 0.5563 + 3 \\ &= 3.5563 \end{aligned}$$

The characteristic of log 3.6 is 0, and the characteristic of log 10^3 is 3. Therefore, the characteristic of log 3,600 is 3, the sum of the characteristics of the two separate logarithms. *Any expression written in scientific notation consists of a number between 1 and 10 multiplied by a power of 10. Since the characteristic of a*

number between 1 and 10 is 0, the power of 10 determines the characteristic of the logarithm.

The exponent that we obtain as the power of 10 in scientific notation is indicated by the number of digits between the actual position of the decimal point in the original number and the standard position of the decimal point. The standard position is immediately after the first nonzero digit in the number. For example, in the number 3,600, the decimal point is understood to be after the second 0 in the original number. This is 3 digits to the right of the standard position, so the exponent of 10 for scientific notation is 3. This exponent is also the characteristic for log 3,600.

NEGATIVE CHARACTERISTICS

If the decimal point in the original number had been to the left of the standard position, the exponent of 10 (and therefore the characteristic) would have been negative. *When the logarithm of a positive number less than 1 is obtained, a negative characteristic occurs.* For example,

$$\begin{aligned} \log 0.036 &= \log (3.6 \times 10^{-2}) \\ &= \log 3.6 + \log 10^{-2} \\ &= \log 3.6 + (-2) \\ &= 0.5563 - 2 \end{aligned}$$

Since logarithm tables do not list negative characteristics, we do not subtract the characteristic from the mantissa to obtain the final form of the logarithm. Perhaps the most universal form for negative characteristics is to use a positive integer minus 10 or an integral multiple of 10 as follows:

$$\log 0.036 = 8.5563 - 10$$

This form is numerically equal to

$$\log 0.036 = 0.5563 - 2$$

Negative numbers and 0 do not have logarithms. When logarithms are used in calculations involving negative numbers, we first determine the sign of the final answer. Next we compute the results as if all the numbers were positive, and then we apply the predetermined sign to the final answer.

PRACTICE PROBLEMS:

Determine the characteristics of the logarithm for each of the following numbers:

1. 32

2. 476

3. 0.25

4. 0.0074

ANSWERS:

1. 1

2. 2

3. −1 or 9 − 10

4. −3 or 7 − 10

LOGARITHM TABLES

Tables of logarithms normally contain only mantissas. Table 1 is an excerpt from appendix I, Common Logarithms of Numbers. Observe that the table of logarithms has headings consisting of the abbreviation "No." (representing Number) and the digits 0 through 9. *The first two digits of any number whose logarithm we seek are found in the No. column. The third digit*

Table 1-1.—Appendix I Excerpt, Common Logarithms of Numbers

No.	0	1	2	3	4	5	6	7	8	
1.0	.0000	.0043	.0086	.0128	.0170	.0212	.0253	.0294	.0334	.0
1.1	.0414	.0453	.0492	.0531	.0569	.0607	.0645	.0682	.0719	.0

is found as one of the column headings, 0 through 9. The mantissa for the logarithm of any three-digit number is found opposite the first two digits and below the third digit.

Steps in determining the logarithm of a three-digit number are as follows:

1. Determine the characteristic of the logarithm of the number.
2. Locate the first two digits of the number on the left side of the table.
3. Locate the third digit of the number at the top of the table.
4. Locate the mantissa corresponding to these values.
5. Determine the logarithm using the characteristic, found in step 1, and the mantissa, found in step 4.

EXAMPLE: Find the logarithm of 1.24 using table 1-1.

SOLUTION:

1. The number 1.24 is in standard form, so the characteristic is 0.
2. Locate 1.2 on the left side of the table.
3. Locate 4 at the top of the table.
4. The mantissa corresponding to the values is .0934.
5. Therefore, the logarithm of 1.24 is 0.0934.

EXAMPLE: Find the logarithm of 13.7.

SOLUTION:

1. The characteristic of 13.7 is 1.
2. Locate 1.3 on the left side of the table.
3. Locate 7 at the top of the table.
4. The mantissa corresponding to the values is .1367.
5. Therefore, the logarithm of 13.7 is 1.1367.

EXAMPLE: Find the logarithm of 0.0682 using appendix I.

SOLUTION:

1. The characteristic of 0.0682 is -2.
2. Locate 6.8 on the left side of the table.
3. Locate 2 at the top of the table.
4. The mantissa corresponding to the values is .8338.
5. Therefore, the logarithm of 0.0682 is 0.8338 $-$ 2 or 8.8338 $-$ 10.

PRACTICE PROBLEMS:

Use table 1-1 or appendix I to find the logarithms of the following numbers:

1. 118
2. 3,420
3. 14.6
4. 5.48

ANSWERS:

1. 2.0719
2. 3.5340
3. 1.1644
4. 0.7388

INTERPOLATION

Interpolation of a mantissa is the process of calculating the mantissa for the logarithm of a number having one more digit than the entries in the table. For example, to find the logarithm of 1,125, we interpolate. Refer to table 1-1 or appendix I. The characteristic of 1,125 is 3, since $1{,}125 = 1.125 \times 10^3$. The logarithm of 1.125 is halfway between the logarithms of 1.120 and 1.130. Therefore, we find the mantissas for the logarithms of these two numbers and then determine the mantissa that is halfway between them. The interpolation for the number 1.125 can be performed as follows:

	NUMBER	MANTISSA		
.010	.005	1.120	.0492	.0039
		1.125	?	
		1.130	.0531	

x

We analyze the foregoing tabulation in terms of the difference between the numbers and the difference between the mantissas. The large bracket on the number side indicates a total difference of .010, and the small bracket on the number side indicates a total difference of .005. The large bracket on the mantissa side indicates a total difference of .0039. Since our number (1.125) is .005/.010, or 5/10 of the way between the two numbers (1.120 and 1.130) in the table, then the mantissa corresponding to our number should be 5/10 of the way between the mantissas (.0492 and .0531) in the table.

Writing the proportions, we have

$$\frac{5}{10} = \frac{x}{.0039}$$

Solving for x gives

$$x = \frac{5}{10}(.0039)$$

$$= .00195$$

$$= .0020 \text{ (rounded to 4 places)}$$

Adding .0020 to .0492, we obtain the mantissa corresponding to 1.125, which is .0512. Therefore,

$$\log 1{,}125 = 3.0512$$

Steps in determining the logarithm of a number by interpolation are as follows:

1. Determine the characteristic of the logarithm of the number.
2. Determine the numbers your number (in standard form) lies between and their corresponding mantissas.
3. Interpolate to obtain the mantissa for your number.
4. Determine the logarithm using the characteristic, found in step 1, and the mantissa, found in step 3.

EXAMPLE: Find log 25.67.

SOLUTION:

1. The characteristic of 25.67 is 1

2. The number 2.567 lies between 2.560 and 2.570; and their corresponding mantissas are .4082 and .4099, respectively.
3. Interpolate:

		NUMBER	MANTISSA		
.010	.007	2.560	.4082	x	.0017
		2.567	?		
		2.570	.4099		

Our number is .007/.010, or 7/10 of the way between 2.560 and 2.570. Therefore, the mantissa must also be 7/10 of the way between .4082 and .4099.

Writing the proportions, we have

$$\frac{7}{10} = \frac{x}{.0017}$$

Solving for x gives

$$x = \frac{7}{10}(.0017)$$

$$= .00119$$

$$= .0012 \text{ (rounded to 4 places)}$$

So the mantissa corresponding to 2.567 is

$$.4082 + .0012 = .4094$$

4. Therefore, log 25.67 = 1.4094.

PRACTICE PROBLEMS:

Find the logarithms of the following numbers:

1. 0.2355

2. 5.432

3. 473.6

4. 9,817

ANSWERS:

1. 9.3720 − 10

2. 0.7350

3. 2.6754

4. 3.9920

ANTILOGARITHMS

The procedure for finding a number when we know its logarithm is called finding the *antilogarithm*. The word "antilogarithm" is abbreviated "antilog," and a symbol sometimes used to indicate the antilog is $\log^{-1}$. The -1 in a symbol of this kind tends to be confusing since it is not an exponent. It is an indicator that emphasizes the inverse relationship between logs and antilogs.

Steps in determining the antilogarithm of an exact table entry of a mantissa are as follows:

1. Locate the mantissa of the logarithm of the number in the table.
2. Locate the two-digit number to the left of the mantissa.
3. Locate the one-digit number directly above the mantissa.
4. Combine the two-digit number and the one-digit number to obtain the three-digit number corresponding to the mantissa.
5. Determine the antilogarithm using the three-digit number, found in step 4, and place the decimal either to the left or right using the characteristic of the original logarithm.

EXAMPLE: Find antilog 1.1271.

SOLUTION:

1. The mantissa of 1.1271 is .1271.
2. The two-digit number to the left of the mantissa is 1.3.
3. The one-digit number directly above the mantissa is 4.
4. The three-digit number corresponding to the mantissa is, then, 1.34.
5. Therefore, antilog 1.1271 = 13.4.

Steps in determining the antilogarithm of a mantissa that is not an exact table entry are as follows:

1. Determine the mantissas your mantissa lies between and their corresponding three-digit numbers.
2. Interpolate to obtain a four-digit number corresponding to your mantissa.
3. Determine the antilogarithm using the four-digit number, found in step 2, and place the decimal either to the left or right using the characteristic of the original logarithm.

EXAMPLE: Find the antilogarithm of 8.5124 − 10.

SOLUTION:

1. The mantissa .5124 lies between the mantissas .5119 and .5132; their corresponding three-digit numbers are 3.25 and 3.26, respectively.
2. Interpolate:

		NUMBER	MANTISSA		
.01	x	3.25	.5119	.0005	.0013
		?	.5124		
		3.26	.5132		

Our mantissa is .0005/.0013, or 5/13 of the way between .5119 and .5132. Therefore, our number is 5/13 of the way between 3.25 and 3.26.

Writing the proportions, we have

$$\frac{x}{.01} = \frac{5}{13}$$

Solving for x gives

$$x = \frac{5}{13}(.01)$$

$$= .0038$$

$$= .004 \text{ (rounded to 3 places)}$$

So the number corresponding to the mantissa .5124 is

$$3.25 + .004 = 3.254$$

3. Therefore, the antilogarithm of $8.5124 - 10$ is 0.03254.

PRACTICE PROBLEMS:

Find the antilogarithm of the following logarithms:

1. $9.3636 - 10$
2. 1.8451
3. 2.7030
4. 0.3842

ANSWERS:

1. 0.231
2. 70.0
3. 504.7
4. 2.422

NATURAL LOGARITHMS

Natural logarithms are so named because the number e, the base of the natural logarithm system, is involved in the law of nature that governs growth and decay. The law is stated in symbols as

$$A = A_o e^{rt}$$

In the foregoing equation, A represents the total amount after a period of growth, and A_o represents the amount at the beginning of the growth period. The letter r represents the continuous rate of growth, and t represents the time during which growth occurs. The same remarks apply for a period of decay.

By means of higher mathematics, *the number e is found to have the value*

$$e = 2.71828 \text{ (rounded to 5 places)}$$

This number is the base of the natural logarithm system.

The relationship between the common logarithm of a number and its natural logarithm is

$$\ln N = 2.3026 \log N$$

Observe that the special abbreviation ln N is used to represent $\log_e N$.

If $e^x = N$, where N is any number, then by taking the natural logarithm of both sides, we have

$$x \ln e = \ln N$$

Since

$$\ln e = \log_e e$$

then

$$\ln e = 1$$

Therefore, by substitution,

$$x = \ln N$$

and

$$\frac{\ln N}{1} = N$$

The value of $\ln N$ can also be obtained from the basic definition of logarithms. Taking common logarithms of both sides in the expression

$$e^x = N$$

gives

$$\log e^x = \log N$$

$$x \log e = \log N$$

$$x = \frac{\log N}{\log e}$$

Equating the two expressions we have obtained for x gives

$$\ln N = \frac{\log N}{\log e}$$

From the table of common logarithms, we find that log 2.71828 is approximately 0.4343, so

$$\ln N = \frac{\log N}{0.4343}$$

Since the reciprocal of 0.4343 is 2.3026 (rounded), then

$$\ln N = 2.3026 \log N$$

EXAMPLE: Find the natural logarithm of 36.

SOLUTION:

$$\ln 36 = 2.3026 \log 36$$

$$= 2.3026(1.5563)$$

$$= 3.5835$$

EXAMPLE: Find ln 0.053.

SOLUTION:

$$\ln 0.053 = 2.3026 \log 0.053$$

$$= 2.3026(8.7243 - 10)$$

Transform $8.7243 - 10$ to the equivalent form -1.2757 to multiply as follows:

$$\ln 0.053 = 2.3026(-1.2757)$$

$$= -2.9374$$

Now write -2.9374 in the universal form for negative characteristics:

$$-2.9374 = 7.0626 - 10$$

Therefore,

$$\ln 0.053 = 7.0626 - 10$$

PRACTICE PROBLEMS:

Find the natural logarithm of the following numbers:

1. 15
2. 8,014
3. 29
4. 352

ANSWERS:

1. 2.7081
2. 8.9889
3. 3.3673
4. 5.8636

SUMMARY

The following are the major topics covered in this chapter:

1. **Definition of a logarithm:** Every logarithm is an exponent.

   If b is a positive number other than 1 and x is a real number (positive, negative, or zero) in the equation $b^x = a$, then $x = \log_b a$.

   The *exponential form* is $b^x = a$ and the *logarithmic form* is $x = \log_b a$.

2. **Two abilities necessary for logarithmic calculations:**

   1. Recognition of logarithms as exponents
   2. Knowledge of the Laws for Logarithms

3. **Laws for Logarithms:**

   Law 1. *The logarithm of a product is equal to the sum of the logarithms of the factors.*

   $$\log_{10} AB = \log_{10} A + \log_{10} B$$

   Law 2. *The logarithm of a quotient is equal to the difference of the logarithms of the dividend and the divisor.*

   $$\log_{10} \frac{A}{B} = \log_{10} A - \log_{10} B$$

   Law 3. *The logarithm of a number raised to a power is equal to the exponent times the logarithm of the number.*

   $$\log_{10} A^n = (\log_{10} A)n = n \log_{10} A$$

   Law 4. *The logarithm of the nth root of a number is equal to the logarithm of the number divided by n, the index of the root.*

   $$\log_{10} \sqrt[n]{A} = \frac{\log_{10} A}{n} = \frac{1}{n} \log_{10} A$$

4. **Common logarithms:** Logarithms to the base 10 are called *common logarithms.*

   The expression log A is understood to mean the base 10 logarithm of A or $\log_{10} A$.

Every logarithm consists of an integral part, the *characteristic*, and a fractional part, the *mantissa*.

Scientific notation provides a convenient method for determining the characteristic. Any expression written in scientific notation consists of a number between 1 and 10 multiplied by a power of 10. Since the characteristic of a number between 1 and 10 is 0, the power of 10 determines the characteristic of the logarithm.

When the logarithm of a positive number less than 1 is obtained, a negative characteristic occurs. Since logarithm tables do not list negative characteristics, the characteristic is not subtracted from the mantissa to obtain the final form of the logarithm. The most universal form for negative characteristics is to use a positive integer minus 10 or an integral multiple of 10.

Negative numbers and 0 do not have logarithms. When logarithms are used in calculations involving negative numbers, first determine the sign of the final answer. Next compute the results as if all the numbers were positive, and then apply the predetermined sign to the final answer.

5. **Common Logarithm Tables:** The first two digits of any number whose logarithm we seek are found in the No. column. The third digit is found as one of the column headings, 0 through 9. The mantissa for the logarithm of any three-digit number is found opposite the first two digits and below the third digit.

6. **Steps in determining the logarithm of a three-digit number:**

    1. Determine the characteristic of the logarithm of the number.
    2. Locate the first two digits of the number on the left side of the table.
    3. Locate the third digit of the number at the top of the table.
    4. Locate the mantissa corresponding to these values.
    5. Determine the logarithm using the characteristic, found in step 1, and the mantissa, found in step 4.

7. **Interpolation of a mantissa:** *Interpolation of a mantissa* is the process of calculating the mantissa for the logarithm of a

8. **Steps in determining the logarithm of a number by interpolation:**

    1. Determine the characteristic of the logarithm of the number.
    2. Determine the numbers your number (in standard form) lies between and their corresponding mantissas.
    3. Interpolate to obtain the mantissa for your number.
    4. Determine the logarithm using the characteristic, found in step 1, and the mantissa, found in step 3.

9. **Antilogarithms**: The procedure for finding a number when we know its logarithm is called finding the *antilogarithm*. The word "antilogarithm" is abbreviated "antilog," and a symbol sometimes used to indicate the antilog is $\log^{-1}$.

10. **Steps in determining the antilogarithm of an exact table entry of a mantissa:**

    1. Locate the mantissa of the logarithm of the number in the table.
    2. Locate the two-digit number to the left of the mantissa.
    3. Locate the one-digit number directly above the mantissa.
    4. Combine the two-digit number and the one-digit number to obtain the three-digit number corresponding to the mantissa.
    5. Determine the antilogarithm using the three-digit number, found in step 4, and place the decimal either to the left or right using the characteristic of the original logarithm.

11. **Steps in determining the antilogarithm of a mantissa that is not an exact table entry:**

    1. Determine the mantissas your mantissa lies between and their corresponding three-digit numbers.
    2. Interpolate to obtain a four-digit number corresponding to your mantissa.
    3. Determine the antilogarithm using the four-digit number, found in step 2, and place the decimal either to the left or right using the characteristic of the original logarithm.

12. **Natural logarithms:** *Natural logarithms* are so named because the number e, the base of the natural logarithm system, is involved in the law of nature that governs growth and decay. The law is stated in symbols as

$$A = A_0 e^{rt}$$

where A represents the total amount after a period of growth or decay, A_o represents the amount at the beginning of the growth or decay period, r represents the continuous rate of growth or decay, and t represents the time during which growth or decay occurs.

The number e (rounded to 5 places) is

$$e = 2.71828$$

The relationship between the common logarithm of a number and its natural logarithm is

$$\ln N = 2.3026 \log N$$

ADDITIONAL PRACTICE PROBLEMS

1. Change $3^{-2} = 1/9$ to logarithmic form.
2. Change $\log_{10}2 = 0.3010$ to exponential form.
3. Multiply $100^3 \times 10^{-4}$ using logarithms.
4. Divide $\sqrt{10{,}000}$ by 1,000 using logarithms.
5. Determine the characteristic of the logarithm of 89,000.
6. Find the logarithm of 0.00801.
7. Find the logarithm of 99,660.
8. Find the antilogarithm of $6.7404 - 10$.
9. Find the antilogarithm of 3.6060.
10. Find the natural logarithm of 0.00673.

ANSWERS TO ADDITIONAL PRACTICE PROBLEMS

1. $\log_3(1/9) = -2$
2. $10^{0.3010} = 2$
3. 100
4. 0.1
5. 4
6. 7.9036 − 10
7. 4.9985
8. 0.000550
9. 4,036
10. 4.9988 − 10

CHAPTER 2

COMPUTATIONS WITH LOGARITHMS

LEARNING OBJECTIVES

Upon completion of this chapter, you should be able to do the following:

1. Multiply and divide numbers using logarithms.
2. Compute the power of a number and the root of a number using logarithms.
3. Apply the laws for logarithms to algebraic operations and to problem solving.

INTRODUCTION

In this chapter additional mention of the Laws for Logarithms will be given followed by algebraic operations and applications using logarithms.

Laws for Powers and Roots are listed in table 2-1 for reference and review.

Table 2-1.—Laws For Powers and Roots

APPLICATION	LAW	EXAMPLE
Multiplication	$a^n a^m = a^{n+m}$	$3^4 \cdot 3^2 = 3^{4+2} = 3^6$
Division	$\frac{a^n}{a^m} = a^{n-m}$	$\frac{3^4}{3^2} = 3^{4-2} = 3^2$
Power Raised to a Power	$(a^n)^m = a^{nm}$	$(3^4)^2 = 3^{4 \cdot 2} = 3^8$
Product Raised to a Power	$(ab)^n = a^n b^n$	$(3 \cdot 5)^4 = 3^4 \cdot 5^4$
Quotient Raised to a Power	$\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n}$, if $b \neq 0$	$\left(\frac{3}{5}\right)^4 = \frac{3^4}{5^4}$
Negative Power	$a^{-n} = \frac{1}{a^n}$	$3^{-4} = \frac{1}{3^4}$
Square Root	$\sqrt{a} = a^{1/2}$	$\sqrt{3} = 3^{1/2}$
nth Root	$\sqrt[n]{a} = a^{1/n}$	$\sqrt[4]{3} = 3^{1/4}$
nth Root of a Power	$\sqrt[n]{a^m} = a^{m/n}$	$\sqrt[4]{3^2} = 3^{2/4}$

All calculations by means of logarithms in this chapter use 10 as the base. By the convention established in chapter 1, the expression $\log A$ is understood to mean the base 10 logarithm of A.

Computations using logarithms will be evaluated to four significant digits to correspond with interpolation procedures introduced in chapter 1. All the digits of an approximate number except zeros, which serve only to fix the position of the decimal point, are called *significant digits.*

MULTIPLICATION

Law 1. *The logarithm of a product is equal to the sum of the logarithms of the factors;* that is,

$$\log AB = \log A + \log B$$

EXAMPLE: Use logarithms to find the product of 386×254 to four significant digits.

SOLUTION:

Exponential solution:

$$386 = 10^{2.5866}$$

$$254 = 10^{2.4048}$$

$$386 \times 254 = 10^{2.5866} \times 10^{2.4048}$$

$$= 10^{(2.5866 + 2.4048)}$$

$$= 10^{4.9914}$$

Logarithmic solution:

$$\log 386 = 2.5866$$

$$\log 254 = 2.4048$$

$$\log (386 \times 254) = \log 386 + \log 254$$

$$= 2.5866 + 2.4048$$

To find the antilogarithm of 4.9914, we will interpolate:

	NUMBER	MANTISSA	
	9.80	.9912	
x			.0002
.01	?	.9914	.0005
	9.81	.9917	

$$x = \frac{.0002}{.0005}(.01) = \frac{2}{5}(.01) = .004$$

Therefore, antilog 4.9914 = 98,040, which is very close to the actual calculated value of 98,044.

The exponential solution shown in the example is not a part of normal calculations involving logarithms. It was shown in this first example problem solely for the purpose of reemphasizing the relationship between exponents and logarithms.

NOTE: The logarithmic value of 98,040 is not the same as the actual value of 98,044, because the logarithmic table values (used in this book) are only significant to four digits. The larger the table values, the closer the logarithmic value is to the actual value.

EXAMPLE: Use logarithms to find the product of (126)(−33) to four significant digits.

SOLUTION: Recall from chapter 1 that negative numbers do not have logarithms. In using logarithms to solve problems that involve negative numbers, we first determine the sign of the final answer. After this sign is determined, calculations are performed as if all numbers are positive, and then the predetermined sign is applied to the answer.

In our example, dealing first with signs only, we determine the answer to be negative; that is, (+)(−) = (−). At this point the problem can be restated: Use logarithms to find the product of −(126 × 33).

$$\begin{aligned} \log (126 \times 33) &= \log 126 + \log 33 \\ &= 2.1004 + 1.5185 \\ &= 3.6189 \end{aligned}$$

and by interpolation,

$$\text{antilog } 3.6189 = 4{,}158$$

Therefore, $(126)(-33) = -4{,}158$. This value is the same as the actual value.

EXAMPLE: Use logarithms to find the product of $1.73 \times 0.0024 \times 0.08$ to four significant digits.

SOLUTION:

$$\begin{aligned} &\log (1.73 \times 0.0024 \times 0.08) \\ &= \log 1.73 + \log 0.0024 + \log 0.08 \\ &= 0.2380 + (7.3802 - 10) + (8.9031 - 10) \\ &= 16.5213 - 20 \\ &= 6.5213 - 10 \end{aligned}$$

To find antilog $(6.5213 - 10)$, we will interpolate:

	NUMBER	MANTISSA	
	3.32	.5211	
x			.0002
.01	?	.5213	.0013
	3.33	.5224	

$$x = \frac{.0002}{.0013}(.01) = \frac{2}{13}(.01) = .002$$

So, antilog $(6.5213 - 10) = 0.0003322$. The logarithmic value is again the same as the actual value to four significant digits.

PRACTICE PROBLEMS:

Use logarithms to find the product of the following to four significant digits:

1. $(53)(-76)(-0.021)(153)$

3. $(0.00432)(-0.00106)(15)$

4. $0.102 \times 103.5 \times 76.2$

ANSWERS:

1. 12,940

2. 4,856,000

3. -6.869×10^{-5}

4. 804.4

DIVISION

Law 2. *The logarithm of a quotient is equal to the difference of the logarithms of the dividend and the divisor;* that is,

$$\log \frac{A}{B} = \log A - \log B$$

EXAMPLE: Find the quotient of 37.4/1.7 by use of logarithms to four significant digits.

SOLUTION:

$$\log (37.4/1.7) = \log 37.4 - \log 1.7$$

$$= 1.5729 - 0.2304$$

$$= 1.3425$$

and

$$\text{antilog } 1.3425 = 22.01$$

and 22.00 is the actual value.

EXAMPLE: Use logarithms to find the quotient of 16.3/0.008

SOLUTION:

$$\log (16.3/0.008) = \log 16.3 - \log 0.008$$

$$= 1.2122 - (7.9031 - 10)$$

To prevent the complication of subtracting a larger characteristic (7) from a smaller characteristic (1), we add 10 to and subtract 10 from the logarithm of the dividend. Note that this does not change the value of the logarithm. Thus,

$$\begin{aligned} \log 16.3 &= 11.2122 - 10 \\ -\log 0.008 &= \underline{7.9031 - 10} \\ &\quad 3.3091 \end{aligned}$$

and

$$\text{antilog } 3.3091 = 2{,}038$$

Therefore, the logarithmic value of 16.3/0.008 is 2,038, while the actual value is 2,037.5.

PRACTICE PROBLEMS:

Use logarithms to solve the following problems to four significant digits:

1. 635.6/25.4

2. 0.26/0.061

3. 0.126/0.00542

4. 874/26.3

ANSWERS:

1. 25.03

3. 23.25

4. 33.23

POWERS

Law 3. *The logarithm of a number raised to a power is equal to the exponent times the logarithm of the number;* that is,

$$\log A^n = n \log A$$

EXAMPLE: Use logarithms to find the value of $(18.53)^5$ to four significant digits.

SOLUTION:

$$\begin{aligned} \log (18.53)^5 &= 5 \log 18.53 \\ &= 5(1.2679) \\ &= 6.3395 \end{aligned}$$

and

$$\text{antilog } 6.3395 = 2{,}185{,}000$$

So the logarithmic value of $(18.53)^5$ is 2,185,000, while the actual value is 2,184,626.

ROOTS

Law 4. *The logarithm of the nth root of a number is equal to the logarithm of the number divided by n, the index of the root;* that is,

$$\log \sqrt[n]{A} = \frac{1}{n} \log A$$

EXAMPLE: Use logarithms to find the value of $\sqrt[5]{327.6}$ to

SOLUTION:

$$\log \sqrt[5]{327.6} = \frac{1}{5} \log 327.6$$

$$= \frac{1}{5}(2.5153)$$

$$= 0.5031$$

where

$$\text{antilog } 0.5031 = 3.185$$

which is the logarithmic value and the actual of $\sqrt[5]{327.6}$ to four significant digits.

When a logarithm with a negative characteristic is to be divided, adding and subtracting a number that will, after dividing, leave a minus 10 at the right is advisable. This is done to keep the logarithm in standard form.

EXAMPLE: Find the value of $\sqrt[5]{0.0018}$ to four significant digits using logarithms.

SOLUTION:

$$\log \sqrt[5]{0.0018} = \frac{1}{5} \log 0.0018$$

$$= \frac{1}{5}(7.2553 - 10)$$

To keep a minus 10 in the final logarithm, we must add and subtract 40 before dividing. Thus,

$$\log \sqrt[5]{0.0018} = \frac{1}{5}(47.2553 - 50)$$

$$= 9.4511 - 10$$

where

$$\text{antilog } (9.4511 - 10) = 0.2826$$

Therefore, the logarithmic value of $\sqrt[5]{0.0018}$ is 0.2826, while the

PRACTICE PROBLEMS:

Evaluate the following to four significant digits using logarithms:

1. $(3.276)^3$

2. $(0.00468)^2$

3. $\sqrt[6]{0.00867}$

4. $\sqrt[5]{237.7}$

ANSWERS:

1. 35.15

2. 0.00002190

3. 0.4532

4. 2.987

ALGEBRAIC OPERATIONS

This chapter has demonstrated the use of logarithms in nerical calculations. *Practical applications in many fields olve calculations including algebraic expressions in which arithms are useful. In these problems both the laws for algebra l the laws for logarithms in algebraic operations are valid.* For mple,

$$\log (x + 2)(x + 5) = \log (x + 2) + \log (x + 5)$$

EXAMPLE: Solve for x in the equation

$$\log (x^2 - 5x - 6) - \log (x + 1) = 1$$

TE: $\log 10 = 1$.

SOLUTION:

$$\log \frac{x^2 - 5x - 6}{x + 1} = \log 10$$

$$\log \frac{(x - 6)(x + 1)}{(x + 1)} = \log 10$$

$$\log (x - 6) = \log 10$$

$$x - 6 = 10$$

$$x = 16$$

EXAMPLE: Solve for x to four significant digits using logarithms:

$$26^x = 195$$

SOLUTION: Take the logarithm of both sides of the equation:

$$\log 26^x = \log 195$$

$$x \log 26 = \log 195$$

$$x = \frac{\log 195}{\log 26}$$

$$= \frac{2.2900}{1.4150}$$

$$= 1.618$$

In complicated problems we may not be able to solve for the unknown as directly as we did in the previous example. In that case we can continue to use our knowledge of logarithms. For instance, return to the step where $x = 2.29/1.415$; take the logarithm of both sides of the equation

$$\log x = \log \frac{2.29}{1.415}$$

$$= \log 2.29 - \log 1.415$$

$$= 0.3598 - 0.1508$$

$$= 0.2090$$

Now take the antilogarithm of both sides:

$$x = \text{antilog } 0.2090$$

$$= 1.618$$

The logarithmic and actual values of x are both 1.618 to four significant digits, so $26^{1.618} \approx 195$.

EXAMPLE: Solve for x using logarithms:

$$x^{3/2} = 729$$

SOLUTION:

$$\log x^{3/2} = \log 729$$

$$(3/2) \log x = \log 729$$

$$\log x = (2/3) \log 729$$

$$= (2/3)(2.8627)$$

$$= 1.9085$$

$$x = \text{antilog } 1.9085$$

$$= 81$$

The logarithmic and actual values are both 81, so $81^{3/2} = 729$.

PRACTICE PROBLEMS:

Use logarithms to solve for x to four significant digits in the following:

1. $1.7^x = 3.1$

2. $x^{8/3} = 6.35$

ANSWERS:

1. 2.133

2. 2.000

APPLICATIONS

The use of logarithms can simplify the solution of many problems encountered in mathematics, science, and engineering. Applying the operations described in this chapter can reduce many complicated equations to addition and subtraction problems.

EXAMPLE: Find the volume of a circular cone having a height of 3.71 inches and a base radius of 2.71 inches.

SOLUTION: The formula for volume of a circular cone is

$$v = \frac{\pi r^2 h}{3}$$

where v is volume, r is radius, h is height, and π (pi) is equal to 3.142 to four significant digits.

Take the logarithm of both sides of the equation as the first step in the solution and continue with the Laws for Logarithms.

$$\log v = \log \left(\frac{\pi r^2 h}{3}\right)$$

$$= \log \pi + \log r^2 + \log h - \log 3$$

$$= \log (3.142) + 2 \log (2.71) + \log (3.71) - \log (3)$$

$$= 0.4972 + 2(0.4330) + 0.5694 - 0.4771$$

$$= 1.4555$$

$$v = \text{antilog } 1.4555$$

$$= 28.54$$

The volume of the circular cone using logarithms to four significant digits is 28.54 inches cubed, while the actual value to four significant digits is 28.53 inches cubed.

Many electronics problems can be simplified by using logarithms. Some electronics problems include common logarithms in the basic formulas. An example of a formula that includes a logarithmic expression is the formula for finding gain in decibels, where

$$\text{decibels} = 10 \log \frac{P_1}{P_2}$$

Engineering and electronics problems frequently deal with numbers in the millions and decimal fractions in the millionths. These values are easily expressed as exponentials to the base 10, and common logarithms are then a natural and convenient means of simplifying these problems.

EXAMPLE: $X_C = \frac{1}{2\pi fC}$ is a formula used to analyze alternating current circuits. Use logarithms to find X_C to four significant digits, if $f = 22{,}000{,}000$ Hertz and $C = 1.5 \times 10^{-9}$ farads.

SOLUTION: Taking the logarithm of both sides of the formula $X_C = \frac{1}{2\pi fC}$ gives

$$\log X_C = \log 1 - \log (2\pi fC)$$

$$= 0 - [\log 2 + \log 3.142 + \log (2.2 \times 10^7) + \log (1.5 \times 10^{-9})]$$

$$= - [0.3010 + 0.4972 + 7.3424 + (1.1761 - 10)]$$

$$= - (9.3167 - 10)$$

$$= - 9.3167 + 10$$

$$= 0.6833$$

$$X_C = \text{antilog } 0.6833$$

$$= 4.823 \text{ ohms}$$

PRACTICE PROBLEMS:

Use logarithms to solve for the numerical value of the unknown in the following problems to four significant digits:

1. Find the volume (v) of a sphere, given the formula

$$v = \frac{4\pi r^3}{3}$$

where the radius (r) is 7.59 and π is 3.142

2. Find the value of I in the formula

$$P = I^2R$$

when $P = 217$ and $R = 550{,}000$.

ANSWERS:

1. 1,831

2. 0.01986

SUMMARY

The following are the major topics covered in this chapter:

1. **Significant digits:** All the digits of an approximate number except zeros, which serve only to fix the position of the decimal point, are called *significant digits.*

2. **Multiplication:**

   Law 1. *The logarithm of a product is equal to the sum of the logarithms of the factors.*

3. **Division:**

   Law 2. *The logarithm of a quotient is equal to the difference of the logarithms of the dividend and the divisor.*

   To prevent the complication of subtracting a larger characteristic from a smaller characteristic, add 10 to and subtract 10 from the logarithm of the dividend.

4. **Powers:**

   Law 3. *The logarithm of a number raised to a power is equal to the exponent times the logarithm of the number.*

5. **Roots:**

   Law 4. *The logarithm of the nth root of a number is equal to the logarithm of the number divided by n, the index of the root.*

   When a logarithm with a negative characteristic is to be divided, add and subtract a number that will, after dividing, leave a minus 10 at the right. This is done to keep the logarithm in standard form.

6. **Algebraic operations:** Practical applications in many fields involve calculations in which logarithms are useful. In these problems both the laws for algebra and the laws for logarithms in algebraic operations are valid.

7. **Applications:** The use of logarithms can simplify the solution of many problems encountered in mathematics, science, and engineering. Applying operations can reduce many complicated

ADDITIONAL PRACTICE PROBLEMS

Use logarithms to solve the following problems to four significant digits:

1. $\dfrac{(-46.3)(189)}{(-2.13)}$

2. $\dfrac{(815)}{(7.95)^4}$

3. $(-2.46)^3\ (1.11)^5$

4. $\sqrt[5]{\dfrac{(49.9)(5.00)}{(0.0348)}}$

5. Solve for x:

$$5 \cdot 4^x = 6 \cdot 3^x$$

HINT: $\dfrac{4^x}{3^x} = \left(\dfrac{4}{3}\right)^x$

6. $H.P. = \dfrac{1.28apv^2}{1{,}100}$ is a formula used in aeronautics.

Find $H.P.$ when $p = 0.003$, $a = 5.5$ and $v = 253.7$.

7. The chemist defines the pH (hydrogen potential) of a solution by

$$pH = \log \frac{1}{[H^+]}$$

where $[H^+]$ is a numerical value for the concentration of hydrogen ions in an aqueous solution in moles per liter. Calculate the pH of a solution whose hydrogen ion concentration is 3.7×10^{-6}.

ANSWERS TO ADDITIONAL PRACTICE PROBLEMS

1. 4,109
2. 0.2040
3. -25.07
4. 5.903
5. 0.6336
6. 1.236
7. 5.4318

CHAPTER 3

TRIGONOMETRIC MEASUREMENTS

LEARNING OBJECTIVES

Upon completion of this chapter, you should be able to do the following:

1. Measure angles in degrees, radians, and mils.
2. Find angular velocity and the area of a sector using radians.
3. Apply the Pythagorean theorem and properties of similar right triangles to problem solving.
4. Apply trigonometric ratios, functions, and tables to problem solving.

INTRODUCTION

This is the first of several chapters in this course dealing with the subject of trigonometry. Chapters 4, 5, and 6 also deal directly with triangles and trigonometry. Chapter 7 deals with vectors and forces. The study of vectors and forces is so closely related to trigonometry that it is normally included in a trigonometry course.

Mathematics, Volume 1, introduces numerical trigonometry and some applications in problem solving. However, trigonometry is not restricted to solving problems involving triangles; it also forms a foundation for some advanced mathematical concepts and subject areas. Trigonometry is both algebraic and geometric in nature, and in this course both of these qualities will be applied.

MEASURING ANGLES

Mathematics, Volume 1, pointed out that an angle is formed when two straight lines intersect. In this course, an angle is considered to be generated when a line having a set direction is rotated about a point, as depicted in figure 3-1.



Figure 3-1.—Generat

In figure 3-1, line OA is laid out as a reference line having set direction. One end of the line is used as a pivot point and e line is rotated from its initial position (line OA) to another osition (line OB), as in opening a door. As the line turns on its vot point, it generates the angle AOB.

The following terminology is used in this and subsequent apters:

1. *Radius vector*—The line that is rotated to generate an angle.
2. *Initial position*—The original position of the radius vector; corresponds to line OA in figure 3-1.
3. *Terminal position*—The final position of the radius vector; corresponds to line OB in figure 3-1.
4. *Positive angle*—The angle generated by rotating the radius vector counterclockwise from the initial position.
5. *Negative angle*—The angle generated by rotating the radius vector clockwise from the initial position.

The convention of identifying angles by use of Greek letters followed in this text. When only one angle is involved, it will e symbolized by θ (theta). Other Greek letters will be used when ore than one angle is involved. The additional symbols used will e ϕ (phi), α (alpha), and β (beta).

EGREES

The degree system is the most common system of angular easurement. In this system a complete revolution is divided into 60 equal parts called *degrees*; so,

$$1 \text{ revolution} = 360°$$

or accuracy, each degree is divided into 60 minutes; o,

$$1° = 60'$$

ach minute is divided into 60 seconds; so,

$$1' = 60''$$

For convenience in working with angles, the 360° re divided into four equal parts of 90° each, similar o the rectangular coordinate system. The 90° sectors, alled *quadrants*, are numbered according to the onvention shown in figure 3-2.



Figure 3-2.—Quadrant positi

If the angle generated by rotating the radius vector in a positive (counterclockwise) direction is between 0° and 90°, then the angle is in the first quadrant. If the angle is between 90° and 180°, then the angle is in the second quadrant. If the angle is between 180° and 270°, then the angle is in the third quadrant. And if the angle is between 270° and 360°, then the angle is in the fourth quadrant.

If the angle generated by rotating the radius vector in a positive direction is more than 360°, then the quadrant in which the angle lies is found by subtracting from the angle the largest multiple of 360° that the angle contains. The quadrant in which the remainder angle lies is determined as described in the previous paragraph. *The original angle lies in the same quadrant as the remainder angle.*

EXAMPLE: In which quadrant is the angle 130°?

SOLUTION: Since 130° is between 90° and 180°, it is in the second quadrant. (See fig. 3-3, view A).

EXAMPLE: In which quadrant is the angle 850°?

SOLUTION: The largest multiple of 360° contained in 850° is 720°; so, 850° − 720° = 130°. Since 130° is in the second quadrant, then 850° also lies in the second quadrant. This relationship is shown in figure 3-3, view B.



Figure 3-3.—Angle generation.

PRACTICE PROBLEMS:

Determine the quadrant in which each of the following angles lies:

1. 260°

2. 290°

3. 800°

4. 1,930°

ANSWERS:

1. 3rd

2. 4th

3. 1st

4. 2nd

RADIANS

Another even more fundamental method of angular measurement involves the *radian*. It has certain advantages over the degree method. Radian measurement greatly simplifies work with trigonometric functions in calculus. Radian measurement also relates the length of arc generated to the size of an angle.

A *radian* is defined as an angle that, if its vertex is placed at the center of a circle, intercepts an arc equal in length to the radius vector of the circle. Assume that an angle is generated, as shown in figure 3-4, view A. If we impose the condition that the length of the arc, *s*, described by the extremity of the line segment generating the angle, must



Figure 3-4.—Radian measure.

equal the length of the radius vector, r, then we would describe an angle exactly one radian in size; that is, for 1 radian,

$$s = r$$

In a broader sense, *the radian measure of an angle, θ, is the ratio of the length of the arc, s, it subtends to the length of the radius vector, r, of the circle in which it is the central angle*; that is,

$$\theta = \frac{s}{r}$$

For angle θ, in figure 3-4, view B, which intercepts an arc equal to two times the length of the radius vector, θ equals two radians. For angle θ, in figure 3-4, view C, which intercepts an arc equal to three times the length of the radius vector, θ equal three radians.

EXAMPLE: Find the radian measure of the central angle in a circle with a radius of 10 inches if the angle subtends an arc of 5 inches.

SOLUTION:

$$\theta = \frac{s}{r}$$

$$= \frac{5}{10}$$

$$= 0.5 \text{ radians}$$

Recall from plane geometry that the circumference of a circle is 2π times the radius or

$$C = 2\pi r$$

Hence, the radius vector can be laid off on the circumference 2π times. (See fig. 3-4, view D).

Since the arc length of the circumference is 2π radians and the circumference encompasses one complete revolution of 360°, then

$$2\pi \text{ radians} = 360°$$

One-half of a revolution equals 180° or π radians; so,

$$\text{radians} = \frac{180°}{\pi} \qquad (3\text{-}1)$$

By dividing both sides of equation (3.1) by π, we find that

$$1 \text{ radian} = \frac{180^\circ}{\pi}$$

$$= 57.2958^\circ \text{ (rounded)}$$

$$= 57^\circ\ 17'\ 45''$$

By dividing both sides of equation (3.1) by 180, we find that

$$1^\circ = \frac{\pi}{180} \text{ radians}$$

$$= 0.01745 \text{ radians (rounded)}$$

NOTE: The degree symbol ($^\circ$) is customarily used to indicate degrees, and a pure number with no symbol attached is used to indicate radians. For example, sin 3 should be understood to represent "sine of 3 radians," whereas the "sine of 3 degrees" would be written sin 3°.

The following list indicates other relationships frequently used in trigonometric problems:

Radians	Degrees
$\pi/6$	30
$\pi/4$	45
$\pi/3$	60
$\pi/2$	90
π	180
$3\pi/2$	270
2π	360

EXAMPLE: Express 160° in radians, using π in the answer.

SOLUTION:

$$1^\circ = \frac{\pi}{180} \text{ radians}$$

$$160^\circ = 160 \times 1^\circ$$

$$= 160 \times \frac{\pi}{180} \text{ radians}$$

$$= \frac{8\pi}{9} \text{ radians}$$

EXAMPLE: Express $\pi/20$ in degrees.

SOLUTION:

$$1 \text{ radian} = \frac{180^\circ}{\pi}$$

$$\frac{\pi}{20} \text{ radians} = \frac{\pi}{20} \times 1 \text{ radian}$$

$$= \frac{\pi}{20} \times \frac{180^\circ}{\pi}$$

$$= \frac{180^\circ}{20}$$

$$= 9^\circ$$

Refer to figure 3-5. We can see that if θ represents the number of radians in a central angle, r the length of the radius of the circle, and s the length of the intercepted arc, then the *length of the arc* equals the number of radians multiplied by the length of the radius or

$$s = \theta r$$

Figure 3-5

EXAMPLE: In a circle having a radius of 11 inches, an arc subtends a central angle of 3 radians. Find the length of the arc in inches.

SOLUTION:

$$s = \theta r$$

$$= 3 \cdot 11$$

$$= 33 \text{ inches}$$

PRACTICE PROBLEMS:

1. Find the number of radians in the central angle subtended by an arc 18 inches long in a circle whose radius is 8 inches.

Express the following angles in radians, using π in the answer:

2. 420°

3. 135°

Express the following angles in degrees:

4. 20π

5. $5\pi/6$

6. In a circle whose radius, r, is 4 inches, find in inches the length of arc, s, whose central angle is 1 1/4 radians.

ANSWER:

1. 9/4 radians

2. $7\pi/3$

3. $3\pi/4$

4. 3,600°

5. 150°

6. 5 inches

Because of the relationship of the radian to arc length, the radian has some special applications in measurements of angular velocity and area of a sector.

Angular Velocity

Another type of problem that radian measurement simplifies is that which relates the rotating motion of the wheels of a vehicle

to its forward motion. Here we will not be dealing with angles alone but also with *angular velocity*. Let's analyze this type of motion.

Consider the circle at the left in figure 3-6 to indicate the original position of a wheel. As the wheel turns, it rolls so that the center moves along the line CC', where C' is the center of the wheel at its final position. The contact point at the bottom of the wheel moves an equal distance PP'; but as the wheel turns through angle θ, arc s is made to coincide with line PP'; so,



Figure 3-6.—Angular rotat

$$s = PP' = d$$

or the length of arc is equal to the forward distance, d, the wheel travels. But since

$$s = r\theta$$

then the forward distance that the wheel travels is

$$d = r\theta$$

Dividing both sides of the previous equation by t gives

$$\frac{d}{t} = \frac{r\theta}{t}$$

When a vehicle moves with a constant velocity, v, in time, t, the distance, d, the vehicle travels is expressed by the formula

$$d = vt$$

Solving this formula for v, we have

$$v = \frac{d}{t}$$

The fraction d/t expresses the *linear velocity* of the vehicle, and θ/t is the *angular velocity*. If we let ω (Greek letter omega) stand for the angular velocity, then the equation

$$\frac{d}{t} = \frac{r\theta}{t}$$

becomes

$$v = r\omega$$

where ω is measured in radians per unit time.

EXAMPLE: A car wheel is rotating at 1,050 revolutions per minute (rpm). Find

1. the angular velocity in radians per second.
2. the linear velocity in meters per second on the tire tread, 25 centimeters from the center.

SOLUTION:

1. To find the angular velocity, we need to convert rev/min to rad/sec. To do this, we will apply unit conversions (multiples of one) as follows:

$$\omega = 1{,}050 \frac{\text{rev}}{\text{min}} \times 2\pi \frac{\text{rad}}{\text{rev}} \times \frac{1}{60} \frac{\text{min}}{\text{sec}}$$

$$= \frac{(1{,}050)(2\pi)}{60} \frac{\text{rad}}{\text{sec}}$$

$$= 35\pi \text{ radians per second}$$

2. We find the linear velocity as follows:

$$v = r\omega$$

$$= 25 \text{ cm} \times 35\pi \frac{\text{rad}}{\text{sec}}$$

$$= 875\pi \frac{\text{cm}}{\text{sec}}$$

NOTE: When no unit of angular measure is indicated, the angle is understood to be expressed in radians.

We now need to convert cm/sec to m/sec. We will again apply a unit conversion:

$$v = 875\pi \frac{\text{cm}}{\text{sec}} \times \frac{1}{100} \frac{\text{m}}{\text{cm}}$$

$$= \frac{875\pi}{100} \frac{\text{m}}{\text{sec}}$$

$$= 8.75\pi \text{ meters per second}$$

EXAMPLE: A car is traveling 40 miles per hour. If the wheel radius is 16 inches, what is the angular velocity of the wheels in

1. radians per minute?
2. revolutions per minute?

SOLUTION:

1. We know that

$$v = r\omega$$

Thus,

$$\omega = \frac{v}{r}$$

$$= \frac{40}{16}\frac{\text{mi/hr}}{\text{in}}$$

$$= \frac{5}{2}\frac{\text{mi}}{\text{hr} \times \text{in}} \times \frac{5{,}280}{1}\frac{\text{ft}}{\text{mi}} \times \frac{12}{1}\frac{\text{in}}{\text{ft}} \times \frac{1}{60}\frac{\text{hr}}{\text{min}}$$

$$= \frac{(5)(5{,}280)(12)}{(2)(60)}\frac{\text{rad}}{\text{min}}$$

$$= 2{,}640 \text{ radians per minute}$$

2. Since 2π radians $= 360°$ and $360° = 1$ revolution, then

$$\omega = 2{,}640\frac{\text{rad}}{\text{min}} \times \frac{1}{2\pi}\frac{\text{rev}}{\text{rad}}$$

$$= \frac{2{,}640}{2\pi}\frac{\text{rev}}{\text{min}}$$

$$= 420.2 \text{ revolutions per minute}$$

EXAMPLE: Determine the distance a truck will travel in 1 minute if the wheels are 3 feet in diameter and are turning at the rate of 5 revolutions per second. HINT: Diameter $= 2 \times$ radius

SOLUTION:

$$v = r\omega$$

$$\frac{d}{t} = r\omega$$

$$d = rt\omega$$

$$= \frac{3}{2}\text{ ft} \times 1 \text{ min} \times \left(5\frac{\text{rev}}{\text{sec}} \times \frac{2\pi}{1}\frac{\text{rad}}{\text{rev}}\right)$$

$$= \frac{3}{2}\text{ ft} \times 1 \text{ min} \times 10\pi\frac{\text{rad}}{\text{sec}} \times \frac{60}{1}\frac{\text{sec}}{\text{min}}$$

$$= \frac{(3)(10\pi)(60)}{2}\text{ ft}$$

$$= 2{,}827.43 \text{ feet}$$

ea of a Sector

From plane geometry we find that the area of the sector of ircle is proportional to the angle enclosed in the sector.

Consider sector AOB of the circle shown in figure 3-7. If θ ncreased to 2π radians (360°), it encompasses the entire circle; the area of the circle is proportional to 2π radians. Hence,



Figure 3-7.—Sect circle.

$$\frac{\text{area of sector}}{\text{area of circle}} = \frac{\theta}{2\pi}$$

t the area of a circle can be found by the formula

$$A = \pi r^2$$

substitution, we find

$$\text{area of sector} = \frac{\theta}{2\pi}(\pi r^2)$$

$$= \frac{\theta r^2}{2}$$

erefore, the *area of a sector of a circle* can be found by the rmula

$$A = \frac{1}{2} r^2 \theta$$

ere θ is expressed in radians.

EXAMPLE: Find the area of a sector of a circle with a radius 6 inches having a central angle of 60°.

SOLUTION:

$$A = \frac{1}{2} r^2 \theta$$

$$= \frac{1}{2}(6 \text{ in})^2 \left(60° \times \frac{\pi}{180°}\right)$$

$$= \frac{36\pi}{(2)(3)} \text{ in}^2$$

$$= 6\pi \text{ square inches}$$

The area of a sector of a circle can also be found if the radius and arc length are known. Since

$$s = r\theta$$

then

$$A = \frac{1}{2}r^2\theta$$

$$= \frac{1}{2}r(r\theta)$$

$$= \frac{1}{2}rs$$

EXAMPLE: What is the diameter of a circle if a sector of the circle has an arc length of 9 inches and an area of 18 square inches?

SOLUTION:

If

$$A = \frac{1}{2}rs$$

then

$$r = \frac{2A}{s}$$

$$= \frac{2(18 \text{ in}^2)}{9 \text{ in}}$$

$$= 4 \text{ in}$$

But

$$d = 2r$$

Therefore,

$$d = 2(4 \text{ in})$$

PRACTICE PROBLEMS:

1. A car travels 4,500 feet in 1 minute. The diameter of the wheels is 36 inches. What is the angular velocity of the wheels in radians per minute?

2. How far in feet does a car travel in 1 minute if the radius of the wheels is 18 inches and the angular velocity of the wheels is 1,000 radians per minute?

3. Find the area of a sector of a circle whose central angle is $\pi/3$ and whose diameter is 24 inches. Leave the answer in terms of π.

4. Find the area of a sector of a circle in inches whose arc length is 14 inches and whose radius is 2/3 feet.

ANSWERS:

1. 3,000 radians per minute

2. 1,500 feet

3. 24π square inches

4. 56 square inches

MILS

The *mil* is a unit of small angular measurement, which is not widely used but has some military applications in ranging and sighting. The *mil* is defined in two ways:

1. *As 1/6,400 of the circumference of a circle.*

2. *As the angle subtended by an object 1 unit long, perpendicular to the line of sight, at a distance of 1,000 units.*

rom definition 1 we can see that since

$$360° = 6{,}400 \text{ mils}$$

$$1° = \frac{6{,}400}{360} \text{ mils}$$

$$= \frac{160}{9} \text{ mils}$$

$$= 17.78 \text{ mils (rounded)}$$

, since

$$6{,}400 \text{ mils} = 360°$$

n

$$1 \text{ mil} = \frac{360°}{6{,}400}$$

$$= \frac{9°}{160}$$

$$= 0.05625°$$

EXAMPLE: Convert 240 mils to degrees.

SOLUTION:

$$1 \text{ mil} = \frac{9°}{160}$$

$$240 \text{ mils} = 240 \times 1 \text{ mil}$$

$$= 240 \times \frac{9°}{160}$$

$$= \frac{27°}{2}$$

$$= 13.5°$$

EXAMPLE: Convert 27° to mils.

SOLUTION:

$$1° = \frac{160}{9} \text{ mils}$$

$$27° = 27 \times 1°$$

$$= 27 \times \frac{160}{9} \text{ mils}$$

$$= 480 \text{ mils}$$

Since

$$1 \text{ mil} = \frac{9°}{160}$$

and

$$1° = \frac{\pi}{180} \text{ radians}$$

then

$$1 \text{ mil} = \frac{9}{160} \times 1°$$

$$= \frac{9}{160} \times \frac{\pi}{180} \text{ radians}$$

$$= \frac{\pi}{3{,}200} \text{ radians}$$

$$= 0.00098 \text{ radians (rounded)}$$

We see that *1 mil is approximately 0.001* or *1/1,000 radians*. We also see that *1 radian ≈ 1,000 mils.*

EXAMPLE: Convert 25 mils to an approximate radian measure.

SOLUTION:

$$1 \text{ mil} \approx \frac{1}{1{,}000} \text{ radians}$$

$$25 \text{ mils} = 25 \times 1 \text{ mil}$$

$$\approx 25 \times \frac{1}{1{,}000} \text{ radians}$$

$$\approx \frac{25}{1{,}000} \text{ radians}$$

$$\approx 0.025 \text{ radians}$$

EXAMPLE: Convert 6.48 radians to an approximate
asurement in mils.

SOLUTION:

$$1 \text{ radian} \approx 1{,}000 \text{ mils}$$

$$6.48 \text{ radians} = 6.48 \times 1 \text{ radian}$$

$$\approx 6.48 \times 1{,}000 \text{ mils}$$

$$\approx 6{,}480 \text{ mils}$$

Referring to figure 3-8,
n an angle, θ, subtended by
rc, s, is very small and the
us, r, is large, the chord, c,
lmost equal to the arc, s.

The formula for the length
rc of a circle, as previously
ed, is

$$s = r\theta$$

re θ is in radian mea-
ement.



Figure 3-8.—Relationship of chord and arc.

If the measurement of the arc is made in mils, we must divide
mil measure by 1,000 to obtain the radian measure. Since,

$$1 \text{ mil} = \frac{1}{1{,}000} \text{ radians (approximately)}$$

n

$$m \text{ mils} = \frac{m}{1{,}000} \text{ radians}$$

$$s = r\left(\frac{m}{1{,}000}\right)$$

$$= \frac{rm}{1{,}000}$$

Now, since the chord, c, in figure 3-8, is approximately equal to the arc, s, then

$$c = \frac{rm}{1{,}000}$$

Now consider definition 2. If

$$r = 1{,}000 \text{ yds}$$

and

$$m = 1 \text{ mil}$$

then

$$c = \frac{rm}{1{,}000}$$

$$= \frac{1{,}000 \times 1}{1{,}000}$$

$$= 1 \text{ yard}$$

We also know that the arc, s, is approximately equal to 1 yard since $s \approx c$.

The military uses the fact that a mil subtends a yard at a distance of 1,000 yards for quick computations in the field.

EXAMPLE: Find the length of a target if, at a right angle to the line of sight, it subtends an angle of 15 mils at a range of 4,000 yards.

SOLUTION:

$$c = \frac{rm}{1{,}000}$$

$$= \frac{4{,}000 \times 15}{1{,}000}$$

$$= 60 \text{ yards}$$

EXAMPLE: A building known to be 80 feet long and perpendicular to the line of sight subtends an angle of 100 mils. What is the approximate range to the building?

SOLUTION:

Since

$$c = \frac{rm}{1{,}000}$$

then

$$r = \frac{1{,}000c}{m}$$

$$= \frac{1{,}000 \times 80}{100}$$

$$= 800 \text{ feet}$$

PRACTICE PROBLEMS:

1. Convert 3,456 mils to degrees.
2. Convert 12 degrees to mils.
3. Convert 27,183 mils to an approximate radian measure.
4. Convert 431 radians to an approximate measurement in mils.
5. A tower 500 feet away subtends a vertical angle of 250 mils. What is the height of the tower?
6. If points X and Y are 48 yards apart and are 4,000 yards from an observer, how many mils do they subtend?

ANSWERS:

1. 194.4°
2. 213.3 mils
3. 27.183 radians

4. 431,000 mils

5. 125 feet

6. 12 mils

PROPERTIES OF RIGHT TRIANGLES

Mathematics, Volume 1, contains information on the trigonometric ratios and other properties of triangles. This section restates some of the properties of right triangles for review and reference.

PYTHAGOREAN THEOREM

The *Pythagorean theorem* states that in a right triangle, the square of the length of the hypotenuse (longest side) is equal to the sum of the squares of the lengths of the other two sides. In the right triangle shown in figure 3-9, this relationship is expressed as

$$r^2 = x^2 + y^2$$

where r is the length of the hypotenuse and x and y are the lengths of the other two sides.



Figure 3-9.-

This relationship is useful in solving many problems and in developing trigonometric concepts.

EXAMPLE: In figure 3-10, what is the length of the hypotenuse of the right triangle if the lengths of the other two sides are 3 and 4?



Figure 3-1
with hyp

SOLUTION:

$$\begin{aligned} r^2 &= x^2 + y^2 \\ &= 4^2 + 3^2 \\ &= 16 + 9 \\ &= 25 \end{aligned}$$

So,

$$\begin{aligned} r &= \sqrt{25} \\ &= 5 \end{aligned}$$

NOTE: We will use the positive value of the square root since we are dealing with lengths.

EXAMPLE: Figure 3-11 shows a right triangle with a hypotenuse equal to 40 and one of the other sides equal to 10. What is the length of the remaining side?



Figure 3-11.—Right triangl unknown.

SOLUTION:

$$r^2 = x^2 + y^2$$

or

$$x^2 = r^2 - y^2$$

$$= 40^2 - 10^2$$

$$= 1{,}600 - 100$$

$$= 1{,}500$$

So,

$$x = \sqrt{1{,}500}$$

$$= 38.7 \text{ (rounded)}$$

SIMILAR RIGHT TRIANGLES

Another relationship of right triangles that is useful in trigonometry concerns *similar triangles*. Whenever the angles of one triangle are equal to the corresponding angles in another triangle, the two triangles are said to be *similar*.

For example, right triangle A in figure 3-12 is similar to right triangle B. Since the two triangles are similar by definition, the following proportions involving the lengths of the corresponding sides are true:

$$\frac{a}{a'} = \frac{b}{b'} = \frac{c}{c'}$$

This relationship can be used to find the lengths of unknown sides in similar triangles.



Figure 3-12.—Similar triangl

EXAMPLE: Assume right triangles A and B in figure 3-13 are similar with lengths as shown. Find the lengths of sides b' and c'.



Figure 3-13.—Simila exar

SOLUTION:

Since

$$\frac{a}{a'} = \frac{b}{b'} = \frac{c}{c'}$$

then

$$\frac{10}{7} = \frac{11.18}{b'} = \frac{5}{c'}$$

Side b' can be solved for using the first two ratios:

$$\frac{10}{7} = \frac{11.18}{b'}$$

So,

$$b' = \frac{11.18 \times 7}{10}$$

$$= \frac{78.26}{10}$$

$$= 7.826$$

Side c' can be solved for using the first and third ratios:

$$\frac{10}{7} = \frac{5}{c'}$$

So,

$$c' = \frac{5 \times 7}{10}$$

$$= 3.5$$

Recall from plane geometry that *the sum of the interior angles of any triangle is equal to 180°*. Using this fact, we can assume that two triangles are similar if two angles of one are equal to two angles of the other. The remaining angle in any triangle must be equal to 180° minus the sum of the other two angles.

an acute angle of one right triangle is equal to an acute angle other right triangle, the triangles are similar because the right s in the two triangles are also equal to each other.
ence, if θ is one of the acute angles in a right triangle, then $- \theta)$ is the other acute angle, such that

$$90° + \theta + (90° - \theta) = 180°$$

fore, *two right triangles are similar if an acute angle of one le is equal to an acute angle of the other triangle.*
any practical uses of trigonometry are based on the fact that ight triangles are similar if an acute angle of one triangle ıal to an acute angle of the other triangle.
figure 3-14 triangle A is similar to triangle B an acute angle in triangle A is equal to an acute in triangle B. Since triangle A is similar to le B, then

$$\frac{x}{x'} = \frac{y}{y'} = \frac{r}{r'}$$

hanging terms in the proportions gives

$$\frac{x}{y} = \frac{x'}{y'}$$

$$\frac{y}{r} = \frac{y'}{r'}$$

$$\frac{x}{r} = \frac{x'}{r'}$$

are considered among the main ples of numerical trigonometry.



Figure 3-14.—Similar right triangl

RACTICE PROBLEMS:

efer to figure 3-15 in solving the llowing problems:

1. Use the Pythagorean theorem to calculate the unknown length in triangle A.



Figure 3-15.—Triangles for pr problems.

2. Use the Pythagorean theorem to calculate the unknown length in triangle B.

3. Triangles C and D are similar triangles. Find the length of sides a and b in triangle D.

ANSWERS:

1. $\sqrt{5}$

2. $2\sqrt{2}$

3. $a = 15\sqrt{2}/4$, $b = 3/2$

TRIGONOMETRIC RATIOS, FUNCTIONS, AND TABLES

The properties of triangles given in the previous section provide a means for solving many practical problems. Certain practical problems, however, require knowledge of right triangle relationships other than the Pythagorean theorem or the relationships of similar triangles before solutions can be found.

For example, the following two problems require additional knowledge:

1. Find the values of the unknown sides and angles in a right triangle when the values of one side and one acute angle are given.

2. Find the value of the unknown side and the values of the angles in a right triangle when two sides are known.

The additional relationships between the sides and angles of a right triangle are called *trigonometric ratios*. These ratios were introduced in *Mathematics*, Volume 1, and are reviewed in the following paragraphs. The basic foundations of trigonometry rest upon these ratios and their associated trigonometric functions.

GONOMETRIC RATIOS D FUNCTIONS

The sides of a right triangle form six ratios. In re 3-16 we will use the acute angle θ and the three s x, y, and r two at a time to define the onometric ratios. These ratios and the onometric functions associated with each ratio are d as follows:

$$\text{the sine of } \theta = \frac{y}{r}, \text{ written } \sin \theta$$

$$\text{the cosine of } \theta = \frac{x}{r}, \text{ written } \cos \theta$$

$$\text{the tangent of } \theta = \frac{y}{x}, \text{ written } \tan \theta$$

$$\text{the cotangent of } \theta = \frac{x}{y}, \text{ written } \cot \theta$$

$$\text{the secant of } \theta = \frac{r}{x}, \text{ written } \sec \theta$$

$$\text{the cosecant of } \theta = \frac{r}{y}, \text{ written } \csc \theta$$



Figure 3-16.—Right triangle for dete ratios.

The trigonometric functions of a right triangle are embered easier by the convention of naming the sides. er to figure 3-17. The side of length y is called the *opposite* angle θ, the side of length x is called the *adjacent* to angle θ, and the side of length r is called *hypotenuse*. Using this terminology causes the six onometric functions to be defined as:

$$\sin \theta = \frac{y}{r} = \frac{\text{opposite}}{\text{hypotenuse}}$$

$$\cos \theta = \frac{x}{r} = \frac{\text{adjacent}}{\text{hypotenuse}}$$

$$\tan \theta = \frac{y}{x} = \frac{\text{opposite}}{\text{adjacent}}$$

$$\cot \theta = \frac{x}{y} = \frac{\text{adjacent}}{\text{opposite}}$$

$$\sec \theta = \frac{r}{x} = \frac{\text{hypotenuse}}{\text{adjacent}}$$

$$\csc \theta = \frac{r}{y} = \frac{\text{hypotenuse}}{\text{opposite}}$$



Figure 3-17.—Names of sid right triangle.

Remember that the six trigonometric ratios apply only to the acute angles of a right triangle.

EXAMPLE: Give the values of the trigonometric functions of the angle in the right triangle for figure 3-18, view A.

SOLUTION:



Figure 3-18.—Pr

$$\sin \theta = \frac{y}{r} = \frac{3}{5} = 0.6$$

$$\cos \theta = \frac{x}{r} = \frac{4}{5} = 0.8$$

$$\tan \theta = \frac{y}{x} = \frac{3}{4} = 0.75$$

$$\cot \theta = \frac{x}{y} = \frac{4}{3} = 1.33333 \text{ (rounded)}$$

$$\sec \theta = \frac{r}{x} = \frac{5}{4} = 1.25$$

$$\csc \theta = \frac{r}{y} = \frac{5}{3} = 1.66667 \text{ (rounded)}$$

EXAMPLE: Give the values of the trigonometric functions of the angle in the right triangle for figure 3-18, view B.

SOLUTION: Only two sides are given. To find the third side of the right triangle, use the Pythagorean theorem:

$$r^2 = x^2 + y^2$$

and

$$\begin{aligned} y^2 &= r^2 - x^2 \\ &= 6^2 - 3^2 \\ &= 36 - 9 \\ &= 27 \\ y &= \sqrt{27} \\ &= \sqrt{9 \cdot 3} \\ &= 3\sqrt{3} \end{aligned}$$

Now, using the values of x, y, and r, we find the values of the six trigonometric functions are as follows:

$$\sin\theta = \frac{y}{r} = \frac{3\sqrt{3}}{6} = \frac{\sqrt{3}}{2} = 0.86603 \text{ (rounded)}$$

$$\cos\theta = \frac{x}{r} = \frac{3}{6} = \frac{1}{2} = 0.5$$

$$\tan\theta = \frac{y}{x} = \frac{3\sqrt{3}}{3} = \sqrt{3} = 1.73205 \text{ (rounded)}$$

$$\cot\theta = \frac{x}{y} = \frac{3}{3\sqrt{3}} = \frac{1}{\sqrt{3}} = \frac{\sqrt{3}}{3} = 0.57735 \text{ (rounded)}$$

$$\sec\theta = \frac{r}{x} = \frac{6}{3} = 2$$

$$\csc\theta = \frac{r}{y} = \frac{6}{3\sqrt{3}} = \frac{2}{\sqrt{3}} = \frac{2\sqrt{3}}{3} = 1.15470 \text{ (rounded)}$$

TABLES OF TRIGONOMETRIC FUNCTIONS

Tables of trigonometric functions give the numerical values of the ratios of the sides of a right triangle that correspond to the trigonometric functions. Appendixes II and III are tables of trigonometric functions. These tables give values rounded to five decimal places of trigonometric functions for each minute from 0° to 90°. *Appendix II consists of tables of natural sines and cosines. Appendix III consists of tables of natural tangents and cotangents.*

For example, if we wanted to find sin 3° 25′, we would use appendix II, Natural Sines and Cosines, to first locate 3° on the first row of the table. Next, we would locate sin under 3° on the second row. Then, we would locate 25 along the first column of the table. Now, reading left to right across from 25 and from top to bottom under sin 3°, we find sin 3° 25′ = 0.05960. If we wanted to find cos 86° 35′, we would first locate 86° on the last row of the table. (The degrees on the top row range from 0° to 44°, and the degrees on the last row range from 45° to 90°.) Next, we would locate cos above 86° on the next to the last row. Then, we would locate 35 along the last column of the table. Now, reading right to left across from 35 and from bottom to top above cos 86°, we find cos 86° 35′ = 0.05960. Note that sin 3° 25′ = 0.05960 = cos 86° 35′. The reason for this will be discussed in chapter 4.

The tables in appendix III, Natural Tangents and Cotangents, are arranged in the same format as the tables in appendix II and are used in the same way. NOTE: Scientific calculators will give you the same values rounded to five decimal places as supplied in the tables in appendixes II and III.

Most tables list the sine, cosine, tangent, and cotangent of angles from 0° to 90°. Very few give the secant and cosecant since these functions of an angle are seldom used. When needed, they may be found from the values of the sine and cosine as follows:

$$\sec \theta = \frac{r}{x} = \frac{1}{\frac{x}{r}} = \frac{1}{\cos \theta}$$

and

$$\csc \theta = \frac{r}{y} = \frac{1}{\frac{y}{r}} = \frac{1}{\sin \theta}$$

Hence, the reciprocal of the secant function is the cosine function, and the reciprocal of the cosecant function is the sine function.

The tangent and cotangent functions may also be expressed in terms of the sine and cosine functions as follows:

$$\tan \theta = \frac{y}{x} = \frac{\frac{y}{r}}{\frac{x}{r}} = \frac{\sin \theta}{\cos \theta}$$

and

$$\cot \theta = \frac{x}{y} = \frac{\frac{x}{r}}{\frac{y}{r}} = \frac{\cos \theta}{\sin \theta}$$

In addition, the cotangent function may be determined as the reciprocal of the tangent function as follows:

$$\cot \theta = \frac{x}{y} = \frac{1}{\frac{y}{x}} = \frac{1}{\tan \theta}$$

NOTE: These relationships are the fundamental trigonometric identities that will be used extensively in solving more complex identities in chapter 6.

USE OF TRIGONOMETRIC RATIOS AND FUNCTIONS

The trigonometric ratios and trigonometric functions furnish powerful tools for use in problem solving of right triangles. Finding the remaining parts of a right triangle is possible if, in addition to the right angle, the length of one side and the length of any other side or the value of one of the acute angles is known.

EXAMPLE: Find the length of side y in figure 3-19, view A.

SOLUTION: We can use

$$\tan \theta = \frac{y}{x}$$

since we know one side and one angle. Thus,

$$\tan 35° = \frac{y}{20}$$

From appendix III (or calculator), we find that

$$\tan 35° = 0.70021$$

So,

$$0.70021 = \frac{y}{20}$$

$$y = (0.70021)(20)$$

$$= 14.0042$$

We could have also used cos θ, cot θ, or sec θ to find side y.



Figure 3-19.—Practical use o

EXAMPLE: Find the value of r in figure 3-19, view B.

SOLUTION:

$$\sin \theta = \frac{y}{r}$$

$$\sin 65° = \frac{5}{r}$$

$$r = \frac{5}{\sin 65°}$$

$$r = \frac{5}{0.90631}$$

$$= 5.51688$$

We could have also used csc θ to find side y.

PRACTICE PROBLEMS:

Refer to figure 3-20 in working problems 1 through 4.

1. Find the values of the trigonometric functions of angle θ for the right triangle in view A.

2. Find the value of side y in view B using the sine function.

3. Find the value of side x in view C using the cosine function.

4. Find the value of side y in view D using the tangent function.



Figure 3-20.—Triangles for practice proble

ANSWERS:

1. $\sin \theta = 21/35 = 3/5 = 0.6$

   $\cos \theta = 28/35 = 4/5 = 0.8$

   $\tan \theta = 21/28 = 3/4 = 0.75$

   $\cot \theta = 28/21 = 4/3 = 1.33333$

   $\sec \theta = 35/28 = 5/4 = 1.25$

   $\csc \theta = 35/21 = 5/3 = 1.66667$

2. 294.13394

3. 12.12442

4. 8.48655

SUMMARY

The following are the major topics covered in this chapter:

1. **Terminology:**

   Radius vector—The line that is rotated to generate an angle.

   Initial position—The original position of the radius vector.

   Terminal position—The final position of the radius vector.

   Positive angle—The angle generated by rotating the radius vector counterclockwise from the initial position.

   Negative angle—The angle generated by rotating the radius vector clockwise from the initial position.

2. **Degrees:** The degree system is the most common system of angular measurement. In this system a complete revolution is divided into 360 equal parts called *degrees*.

   $1 \text{ revolution} = 360°$

   $1° = 60'$

   $1' = 60''$

   For convenience, the 360° are divided into four equal parts of 90° each called *quadrants*.

   If $0° < \theta < 90°$, then θ is in quadrant I.

   If $90° < \theta < 180°$, then θ is in quadrant II.

   If $180° < \theta < 270°$, then θ is in quadrant III.

   If $270° < \theta < 360°$, then θ is in quadrant IV.

   If $\theta > 360°$, then θ lies in the same quadrant as $\theta - n(360°)$, where $n = 1, 2, 3, \ldots$ and $n(360°) < \theta$.

3. **Radians:** An even more fundamental method of angular measurement involves the *radian*. A *radian* is defined as an angle that, if its vertex is placed at the center of a circle, intercepts an arc equal in length to the radius of the circle.

   $2\pi \text{ radians} = 360°$

   $\pi \text{ radians} = 180°$

$$1 \text{ radian} = \frac{180°}{\pi}$$

$$1° = \frac{\pi}{180°} \text{ radians}$$

The radian measure of an angle, θ, is the ratio of the length of the arc, s, it subtends to the length of the radius vector, r, of the circle in which it is the central angle or

$$\theta = \frac{s}{r}$$

4. **Other frequently used relationships between radians and degrees:**

Radians	Degrees
$\pi/6$	30
$\pi/4$	45
$\pi/3$	60
$\pi/2$	90
π	180
$3\pi/2$	270
2π	360

5. **Length of arc:**

$$s = \theta r$$

where θ represents the number of radians in a central angle, r the length of the radius of the circle, and s the length of the intercepted arc.

6. **Angular velocity:**

$$\omega = \frac{\theta}{t}$$

where θ is measured in radians and t is the unit time.

7. **Linear velocity:**

$$v = \frac{d}{t}$$

where d is the distance and t is the unit time.

$$v = r\omega$$

where r is the radius and ω is the angular velocity.

8. **Area of a sector of a circle:**

$$A = \frac{1}{2}r^2\theta$$

where θ is expressed in radians.

$$A = \frac{1}{2}rs$$

where r is the radius and s is the arc length.

9. **Mils:** The *mil* is a unit of small angular measurement that has military applications. The *mil* is defined as follows:

   1. 1/6,400 of the circumference of a circle.

      $360° = 6{,}400$ mils

      $1° = \frac{160}{9}$ mils

      $1 \text{ mil} = \frac{9°}{160}$

   2. The angle subtended by an object 1 unit long, perpendicular to the line of sight, at a distance of 1,000 units.

      $1 \text{ mil} \approx \frac{1}{1{,}000}$ radians

      1 radian $\approx$ 1,000 mils

10. **Pythagorean theorem:** The *Pythagorean theorem* states that in a right triangle, the square of the length of the hypotenuse, r, is equal to the sum of the squares of the lengths of the other two sides, x and y, or

$$r^2 = x^2 + y^2$$

11. **Similar triangles**: Whenever the angles of one triangle are equal to the corresponding angles in another triangle, the two triangles are said to be *similar* and the following proportions involving the lengths of their corresponding sides are true:

$$\frac{a}{a'} = \frac{b}{b'} = \frac{c}{c'}$$

12. **Similar right triangles:** Two right triangles are similar if an acute angle of one triangle is equal to an acute angle of the other triangle. The following proportions involving the lengths of their corresponding sides are true:

$$\frac{x}{x'} = \frac{y}{y'} = \frac{r}{r'}$$

13. **Trigonometric ratios and functions:**

$$\sin \theta = \frac{y}{r} = \frac{\text{opposite}}{\text{hypotenuse}}$$

$$\cos \theta = \frac{x}{r} = \frac{\text{adjacent}}{\text{hypotenuse}}$$

$$\tan \theta = \frac{y}{x} = \frac{\text{opposite}}{\text{adjacent}}$$

$$\cot \theta = \frac{x}{y} = \frac{\text{adjacent}}{\text{opposite}}$$

$$\sec \theta = \frac{r}{x} = \frac{\text{hypotenuse}}{\text{adjacent}}$$

$$\csc \theta = \frac{r}{y} = \frac{\text{hypotenuse}}{\text{opposite}}$$

14. **Tables of trigonometric functions:** Tables of trigonometric functions give the numerical values of the ratios of the sides of a right triangle that correspond to the trigonometric functions. Appendix II consists of tables of natural sines and and cosines. Appendix III consists of tables of natural tangents and cotangents.

ADDITIONAL PRACTICE PROBLEMS

1. In which quadrant is the angle 5,370°?
2. Find the radian measure of the central angle in a circle with radius π inches if the angle subtends an arc of $3\pi/5$ inches.
3. Express 4,320° in radians, using π in the answer.
4. Express $11\pi/12$ in degrees.
5. If the length of the radius of a circle is 5 meters, find the length of arc subtended by a central angle with measure π radians.
6. Kim and Tom are riding on a Ferris wheel. Kim observes that it takes 30 seconds to make a complete revolution. Their seat is 35 feet from the axle of the wheel.
   a. What is their angular velocity in radians per second?
   b. What is their linear velocity in feet per minute?
7. Find the area of a sector of a circle if its central angle is 45° and the diameter of the circle is 28 centimeters.
8. Convert 17 7/9 mils to degrees.
9. Convert 3.6 degrees to mils.
10. Convert 9/5 mils to an approximate radian measure.
11. Convert 0.00145 radians to an approximate measurement in mils.
12. An airplane with a wing span of 84 feet is flying toward an observer. What is the distance of the plane from the observer when the plane subtends 7 mils?
13. The length of the hypotenuse of a right triangle is 17, and the length of one of the other sides is 8. What is the length of the remaining side?
14. Assume similar right triangles A and B have sides x, y, r, and x', y', r', respectively. If $x = 6$, $y = 8$, $r = 10$, and $y' = 1/2$, what are the values of x' and r'?
15. Find the values of the trigonometric functions θ of in a right triangle if the hypotenuse is 25 and the side adjacent to θ is 24.
16. If in a right triangle one of the acute angles is 56° 17′ and the hypotenuse is 10, what are the lengths of the other two sides?

ANSWERS TO ADDITIONAL PRACTICE PROBLEMS

1. 4th
2. 3/5
3. 24π
4. $165°$
5. 5π meters
6. a. $\pi/15$ radians per second

   b. 140π feet per minute
7. $49\pi/2$ square centimeters
8. $1°$
9. 64 mils
10. 0.0018 radians
11. 1.45 mils
12. 12,000 feet
13. 15
14. $x' = 3/8$

    $z' = 5/8$
15. $\sin \theta = 7/25 = 0.28$

    $\cos \theta = 24/25 = 0.96$

    $\tan \theta = 7/24 = 0.29167$ (rounded)

    $\cot \theta = 24/7 = 3.42857$ (rounded)

    $\sec \theta = 25/24 = 1.04167$ (rounded)

    $\csc \theta = 25/7 = 3.57143$ (rounded)
16. 5.5509 and 8.3179

CHAPTER 4

TRIGONOMETRIC ANALYSIS

LEARNING OBJECTIVES

Upon completion of this chapter, you should be able to do the following:

1. Use the rectangular coordinate system to determine the algebraic signs and the values of the trigonometric functions and to locate and define the trigonometric functions.
2. Relate any angle in standard position to its reference angle.
3. Determine the trigonometric functions of an angle in any quadrant, of negative angles, of coterminal angles, of frequently used angles, and of quadrantal angles.
4. Express the trigonometric functions of an angle in terms of their complement.
5. Recognize characteristics of the graphs of the sine, cosine, and tangent functions.

INTRODUCTION

This chapter is a continuation of the broad topic of trigonometry introduced in chapter 3. The topic is expanded in this chapter to allow analysis of angles greater than 90°. The chapter is extended as a foundation for analysis of the generalized angle; that is, an angle of any number of degrees. Additionally, the chapter introduces the concept of both positive and negative angles.

RECTANGULAR COORDINATE SYSTEM

The rectangular, or Cartesian, coordinate system introduced in *Mathematics*, Volume 1, was used in solving equations; in this

apter it is used to analyze the generalized angle. ıe following is a brief review of the rectangular ordinate system:

1. The vertical axis (Y axis in fig. 4-1) is considered positive above the origin and negative below the origin.

2. The horizontal axis (X axis in fig. 4-1) is positive to the right of the origin and negative to the left of the origin.

3. A point, $P(x,y)$, anywhere in a rectangular coordinate system may be located by two numbers. The value of x is called the *abscissa*. The value of y is called the *ordinate*. The abscissa and ordinate of a point are its *coordinates*.



Figure 4-1.—Rectangular coordina

4. In notation used to locate points, the coordinates are conventionally placed in parentheses and separated with a comma, with the abscissa always written first. The general form of this notation is $P(x,y)$. Thus, point P in figure 4-1 would have the notation $P(4,-5)$.

5. The quadrants are numbered in the manner described in chapter 3 of this course (shown as Roman numerals in figure 4-1).

6. The x coordinate is positive in the first (I) and fourth (IV) quadrants and negative in the second (II) and third (III) quadrants. The y coordinate is positive in the first and second quadrants and negative in the third and fourth quadrants. The signs of the coordinates are shown in parentheses in figure 4-1. The algebraic signs of the coordinates of a point are used in this chapter for determining the algebraic signs of trigonometric functions.

ANGLES IN STANDARD POSITION

To construct an *angle in standard position*, first lay out a ctangular coordinate system. Then draw the angle, θ, so at its vertex is at the origin of the coordinate system and its itial or original side is lying along the positive X axis as shown in

figure 4-2. The terminal or final side of the angle will lie in any of the quadrants or on one of the axes separating the quadrants. When the terminal side falls on an axis, the angle is called a *quadrantal angle*, which will be discussed later in this chapter. In figure 4-2 the terminal side lies in quadrant II.

The quadrant in which an angle lies is determined by the terminal side. When an angle is placed in standard position, the angle is said to lie in the quadrant containing the terminal side. For example, the negative angle, θ, shown in standard position in figure 4-3, is said to lie in the second quadrant.

When two or more angles in standard position have their terminal sides located at the same position, they are said to be *coterminal*. If θ is any general angle, then θ plus or minus an integral multiple of 360° yields a coterminal angle.

For example, the angles θ, ϕ, and α in figure 4-4 are said to be coterminal angles. If

$$\theta = 45°$$

then

$$\begin{aligned}\phi &= \theta - 360° \\ &= 45° - 360° \\ &= -315°\end{aligned}$$

and

$$\begin{aligned}\alpha &= \theta + 360° \\ &= 45° + 360° \\ &= 405°\end{aligned}$$

The relationship of coterminal angles can be stated in a general form. *For any angle θ measured in degrees, any angle ϕ coterminal with θ can be found by*

$$\phi = \theta + n(360°)$$

where n is any integer (positive, negative, or zero); that is,

$$n = 0, \pm 1, \pm 2, \pm 3, \ldots$$



Figure 4-2.—Ar



Figure 4-3.—
qua



Figure 4-4.—

The principle of coterminal angles is used in developing other trigonometric relationships and other phases of trigonometric analysis. An expansion of this principle, discussed later in this chapter, states that the trigonometric functions of coterminal angles have the same value.

PRACTICE PROBLEMS:

Determine whether or not the following sets of angles are coterminal:

1. $60°, -300°, 420°$

2. $0°, 360°, 180°$

3. $45°, -45°, 345°$

4. $735°, -345°, -705°$

ANSWERS:

1. Coterminal

2. Not coterminal

3. Not coterminal

4. Coterminal

DEFINITIONS OF THE TRIGONOMETRIC FUNCTIONS

So far, the trigonometric functions have been defined as follows:

1. By labeling the sides of a right triangle x, y, and r.
2. By naming the sides of a right triangle adjacent, opposite, and hypotenuse.





Figure 4-5.—Functions of general angles.

In this chapter we will introduce a third set of definitions using the nomenclature of the coordinate system. Note that each definition defines the same functions using different terminology.

To arrive at the third set of definitions, construct an angle in standard position on a coordinate system as shown in figure 4-5, view A. Choose point $P(x,y)$ on the final position of the radius vector. Distance OP is denoted by the positive number r for the length of the radius.

By constructing a right triangle using $P(x,y)$ and r, as in figure 4-5, view B, the six trigonometric functions are classified as follows:

$$\sin \theta = \frac{y}{r} = \frac{\text{ordinate}}{\text{length of radius}}$$

$$\cos \theta = \frac{x}{r} = \frac{\text{abscissa}}{\text{length of radius}}$$

$$\tan \theta = \frac{y}{x} = \frac{\text{ordinate}}{\text{abscissa}}$$

$$\cot \theta = \frac{x}{y} = \frac{\text{abscissa}}{\text{ordinate}}$$

$$\sec \theta = \frac{r}{x} = \frac{\text{length of radius}}{\text{abscissa}}$$

$$\csc \theta = \frac{r}{y} = \frac{\text{length of radius}}{\text{ordinate}}$$

The value of each function is dependent on angle θ and not on the selection of point $P(x,y)$. If a different point were chosen, the length of r, as well as the values of the x and y coordinates, would change proportionally, but the ratios would be unchanged.

EXAMPLE: Find the sine and cosine of angle θ in figure 4-5, view A, for the point $P(3,4)$.

SOLUTION: To determine the sine and cosine of θ, we must find the value of r. Since the values of the x and y coordinates correspond to the lengths of the sides x and y in figure 4-5, view B, we can determine the length of r by using the Pythagorean theorem or by recalling from *Mathematics*, Volume 1, the 3-4-5 triangle. In either case, the length of r is 5 units. Hence,

$$\sin \theta = \frac{\text{ordinate}}{\text{length of radius}}$$

$$= \frac{4}{5}$$

and

$$\cos \theta = \frac{\text{abscissa}}{\text{length of radius}}$$

$$= \frac{3}{5}$$

NOTE: For the remainder of this chapter, all angles are understood to be in standard position, unless otherwise stated.

PRACTICE PROBLEMS:

Find the sine, cosine, and tangent of the angles whose radius vectors pass through the following points:

1. $P(5,12)$

2. $P(1,1)$

3. $P(1, \sqrt{3})$

4. $P(3,2)$

ANSWERS:

1. $\sin \theta = 12/13$

   $\cos \theta = 5/13$

   $\tan \theta = 12/5$

2. $\sin \theta = 1/\sqrt{2} = \sqrt{2}/2$

   $\cos \theta = 1/\sqrt{2} = \sqrt{2}/2$

   $\tan \theta = 1/1 = 1$

3. $\sin \theta = \sqrt{3}/2$

   $\cos \theta = 1/2$

   $\tan \theta = \sqrt{3}/1 = \sqrt{3}$

4. $\sin \theta = 2/\sqrt{13} = 2\sqrt{13}/13$

   $\cos \theta = 3/\sqrt{13} = 3\sqrt{13}/13$

   $\tan \theta = 2/3$

QUADRANT SYSTEM

The quadrants formed in the rectangular coordinate system are used to determine the algebraic signs of the trigonometric functions. The quadrants in figure 4-6 show the algebraic signs of the trigonometric functions in the various quadrants.

In the first quadrant the abscissa and ordinate are always positive. The radius vector is always taken as positive. Therefore, *all the trigonometric ratios are positive for angles in the first quadrant. For angles in the second quadrant, only the ratios involving the ordinate and the radius vector are positive. These are the sine and cosecant ratios. For angles in the third quadrant, where the ordinate and abscissa are both negative, only the ratios involving the abscissa and the ordinate are positive.*

II

sin θ = +/+ = +
cos θ = -/+ = -
tan θ = +/- = -
cot θ = -/+ = -
sec θ = +/- = -
csc θ = +/+ = +

III

sin θ = -/+ = -
cos θ = -/+ = -
tan θ = -/- = +
cot θ = -/- = +
sec θ = +/- = -
csc θ = +/- = -

Figure 4-6.—Sig

These are the tangent and cotangent ratios. For angles in the fourth quadrant, ratios involving the radius vector and the abscissa are positive. These are the cosine and the secant ratios.

NOTE: In each quadrant the sine and cosecant have the same sign, the cosine and the secant have the same sign, and the tangent and cotangent have the same sign.

The last group of practice problems involved angles in the first quadrant only, where all of the functions were positive. When an angle lies in one of the other quadrants, the trigonometric functions may be positive or negative.

EXAMPLE: Find all of the trigonometric functions of θ if $\tan \theta = 5/12$, $\sin \theta < 0$, and $r = 13$.

SOLUTION: Reference to figure 4-6 shows that an angle with a positive tangent and a negative sine can only occur in the third quadrant. The point in the third quadrant has coordinates $(-12, -5)$. (See fig. 4-7)

We can now read the trigonometric ratios from the figure:

$$\sin \theta = \frac{\text{ordinate}}{\text{length of radius}} = \frac{-5}{13}$$

$$\cos \theta = \frac{\text{abscissa}}{\text{length of radius}} = \frac{-12}{13}$$

$$\tan \theta = \frac{\text{ordinate}}{\text{abscissa}} = \frac{-5}{-12} = \frac{5}{12}$$

$$\cot \theta = \frac{\text{abscissa}}{\text{ordinate}} = \frac{-12}{-5} = \frac{12}{5}$$

$$\sec \theta = \frac{\text{length of radius}}{\text{abscissa}} = \frac{13}{-12} = \frac{-13}{12}$$

$$\csc \theta = \frac{\text{length of radius}}{\text{ordinate}} = \frac{13}{-5} = \frac{-13}{5}$$



Figure 4-7.—Finding th functions for a third-q

EXAMPLE: Find all of the trigonometric functions of θ if $\csc \theta = -17/15$ and $\cos \theta < 0$.

SOLUTION: The cosecant is negative in the same quadrants as the sine; that is, quadrants III and IV. The cosine is negative

in quadrants II and III. Therefore, the cosecant and cosine are both negative in quadrant III. (Refer to fig. 4-6.) The ordinate in the third quadrant is -15 and the radius is 17.

NOTE: The fraction $-17/15$ indicates that either the numerator or denominator is negative, but not both. In this case, we know that the ordinate (denominator) is negative since the radius (numerator) is always positive.

From the Pythagorean theorem the abscissa in the third quadrant is

$$x^2 = r^2 - y^2$$

$$= (17)^2 - (-15)^2$$

$$= 289 - 225$$

$$= 64$$

$$x = -8$$

Therefore, referring to figure 4-8, the six trigonometric functions are as follows:

$$\sin \theta = -15/17$$

$$\cos \theta = -8/17$$

$$\tan \theta = -15/-8 = 15/8$$

$$\cot \theta = -8/-15 = 8/15$$

$$\sec \theta = 17/-8 = -17/8$$

$$\csc \theta = 17/-15 = -17/15$$



Figure 4-8.—Constructi quadran

EXAMPLE: If $\sec \theta = -25/24$ and $\tan \theta = -7/24$, find the other four trigonometric ratios of θ.

SOLUTION: The secant and tangent are both negative in the second quadrant. In the second quadrant the abscissa is -24, the ordinate is 7, and the radius is 25 (refer to fig. 4-9); so,

$$\sin \theta = 7/25$$

$$\cos \theta = -24/25$$

$$\cot \theta = -24/7$$

$$\csc \theta = 25/7$$



Figure 4-9.—Construction quadrant 2.

PRACTICE PROBLEMS:

Without using tables, find the six trigonometric functions of θ under the following conditions:

1. $\tan \theta = 3/4$, $r = 5$, and θ is not in the first quadrant.
2. $\tan \theta = -21/20$, $r = 29$, and $\cos \theta > 0$.
3. $\cos = -3/5$ and $\cot \theta = 3/4$.
4. $\tan \theta = -8/15$ and $\csc \theta$ is positive.

Indicate the quadrant in which the terminal side of θ lies for the following conditions:

5. $\sin \theta > 0$ and $\cos \theta < 0$
6. $\cos \theta < 0$ and $\csc \theta < 0$
7. $\sec \theta > 0$ and $\cot \theta < 0$

ANSWERS:

1. $\sin \theta = -3/5$

   $\cos \theta = -4/5$

   $\tan \theta = -3/-4 = 3/4$

   $\cot \theta = -4/-3 = 4/3$

   $\sec \theta = 5/-4 = -5/4$

2. $\sin \theta = -21/29$

   $\cos \theta = 20/29$

   $\tan \theta = -21/20$

   $\cot \theta = 20/-21 = -20/21$

   $\sec \theta = 29/20$

   $\csc \theta = 29/-21 = -29/21$

3. $\sin \theta = -4/5$

   $\cos \theta = -3/5$

   $\tan \theta = -4/-3 = 4/3$

   $\cot \theta = -3/-4 = 3/4$

   $\sec \theta = 5/-3 = -5/3$

   $\csc \theta = 5/-4 = -5/4$

4. $\sin \theta = 8/17$

   $\cos \theta = -15/17$

   $\tan \theta = 8/-15 = -8/15$

   $\cot \theta = -15/8$

   $\sec \theta = 17/-15 = -17/15$

   $\csc \theta = 17/8$

5. 2

6. 3

7. 4

REFERENCE ANGLE

The *reference angle*, θ', for any angle, θ, in standard position is the smallest positive angle between the radius vector of θ and

X axis, such that $0° \leq \theta' \leq 90°$. In general, the reference angle θ is

$$\theta' = n(180°) \pm \theta$$

ere n is any integer. Expressed in an equivalent form

$$\theta' = n\pi \pm \theta$$

ere again n is any integer and $0 \leq \theta' \leq \pi/2$.

Refer to figure 4-10. If P is
y point on the radius vector,
erpendicular from P to the
nt A on the X axis forms a
ht triangle with sides OA,
, and OP. We call this
angle the *reference triangle*.
e relationship between θ, θ',
d the reference triangle in
ch quadrant is shown in
ure 4-10.

FUNCTIONS OF ANGLES IN ANY QUADRANT

In addition to the reference
angle, formulas are used for
termining the signs of the
nctions at any angle. These
e called *reduction formulas*.
is section shows the geo-
trical development of some
the most commonly used
luction formulas. In general,
duction formulas provide a
eans of reducing the func-
ns of any angle to an
uivalent expression for the
nction in terms of a positive
ute angle, θ. The reduction
rmulas can be used in the
ution of some trigonometric
ntities and in other appli-
ions requiring analysis of



Figure 4-10.—Reference triangles in each quadrant.

The function of θ and the reduction formulas of the functions of $180° - \theta$, $180° + \theta$, and $360° - \theta$ are summarized in the following paragraphs according to their respective quadrants.

QUADRANT I

Any angle in the first quadrant can be represented by θ; that is,

$$\sin \theta = \frac{y}{r}$$

$$\cos \theta = \frac{x}{r}$$

$$\tan \theta = \frac{y}{x}$$

$$\cot \theta = \frac{x}{y}$$

$$\sec \theta = \frac{r}{x}$$

$$\csc \theta = \frac{r}{y}$$

QUADRANT II

Any angle in the second quadrant can be represented by $180° - \theta$; that is,

$$\sin (180° - \theta) = \frac{y}{r} = \sin \theta$$

$$\cos (180° - \theta) = -\frac{x}{r} = -\cos \theta$$

$$\tan (180° - \theta) = -\frac{y}{x} = -\tan \theta$$

$$\cot (180° - \theta) = -\frac{x}{y} = -\cot \theta$$

$$\sec (180° - \theta) = -\frac{r}{x} = -\sec \theta$$

$$\csc (180° - \theta) = \frac{r}{y} = \csc \theta$$

EXAMPLE: Use a reduction formula and appendix III to find the cotangent of 112°.

SOLUTION: Since 112° is in the second quadrant, where

$$\cot (180° - \theta) = -\cot \theta$$

then

$$\begin{aligned} \cot 112° &= \cot (180° - 68°) \\ &= -\cot 68° \\ &= -0.40403 \end{aligned}$$

QUADRANT III

Any angle in the third quadrant can be represented by $180° + \theta$; that is,

$$\sin (180° + \theta) = -\frac{y}{r} = -\sin \theta$$

$$\cos (180° + \theta) = -\frac{x}{r} = -\cos \theta$$

$$\tan (180° + \theta) = \frac{y}{x} = \tan \theta$$

$$\cot (180° + \theta) = \frac{x}{y} = \cot \theta$$

$$\sec (180° + \theta) = -\frac{r}{x} = -\sec \theta$$

$$\csc (180° + \theta) = -\frac{r}{y} = -\csc \theta$$

EXAMPLE: Use a reduction formula and appendix II to find the sine of 220°.

SOLUTION: Since 220° is in the third quadrant, where

$$\sin (180° + \theta) = -\sin \theta$$

then

$$\sin 220° = \sin (180° + 40°)$$

$$= -\sin 40°$$

$$= -0.64279$$

QUADRANT IV

Any angle in the fourth quadrant can be represented by $360° - \theta$; that is,

$$\sin (360° - \theta) = -\frac{y}{r} = -\sin \theta$$

$$\cos (360° - \theta) = \frac{x}{r} = \cos \theta$$

$$\tan (360° - \theta) = -\frac{y}{x} = -\tan \theta$$

$$\cot (360° - \theta) = -\frac{x}{y} = -\cot \theta$$

$$\sec (360° - \theta) = \frac{r}{x} = \sec \theta$$

$$\csc (360° - \theta) = -\frac{r}{y} = -\csc \theta$$

EXAMPLE: Find cos 324°.

SOLUTION: Since

$$\cos (360° - \theta) = \cos \theta$$

then

$$\cos 324° = \cos (360° - 36°)$$

$$= \cos 36°$$

$$= 0.80902$$

FUNCTIONS OF NEGATIVE ANGLES

The following relationships enable us to change a function with a negative angle into the same function with a positive angle:

$$\sin(-\theta) = -\frac{y}{r} = -\sin\theta$$

$$\cos(-\theta) = \frac{x}{r} = \cos\theta$$

$$\tan(-\theta) = -\frac{y}{x} = -\tan\theta$$

$$\cot(-\theta) = -\frac{x}{y} = -\cot\theta$$

$$\sec(-\theta) = \frac{r}{x} = \sec\theta$$

$$\csc(-\theta) = -\frac{r}{y} = -\csc\theta$$

EXAMPLE: Find $\tan(-350°)$.

SOLUTION: Since

$$\tan(-\theta) = -\tan\theta$$

then

$$\tan(-350°) = -\tan 350°$$

and

$$\begin{aligned} -\tan 350° &= -\tan(360° - 10°) \\ &= -(-\tan 10°) \\ &= 0.17633 \end{aligned}$$

FUNCTIONS OF COTERMINAL ANGLES

For a *coterminal angle* in the form of

here n is any integer θ and is an integral multiple of θ', the igonometric functions of θ' are equal to those of θ. In other ords, θ is the remainder obtained by dividing θ' by 360, and is the number of times 360 will divide into θ'. Thus, we can nd the ratios of a coterminal angle greater than 360° by dividing ' by 360 and finding the functions of the remainder.

EXAMPLE: Find the cosine of $-2{,}080°$. Refer to fig. 4-11.)

SOLUTION: Divide 2,080 by 360.

$$\begin{array}{r} 5 \\ 360\overline{)2{,}080} \\ \underline{1{,}800} \\ 280 \end{array}$$

o,

$$\cos(-2{,}080°) = \cos(-280°)$$

nd

$$\begin{aligned} \cos(-280°) &= \cos(280°) \\ &= \cos(360° - 80°) \\ &= \cos 80° \\ &= 0.17365 \end{aligned}$$



Figure 4-11.—Coterminal angles o

PRACTICE PROBLEMS:

Use reduction formulas and appendixes II and III to find the values of the sine, cosine, and tangent of θ given the following angles:

1. 137°
2. 214°
3. 325°
4. $-70°$
5. 1,554°

ANSWERS:

1. $\sin 137° = \sin 43° = 0.68200$

   $\cos 137° = -\cos 43° = -0.73135$

   $\tan 137° = -\tan 43° = -0.93252$

2. $\sin 214° = -\sin 34° = -0.55919$

   $\cos 214° = -\cos 34° = -0.82904$

   $\tan 214° = \tan 34° = 0.67451$

3. $\sin 325° = -\sin 35° = -0.57358$

   $\cos 325° = \cos 35° = 0.81915$

   $\tan 325° = -\tan 35° = -0.70021$

4. $\sin (-70°) = -\sin 70° = -0.93969$

   $\cos (-70°) = \cos 70° = 0.34202$

   $\tan (-70°) = -\tan 70° = -2.74748$

5. $\sin 1{,}554° = \sin 114° = \sin 66° = 0.91355$

   $\cos 1{,}554° = \cos 114° = -\cos 66° = -0.40674$

   $\tan 1{,}554° = \tan 114° = -\tan 66° = -2.24604$

COFUNCTIONS AND COMPLEMENTARY ANGLES

Complementary angles are angles whose sum is 90°. Two trigonometric functions that have equal values for complementary angles are called *cofunctions*.

Inspect the triangle in figure 4-12. We will compare the six trigonometric functions of θ with the six trigonometric functions of $90° - \theta$.

Functions of θ	Functions of $90° - \theta$
$\sin \theta = \frac{y}{r}$	$\cos (90° - \theta) = \frac{y}{r}$
$\cos \theta = \frac{x}{r}$	$\sin (90° - \theta) = \frac{x}{r}$
$\tan \theta = \frac{y}{x}$	$\cot (90° - \theta) = \frac{y}{x}$
$\cot \theta = \frac{x}{y}$	$\tan (90° - \theta) = \frac{x}{y}$
$\sec \theta = \frac{r}{x}$	$\csc (90° - \theta) = \frac{r}{x}$
$\csc \theta = \frac{r}{y}$	$\sec (90° - \theta) = \frac{r}{y}$

Figure

We see from the above relationships that

$$\sin \theta = \cos (90° - \theta)$$

$$\cos \theta = \sin (90° - \theta)$$

$$\tan \theta = \cot (90° - \theta)$$

$$\cot \theta = \tan (90° - \theta)$$

$$\sec \theta = \csc (90° - \theta)$$

$$\csc \theta = \sec (90° - \theta)$$

Hence, a trigonometric function of an angle is equal to the confunction of its complement.

NOTE: These relationships may explain to you how the cosine, cotangent, and cosecant functions received their names.

The confunction principle accounts for the format of the tables of trigonometric functions in appendixes II and III. For example, in appendix II

$$\sin 21° 30' = 0.36650$$

and

$$\cos 68^\circ 30' = 0.36650$$

Notice that

$$21^\circ 30' + 68^\circ 30' = 90^\circ$$

PRACTICE PROBLEMS:

Express the following as a function of the complementary angle:

1. $\sin 27^\circ$
2. $\tan 38^\circ 17'$
3. $\csc 41^\circ$
4. $\cos 16^\circ 30' 22''$
5. $\sec 79^\circ 37' 16''$
6. $\cos 56^\circ$
7. $\cot 48^\circ$

ANSWERS:

1. $\cos 63^\circ$
2. $\cot 51^\circ 43'$
3. $\sec 49^\circ$
4. $\sin 73^\circ 29' 38''$
5. $\csc 10^\circ 22' 44''$
6. $\sin 34^\circ$
7. $\tan 42^\circ$

SPECIAL ANGLES

Two groups of angles are discussed in this section. The first group of angles is considered because the angles can be determined geometrically and are used frequently in problem solving. The second group is considered because the radius vectors of the angles fall on one of the coordinate axes, not in one of the quadrants.

FREQUENTLY USED ANGLES

As stated previously, the approximate values of the trigonometric functions for any angle can be read directly from tables or can be determined from tables by the use of the principles stated in this text. However, certain frequently used simple angles exist for which the exact function values are often used because these exact values can easily be determined geometrically. In the following paragraphs the geometrical determination of these functions is shown.

30°-60° Angles

The trigonometric functions of 30° and 60° can be determined geometrically. Construct an equilateral triangle with side lengths of 2 units, such as triangle OYA in figure 4-13. (The functions to be determined are not dependent on the lengths of the sides being 2 units; this size was selected for convenience.)

Drop a perpendicular from angle Y to the base of the triangle at point X. The right triangles YXO and YXA are formed by the perpendicular, which also bisects angle Y forming a 30° angle. Moreover, since side OA is 2 units long, then OX is 1 unit long and YX is $\sqrt{3}$ units long (using the Pythagorean theorem).

Figures 4-14 and 4-15 show a 30° and a 60° reference triangle, respectively. From these figures we can determine the trigonometric ratios of 30° and 60°, which are summarized in table 4-1.



Figure 4-13.—Geometr
30° and 60° rig



Figure 4-14.—30° r



Figure 4-15.—60° r

Table 4-1.—Trigonor
Special

θ	sin θ	cos θ	tan
30°	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$
60°	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$
45°	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1

NOTE: The values of the confunctions interchange since 30° and 60° are complementary angles. An easy way to recall the values of the functions of right triangles with 30° and 60° complementary angles is to remember that the ratio of the sides is always 1, 2, and $\sqrt{3}$, where the largest side value represents the length of the hypotenuse.

EXAMPLE: Find the six trigonometric functions of 300°.

SOLUTION: Referring to figure 4-16, 300° is in the fourth quadrant and its reference angle is 60°. Therefore,

$$\sin 300° = -\sqrt{3}/2$$

$$\cos 300° = 1/2$$

$$\tan 300° = -\sqrt{3}/1 = -\sqrt{3}$$

$$\cot 300° = 1/-\sqrt{3} = -\sqrt{3}/3$$

$$\sec 300° = 2/1 = 2$$

$$\csc 300° = 2/-\sqrt{3} = -2\sqrt{3}/3$$



Figure 4-16.—300° positi

45° Angles

Refer to figure 4-17. If one of the acute angles of the right triangle OXY is 45°, then the other acute angle is also 45°. Since triangle OXY is an isosceles triangle, then sides OX and XY are equal. If we let OX and XY be 1 unit long, then by the Pythagorean theorem, the length of OY is $\sqrt{2}$ units.

NOTE: This relationship is true of all 45° triangles and is not altered by the lengths of the legs. The ratio of the sides of right triangles with 45° complementary angles will always be 1, 1, and $\sqrt{2}$, where the largest value represents the length of the hypotenuse.



Figure 4-17.—Geometrical c right triang

Figure 4-18 shows a 45° reference triangle. From is figure we can determine the trigonometric ratios 45°, which are also summarized in table 4-1.

EXAMPLE: Find the six trigonometric functions 135°.

SOLUTION: Referring to figure 4-19, 135° is in e second quadrant and its reference angle is 45°. herefore,

$$\sin 135° = 1/\sqrt{2} = \sqrt{2}/2$$

$$\cos 135° = -1/\sqrt{2} = -\sqrt{2}/2$$

$$\tan 135° = 1/-1 = -1$$

$$\cot 135° = -1/1 = -1$$

$$\sec 135° = \sqrt{2}/-1 = -\sqrt{2}$$

$$\csc 135° = \sqrt{2}/1 = \sqrt{2}$$



Figure 4-18.—45° reference t

UADRANTAL ANGLES

An angle whose terminal side lies on a coordinate is when the angle is in standard position is a *uadrantal angle*. Angles of 0°, ±90°, ±180°, and 270° are quadrantal angles.

The trigonometric functions of the quadrantal gles are defined in the same manner as before, cept for the restriction that a function is undefined hen the denominator of the ratio is zero.

To derive the functions of the quandrantal angles, e choose points on the terminal sides, where $r = 1$, shown in figure 4-20. Then either x or y is zero, d the other coordinate is either positive or negative 1.



Figure 4-19.—135° angle in position.



Figure 4-20.—Functions of quadrantal angles.

Table 4-2.—Functions of Quadrantal Angles

θ		$\sin\theta$	$\cos\theta$	$\tan\theta$	$\cot\theta$	$\sec\theta$	$\csc\theta$
Deg.	Rad.						
0°	0	0	1	0	undefined	1	undefined
90°	$\frac{\pi}{2}$	1	0	undefined	0	undefined	1
180°	π	0	-1	0	undefined	-1	undefined
270°	$\frac{3\pi}{2}$	-1	0	undefined	0	undefined	-1

Consider view C of figure 4-20 in which $\theta = 180°$. For the point $P(-1,0)$ and $r = 1$, we have

$$\sin 180° = 0/1 = 0$$

$$\cos 180° = -1/1 = -1$$

$$\tan 180° = 0/-1 = 0$$

$$\cot 180° = -1/0 \text{ (undefined)}$$

$$\sec 180° = 1/-1 = -1$$

$$\csc 180° = 1/0 \text{ (undefined)}$$

The values of the functions of the other quadrantal angles can be found by a similar procedure and are summarized in table 4-2.

EXAMPLE: Determine the six trigonometric functions of 990°.

SOLUTION: Referring to figure 4-21, we see that 990° lies on the same quadrantal axes as 270°. Therefore, for $P(0,-1)$ and $r = 1$, we have

$$\sin 990° = -1/1 = -1$$

$$\cos 990° = 0/1 = 0$$

$$\tan 990° = -1/0 \text{ (undefined)}$$

$$\cot 990° = 0/-1 = 0$$

$$\sec 990° = 1/0 \text{ (undefined)}$$

$$\csc 990° = 1/-1 = -1$$



Figure 4-21.—990° a

PRACTICE PROBLEMS:

Without using the appendixes, determine the trigonometric functions of problems 1 through 5.

1. $\theta = 210°$

2. $\theta = 360°$

3. $\theta = 585°$

4. $\theta = -180°$

5. $\theta = -315°$

Without using the appendixes, evaluate problems 6 through 8. [Note that $\sin^2 \theta = (\sin \theta)^2$.]

6. $\sin^2 150° + \cos^2 150°$

7. $2 \sin 120° \cos 120°$

8. $\cos^2 135° - \sin^2 135°$

ANSWERS:

1. $\sin 210° = -1/2$

   $\cos 210° = -\sqrt{3}/2$

   $\tan 210° = -1/-\sqrt{3} = \sqrt{3}/3$

   $\cot 210° = -\sqrt{3}/-1 = \sqrt{3}$

   $\sec 210° = 2/-\sqrt{3} = -2\sqrt{3}/3$

   $\csc 210° = 2/-1 = -2$

2. $\sin 360° = 0/1 = 0$

   $\cos 360° = 1/1 = 1$

   $\tan 360° = 0/1 = 0$

   $\cot 360° = 1/0$ (undefined)

   $\sec 360° = 1/1 = 1$

   $\csc 360° = 1/0$ (undefined)

3. $\sin 585° = -1/\sqrt{2} = -\sqrt{2}/2$

   $\cos 585° = -1/\sqrt{2} = -\sqrt{2}/2$

   $\tan 585° = -1/-1 = 1$

   $\cot 585° = -1/-1 = 1$

   $\sec 585° = \sqrt{2}/-1 = -\sqrt{2}$

   $\csc 585° = \sqrt{2}/-1 = -\sqrt{2}$

4. $\sin (-180°) = 0/1 = 0$

   $\cos (-180°) = -1/1 = -1$

   $\tan (-180°) = 0/-1 = 0$

   $\cot (-180°) = -1/0$ (undefined)

   $\sec (-180°) = 1/-1 = -1$

   $\csc (-180°) = 1/0$ (undefined)

5. $\sin (-315°) = 1/\sqrt{2} = \sqrt{2}/2$

   $\cos (-315°) = 1/\sqrt{2} = \sqrt{2}/2$

   $\tan (-315°) = 1/1 = 1$

   $\cot (-315°) = 1/1 = 1$

   $\sec (-315°) = \sqrt{2}/1 = \sqrt{2}$

   $\csc (-315°) = \sqrt{2}/1 = \sqrt{2}$

6. $(1/2)^2 + (-\sqrt{3}/2)^2 = 1$

7. $2(\sqrt{3}/2)(-1/2) = -\sqrt{3}/2$

8. $(-1/\sqrt{2})^2 - (1/\sqrt{2})^2 = 0$



Figure 4-22.—Graph of the sine function.

PERIODS OF THE TRIGONOMETRIC FUNCTIONS

A trigonometric function of an angle is not changed in value when the angle is changed by any multiple of 360° or 2π radians. For this reason the functions are said to be *periodic.*

In the following paragraphs, the graphs of the sine, cosine, and tangent functions are discussed.

GRAPH OF THE SINE FUNCTION

Figure 4-22 shows two periods of the sine function. The graph shows that the value of the sine function varies between $+1$ and -1 and never goes beyond these limits as the angle varies. The graph also shows that the sine function increases from 0 at 0° or 0 radians to a maximum value of $+1$ at 90° or $\pi/2$ radians. It decreases back to 0 at 180° or π radians and continues to decrease to a minimum value of -1 at 270° or $3\pi/2$ radians. It then increases to a value of 0 at 360° or 2π radians. If we extend the graph (in either direction), the sine function will continue to repeat itself every 360° or 2π radians. Therefore, *the period of the sine function is 360° or 2π radians.*

GRAPH OF THE COSINE FUNCTION

The cosine function also has a period of 360° or 2π radians. Figure 4-23 shows two periods of the cosine function. The range



Figure 4-23.—Graph of the cosine function.

of the values the cosine function takes on also lies between $+1$ and -1. However, as seen on the graph, the cosine function decreases from 1 at $0°$ or 0 radians to 0 at $90°$ or $\pi/2$ radians and continues to decrease to a minimum value of -1 at $180°$ or π radians. It then increases to 0 at $270°$ or $3\pi/2$ radians and continues to increase to a maximum value of $+1$ at $360°$ or 2π radians. This completes one period of the cosine function.

GRAPH OF THE TANGENT FUNCTION

Figure 4-24 shows the graph of the tangent function from 0 radians to 2π radians. Notice that the tangent function is 0 at $0°$ or 0 radians and increases to positive infinity (without bounds) between $0°$ and $90°$ or 0 radians and $\pi/2$ radians. Remember that the tangent function is undefined for $90° + n(180°)$ or $\pi/2 + n\pi$, where n is any integer. The dashed vertical lines in figure 4-24 represent the undefined points. The tangent function increases from negative infinity to 0 between $90°$ and $180°$ or $\pi/2$ radians and π radians. At $180°$ or π radians, the tangent function is 0. The function continues to increase from 0 to positive infinity between $180°$ and $270°$ or π radians and $3\pi/2$ radians. Between $270°$ and $360°$ or $3\pi/2$ radians and 2π, it again increases from negative infinity to 0 at $360°$ or 2π radians. If we extend the graph (in either direction), the curve will repeat itself every $180°$ or π radians. Therefore, *the period of the tangent function is $180°$ or π radians.*



Figure 4-24.—Graph of the tange

EXAMPLE: Using the graphs in figures 4-22 through 4-24, determine the values of θ, where $\sin\theta$ and $\tan\theta$ increase together, if $0 \leq \theta \leq \pi$.

SOLUTION: In figure 4-22 the sine function increases between 0 and $\pi/2$ radians for the interval of $0 \leq \theta \leq \pi$. (The sine function does not increase or decrease at points 0 or $\pi/2$.) In figure 4-24 the tangent function also increases between 0 and $\pi/2$ radians for the interval of $0 \leq \theta \leq \pi$. (The tangent function does not increase or decrease at 0 and is undefined at $\pi/2$.) Therefore, the values of θ, where $\sin\theta$ and $\tan\theta$ increase together, are $0 < \theta < \pi/2$.

PRACTICE PROBLEMS:

Use the graphs in figures 4-22 through 4-24 to answer the following problems (use appendixes II and III to verify your answers):

1. For what values of θ does $\cos \theta$ increase if $0 \leq \theta \leq \pi$?

2. For what values of θ do $\sin \theta$ and $\cos \theta$ decrease together if $0 \leq \theta \leq 2\pi$?

3. For what values of θ do $\cos \theta$ and $\tan \theta$ increase together if $\pi/2 \leq \theta \leq 3\pi/2$?

4. For what values of θ do $\sin \theta$, $\cos \theta$, and $\tan \theta$ increase together if $0 \leq \theta \leq 2\pi$?

ANSWERS:

1. None

2. $\pi/2 < \theta < \pi$

3. $\pi < \theta < 3\pi/2$

4. $3\pi/2 < \theta < 2\pi$

SUMMARY

The following are the major topics covered in this chapter:

1. **Angles in standard position:** An angle in standard position on a rectangular coordinate system has its vertex at the origin, its initial side lying along the X axis, and its terminal side lying in any of the quadrants or on one of the axes.

2. **Coterminal angles**: When two or more angles in standard position have their terminal sides located at the same position, they are said to be *coterminal.*

   For any general angle θ measured in degrees, any angle ϕ coterminal with θ can be found by

   $$\phi = \theta + n(360°)$$

   where n is any integer.

3. **Definitions of the trigonometric functions:**

   $$\sin \theta = \frac{y}{r} = \frac{\text{ordinate}}{\text{length of radius}}$$

   $$\cos \theta = \frac{x}{r} = \frac{\text{abscissa}}{\text{length of radius}}$$

   $$\tan \theta = \frac{y}{x} = \frac{\text{ordinate}}{\text{abscissa}}$$

   $$\cot \theta = \frac{x}{y} = \frac{\text{abscissa}}{\text{ordinate}}$$

   $$\sec \theta = \frac{r}{x} = \frac{\text{length of radius}}{\text{abscissa}}$$

   $$\csc \theta = \frac{r}{y} = \frac{\text{length of radius}}{\text{ordinate}}$$

4. **Signs of the trigonometric ratios in the quadrant system**: All the trigonometric ratios are positive for angles in the first quadrant. Only the sine and cosecant ratios are positive in the second quadrant. Only the tangent and cotangent ratios are positive in the third quadrant. Only the cosine and secant ratios are positive in the fourth quadrant.

5. **Reference angle**: The *reference angle*, θ', for any angle, θ, in standard position is the smallest positive angle between the radius vector of θ and the X axis, such that $0° \leq \theta' \leq 90°$. In general, for any integer n,

$$\theta' = n(180°) \pm \theta$$

or

$$\theta' = n\pi \pm \theta$$

where $0 \leq \theta' \leq \pi/2$.

6. **Reference triangle:** The right triangle formed from the reference angle when you connect a point on the radius vector of the reference angle perpendicular to the X axis is called the *reference triangle.*

7. **Reduction formulas:** *Reduction formulas* are formulas used to determine the signs of the functions of any angle. They provide a means of reducing the functions of any angle to an equivalent expression for the function in terms of a positive acute angle.

8. **Quadrant I angles**: An angle in quadrant I is represented by θ.

$$\sin \theta = y/r$$

$$\cos \theta = x/r$$

$$\tan \theta = y/x$$

$$\cot \theta = x/y$$

$$\sec \theta = r/x$$

$$\csc \theta = r/y$$

9. **Quadrant II angles**: An angle in quadrant II is represented by $180 - \theta$.

$$\sin (180° - \theta) = \sin \theta$$

$$\cos (180° - \theta) = -\cos \theta$$

$$\tan (180° - \theta) = -\tan \theta$$

$$\cot (180° - \theta) = -\cot \theta$$

$$\sec (180° - \theta) = -\sec \theta$$

$$\csc (180° - \theta) = \csc \theta$$

10. **Quadrant III angles:** An angle in quadrant III is represented by $180° + \theta$.

$$\sin (180° + \theta) = -\sin \theta$$

$$\cos (180° + \theta) = -\cos \theta$$

$$\tan (180° + \theta) = \tan \theta$$

$$\cot (180° + \theta) = \cot \theta$$

$$\sec (180° + \theta) = -\sec \theta$$

$$\csc (180° + \theta) = -\csc \theta$$

11. **Quadrant IV angles:** An angle in quadrant IV is represented by $360° - \theta$.

$$\sin (360° - \theta) = -\sin \theta$$

$$\cos (360° - \theta) = \cos \theta$$

$$\tan (360° - \theta) = -\tan \theta$$

$$\cot (360° - \theta) = -\cot \theta$$

$$\sec (360° - \theta) = \sec \theta$$

$$\csc (360° - \theta) = -\csc \theta$$

12. **Functions of negative angles:**

$$\sin (-\theta) = -\sin \theta$$

$$\cos (-\theta) = \cos \theta$$

$$\tan (-\theta) = -\tan \theta$$

$$\cot (-\theta) = -\cot \theta$$

$$\sec (-\theta) = \sec \theta$$

$$\csc (-\theta) = -\csc \theta$$

13. **Functions of coterminal angles:** For a coterminal angle in the form of

$$\theta' = n(360°) + \theta$$

where n is any integer and θ is an integral multiple of θ', the trigonometric functions of θ' are equal to those of θ.

14. **Cofunctions and complementary angles:** *Complementary angles* are angles whose sum is 90°. Two trigonometric functions that have equal values for complementary angles are called *cofunctions.*

$$\sin \theta = \cos (90° - \theta)$$

$$\cos \theta = \sin (90° - \theta)$$

$$\tan \theta = \cot (90° - \theta)$$

$$\cot \theta = \tan (90° - \theta)$$

$$\sec \theta = \csc (90° - \theta)$$

$$\csc \theta = \sec (90° - \theta)$$

15. **Frequently used angles:** The trigonometric functions of 30°, 60°, and 45° can be determined geometrically. The trigonometric ratios corresponding to these functions are summarized in table 4-1.

16. **Quadrantal angles:** An angle whose terminal side lies on a coordinate axis when the angle is in standard position is a *quadrantal angle.* The trigonometric ratios corresponding to the functions of the quadrantal angles are summarized in table 4-2.

17. **Periods of the trigonometric functions:** A trigonometric function of an angle is not changed in value when the angle is changed by any multiple of 360° or 2π radians. For this reason the functions are said to be *periodic.* The periods of the sine and cosine functions are 360° or 2π radians. The period of the tangent function is 180° or π radians.

ADDITIONAL PRACTICE PROBLEMS

1. Are the angles $840°$, $-240°$, and $600°$ coterminal?

2. Find the sine, cosine, and tangent of the angle θ whose radius vector passes through the point $P(\sqrt{5}, \sqrt{11})$.

3. Find the six trigonometric functions of θ if $\csc \theta = -37/35$ and $\tan \theta > 0$.

4. Find the sine, cosine, and tangent of $-4{,}010°$.

5. Express $\csc 87° 23' 13''$ as a function of its complementary angle.

6. Without using the appendixes, evaluate $\sec^2(-135°) + \cot^2(-690°) + \csc^2(-600°)$.

7. Without using the appendixes, find the six trigonometric functions of $-3{,}510°$.

8. For what values of θ do $\cos \theta$ and $\tan \theta$ both increase and $\sin \theta$ decrease together if $0 \leq \theta \leq 2\pi$?

ANSWERS TO ADDITIONAL PRACTICE PROBLEMS

1. No

2. $\sin \theta = \sqrt{11}/4$

   $\cos \theta = \sqrt{5}/4$

   $\tan \theta = \sqrt{11}/\sqrt{5} = \sqrt{55}/5$

3. $\sin \theta = -35/37$

   $\cos \theta = -12/37$

   $\tan \theta = 35/12$

   $\cot \theta = 12/35$

   $\sec \theta = -37/12$

   $\csc \theta = -37/35$

4. $\sin \theta = -0.76604$

   $\cos \theta = 0.64279$

   $\tan \theta = -1.19175$

5. sec 2° 36′ 47″

6. 6 1/3

7. $\sin \theta = 1$

   $\cos \theta = 0$

   $\tan \theta$ is undefined

   $\cot \theta = 0$

   $\sec \theta$ is undefined

   $\csc \theta = 1$

8. $\pi < \theta < 3\pi/2$

CHAPTER 5

OBLIQUE TRIANGLES

LEARNING OBJECTIVES

Upon completion of this chapter, you should be able to do the following:

1. Apply the Law of Sines to solve oblique triangles given one side and two angles or two sides and an angle opposite one of them.

2. Apply the Law of Cosines to solve oblique triangles given two sides and the included angle or all three sides.

3. Find the area of an oblique triangle.

INTRODUCTION

The two previous chapters primarily dealt with properties of right triangles in solving trigonometric measurements and functions. In this chapter we will apply properties of oblique triangles in solving trigonometric measurements and functions. *Oblique triangles* are triangles containing no right angles. Oblique triangles are made up of either three acute angles or two acute angles and one obtuse angle. *Acute angles* have measures between 0° and 90°. *Obtuse angles* have measures between 90° and 180°.

In *Mathematics,* Volume 1, a method for solving problems involving oblique triangles was introduced. The method employed the procedures of dividing the original triangle into two or more right triangles and using the properties of right triangles in problem solving.

This chapter develops two methods or laws dealing directly with oblique triangles. The methods consider the parts of the

ngle that are given. The four standard cases for solving ique triangles are as follows:

Case 1. One side and two angles

Case 2. Two sides and an angle opposite one of them

Case 3. Two sides and the included angle

Case 4. All three sides

Also included in this chapter are problems concerning the area a triangle, which combine the area formula of plane geometry h trigonometric properties.

METHODS OF SOLVING OBLIQUE TRIANGLES

This section is concerned with the development and proofs of Law of Sines and the Law of Cosines. The four standard cases solving oblique triangles use applications of these laws.

W OF SINES

Law of Sines. *The lengths of the sides of any triangle are portional to the sines of their opposite angles*; that is,

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

PROOF: Refer to the ique triangle shown in ure 5-1, view A. Let h the length of the pendicular from angle to the side opposite le A. Considering the right triangles formed h, we obtain



A



B

Figure 5-1.—Development of Law of Sines.

$$B = \frac{h}{c} \text{ or } h = c \sin B$$

$$C = \frac{h}{b} \text{ or } h = b \sin C$$

Equating these two values of h, we have

$$c \sin B = b \sin C$$

or in an equivalent form, we have

$$\frac{c}{\sin C} = \frac{b}{\sin B}$$

Now, if we redraw the oblique triangle in figure 5-1, view A, by extending the length of side c until it forms a right angle (is perpendicular) with a line, h', from angle C (see fig. 5-1, view B), then from the newly formed triangle, we obtain

$$\sin B = \frac{h'}{a} \text{ or } h' = a \sin B$$

and

$$\sin (180° - A) = \frac{h'}{b} \text{ or } h' = b \sin (180° - A)$$

Since

$$\sin (180° - A) = \sin A$$

then by substituting $\sin A$ for $\sin (180° - A)$ and equating values of h', we have

$$a \sin B = b \sin A$$

or in an equivalent form

$$\frac{a}{\sin A} = \frac{b}{\sin B}$$

But

$$\frac{b}{\sin B} = \frac{c}{\sin C}$$

Therefore,

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

Case 1. One Side and Two Angles

When one side and two angles of a triangle are given, the third angle can be found since the sum of the angles equals 180°; that is, $A + B + C = 180°$. Then the Law of Sines can be used to find the two remaining sides.

EXAMPLE: Solve the remaining parts of triangle ABC, given $c = 5$, $B = 30°$, and $C = 97° 30'$. Give side accuracy to one decimal place.

SOLUTION: Refer to figure 5-2. Since

$$A + B + C = 180°$$

then

$$A + 30° + 97° 30' = 180°$$

$$A = 180° - 30° - 97° 30'$$

$$= 52° 30'$$



Figure 5-2.—Case 1. One side and

By the Law of Sines,

$$\frac{a}{\sin A} = \frac{c}{\sin C}$$

we obtain

$$\frac{a}{\sin 52° 30'} = \frac{5}{\sin 97° 30'}$$

$$a = \frac{5 \sin 52° 30'}{\sin 97° 30'}$$

$$= \frac{5(0.79335)}{0.99144}$$

$$= 4.0$$

We will use the Law of Sines again to solve for the length of side b:

$$\frac{b}{\sin B} = \frac{c}{\sin C}$$

o,

$$\frac{b}{\sin 30^\circ} = \frac{5}{\sin 97^\circ 30'}$$

$$b = \frac{5 \sin 30^\circ}{\sin 97^\circ 30'}$$

$$= \frac{5(0.50000)}{0.99144}$$

$$= 2.5$$

EXAMPLE: The base of flagpole standing vertically on a hill s inclined at an angle of 15° with the horizontal. A man standing 200 feet downhill from the base of the flagpole notes that his line of sight to the top of the flagpole makes an angle of 40° with the horizontal. How high, to the nearest foot, is the flagpole?

SOLUTION: Refer to figure 5-3. In triangle *ABC* we find

$$A = 40^\circ - 15^\circ$$

$$= 25^\circ$$

From right triangle *ADC* we find

$$C = 180^\circ - 40^\circ - 90^\circ$$

$$= 50^\circ$$

Applying the Law of Sines, we obtain

$$\frac{a}{\sin 25^\circ} = \frac{200}{\sin 50^\circ}$$

$$a = \frac{200 \sin 25^\circ}{\sin 50^\circ}$$

$$= \frac{200(0.42262)}{0.76604}$$

$$= 110 \text{ feet}$$



Figure 5-3.—Case 1. Flagpole

Case 2. Two Sides and an Angle Opposite One of Them

Case 2 is sometimes referred to as the *ambiguous case* since two triangles, one triangle, or no triangle



A



B



C



D



E



F

Figure 5-4.—Case 2. Two sides and an angle opposite one of them.

may result from data given in this form. Consider triangle ABC in figure 5-4. Assuming we are given angle A and sides a and c, the following situations may exist:

For acute angle A:

1. If $a < c$ and sin $C < 1$, then two possible triangles exist; one triangle comprises the acute angle C and the other triangle comprises the obtuse angle $C' = 180° - C$. See figure 5-4, view A.

2. If $a \geq c$, then one triangle exists. See figure 5-4, view B.

3. If $a < c$ and $\sin C = 1$, then one right triangle exists. See figure 5-4, view C.

4. If $a < c$ and $\sin C > 1$, then no triangle is determined. See figure 5-4, view D. (This should be obvious since in the previous chapter we learned that the sine of an angle may have values only between 0 and 1.)

For obtuse angle A:

1. If $a > c$, then one triangle exists. See figure 5-4, view E.

2. If $a \leq c$, then no triangle is determined. See figure 5-4, view F.

When two sides and an angle opposite one of them are given, we can solve for the remaining parts of the triangle using the Law of Sines. Sketches can be helpful.

EXAMPLE: Solve the triangle or triangles if they exist, given $B = 45°$, $b = 3$, and $c = 7$.

SOLUTION: Using the Law of Sines,

$$\frac{b}{\sin B} = \frac{c}{\sin C}$$

we have

$$\frac{3}{\sin 45°} = \frac{7}{\sin C}$$

$$\sin C = \frac{7 \sin 45°}{3}$$

$$= \frac{7(0.70711)}{3}$$

$$= 1.64992$$



Figure 5-5.—Case 2. Acute a
$b < c$ and sin $C >$

Since the sine of an angle cannot exceed 1, then we conclude that no triangle exists. Refer to figure 5-5. Notice that $b < c$ and $\sin C > 1$.

EXAMPLE: Solve the triangle or triangles if they exist, given $A = 22°$, $a = 5.4$, and $c = 14$. Give angle accuracy to the nearest minute and side accuracy to one decimal place.

SOLUTION: Using the Law of Sines,

$$\frac{a}{\sin A} = \frac{c}{\sin C}$$

ve obtain

$$\frac{5.4}{\sin 22^\circ} = \frac{14}{\sin C}$$

$$\sin C = \frac{14 \sin 22^\circ}{5.4}$$

$$= \frac{14(0.37461)}{5.4}$$

$$= 0.97121$$

$$C = 76^\circ\ 13'$$

Since the side opposite the known angle s smaller than the other given side, that s, $a < c$, and $\sin C < 1$, then two possible riangles exist. One triangle is ABC and he other is $AB'C'$. Refer to figure 5-6. Hence,



Figure 5-6.—Case 2. Acute angle A with $a < c$ an

$$C' = 180^\circ - C$$

$$= 180^\circ - 76^\circ\ 13'$$

$$= 103^\circ\ 47'$$

Solving triangle ABC first, we find angle B by

$$B = 180^\circ - (A + C)$$

$$= 180^\circ - (22^\circ + 76^\circ\ 13')$$

$$= 81^\circ\ 47'$$

and by the Law of Sines,

$$\frac{a}{\sin A} = \frac{b}{\sin B}$$

we find the length of side b to be

$$\frac{5.4}{\sin 22^\circ} = \frac{b}{\sin 81^\circ 47'}$$

$$b = \frac{5.4 \sin 81^\circ 47'}{\sin 22^\circ}$$

$$= \frac{5.4(0.98973)}{0.37461}$$

$$= 14.3$$

Now solving triangle $AB'C'$, we find angle B' by

$$B' = 180^\circ - (A + C')$$

$$= 180^\circ - (22^\circ + 103^\circ 47')$$

$$= 54^\circ 13'$$

and by the Law of Sines,

$$\frac{a}{\sin A} = \frac{b'}{\sin B'}$$

we find the length of side b' to be

$$\frac{5.4}{\sin 22^\circ} = \frac{b'}{\sin 54^\circ 13'}$$

$$b' = \frac{5.4 \sin 54^\circ 13'}{\sin 22^\circ}$$

$$= \frac{5.4(0.81123)}{0.37461}$$

$$= 11.7$$

EXAMPLE: Solve the triangle if it exists, given $C = 125^\circ 48'$, $b = 41.8$, and $c = 56.2$. Give angle accuracy to the nearest minute and side accuracy to two decimal places.

SOLUTION: Since the given angle, C, is obtuse and the side opposite the given angle is larger than the other given side, that is, $c > b$, then one triangle exists. Refer to figure 5-7. By the Law of Sines,

$$\frac{b}{\sin B} = \frac{c}{\sin C}$$



Figure 5-7.—Case 2. Obtuse a

we get

$$\frac{41.8}{\sin B} = \frac{56.2}{\sin 125^\circ 48'}$$

$$\sin B = \frac{41.8 \sin 125^\circ 48'}{56.2}$$

$$= \frac{41.8(0.81106)}{56.2}$$

$$= 0.60324$$

$$B = 37^\circ 6'$$

Additionally,

$$A = 180^\circ - (B + C)$$

$$= 180^\circ - (37^\circ 6' + 125^\circ 48')$$

$$= 17^\circ 6'$$

and by the Law of Sines,

$$\frac{a}{\sin A} = \frac{c}{\sin C}$$

we find the length of side a to be

$$\frac{a}{\sin 17^\circ 6'} = \frac{56.2}{\sin 125^\circ 48'}$$

$$a = \frac{56.2 \sin 17^\circ 6'}{\sin 125^\circ 48'}$$

$$= \frac{56.2(0.29404)}{0.81106}$$

$$= 20.37$$

PRACTICE PROBLEMS:

Use the Law of Sines to solve the remaining parts of triangle ABC given the following parts (give angle accuracy to the nearest minute and side accuracy to one decimal place):

1. $A = 59° 36'$, $B = 48° 14'$, and $c = 86.4$

2. $A = 98° 8'$, $C = 25° 25'$, and $b = 2.1$

3. $B = 30° 30'$, $a = 10$, and $b = 10$

4. $C = 100° 21'$, $a = 4.2$, and $c = 3.2$

ANSWERS:

1. $C = 72° 10'$

   $a = 78.3$

   $b = 67.7$

2. $B = 56° 27'$

   $a = 2.5$

   $c = 1.1$

3. $A = 30° 30'$

   $C = 119°$

   $c = 17.2$

4. No solution; C is obtuse and $c \leq a$.

LAW OF COSINES

Law of Cosines. *In a triangle, the square of any side is equal to the sum of the squares of the other two sides minus twice the*

product of the same two sides multiplied by the cosine of the angle between them; that is,

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$b^2 = a^2 + c^2 - 2ac \cos B$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$



Figure 5-8.—Developmer Cosines.

PROOF: Refer to the oblique triangle shown in figure 5-8. Let h be the length of the perpendicular from angle B to the side opposite angle B.

NOTE: $b = b + 0$

$= b + (x - x)$

$= x + (b - x)$

Considering right triangle ADB formed by h, we obtain

$$\cos A = \frac{x}{c} \text{ or } x = c \cos A$$

and

$$h^2 = c^2 - x^2$$

Substituting the value of x into the last equation gives

$$h^2 = c^2 - c^2 \cos^2 A$$

Considering right triangle CDB, we obtain

$$h^2 = a^2 - (b - x)^2$$

$$= a^2 - b^2 + 2bx - x^2$$

Substituting the value x in the last equation for h^2 gives

$$h^2 = a^2 - b^2 + 2bc \cos A - c^2 \cos^2 A$$

Equating the two values of h^2 gives

$$c^2 - c^2 \cos^2 A = a^2 - b^2 + 2bc \cos A - c^2 \cos^2 A$$

Therefore, rearranging and canceling terms gives

$$a^2 = b^2 + c^2 - 2bc \cos A$$

The same procedure can be applied to derive all three forms of the Law of Cosines, which are

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$b^2 = a^2 + c^2 - 2ac \cos B$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$

Case 3. Two Sides and the Included Angle

When two sides and the angle between them are given, we can solve for the remaining parts of the triangle using the Law of Cosines. First, the unknown side is determined; then the two other angles are determined.

EXAMPLE: Solve for the remaining parts of triangle ABC, given $b = 7$, $c = 5$, and $A = 19°$. Give angle accuracy to the nearest minute and side accuracy to one decimal place.

SOLUTION: Refer to figure 5-9. First, find the length of the unknown side using the Law of Cosines,

$$a^2 = b^2 + c^2 - 2bc \cos A$$

Hence,

$$\begin{aligned} a^2 &= 7^2 + 5^2 - 2(7)(5) \cos 19° \\ &= 49 + 25 - 70(0.94552) \\ &= 7.8136 \\ a &= \sqrt{7.8136} \\ &= 2.8 \end{aligned}$$



Figure 5-9.—Case 3. Two included angle

Next, compute the remaining angles using a rearrangement of the Law of Cosines:

$$\begin{aligned} \cos B &= \frac{a^2 + c^2 - b^2}{2ac} \\ &= \frac{(2.8)^2 + (5)^2 - (7)^2}{2(2.8)(5)} \\ &= \frac{7.84 + 25 - 49}{28} \\ &= -0.57714 \\ &= -\cos 54° 45' \end{aligned}$$

Angle B is an obtuse angle since cos B is negative. Therefore,

$$B = 125^\circ\ 15'$$

and

$$\cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

$$= \frac{(2.8)^2 + (7)^2 - (5)^2}{2(2.8)(7)}$$

$$= \frac{7.84 + 49 - 25}{39.2}$$

$$= 0.81224$$

$$C = 35^\circ\ 41'$$

NOTE: Since we are solving angles to the nearest minute, the sum of the angles may not equal exactly 180°.

EXAMPLE: Two points, A and B, are separated by a pond. The distance from A to a third point, C, is 10.2 feet; the distance from C to B is 13.8 feet; and angle C is 52° 40′. Find the distance from A to B to two decimal places and angles A and B to the nearest minute.

SOLUTION: We will first find the distance from point A to point B. Using the Law of Cosines, we find that

$$c^2 = a^2 + b^2 - 2ab \cos C$$

$$= (13.8)^2 + (10.2)^2 - 2(13.8)(10.2) \cos 52^\circ\ 40'$$

$$= 190.44 + 104.04 - (281.52)(0.60645)$$

$$= 123.75$$

$$c = \sqrt{123.75}$$

$$= 11.12 \text{ feet}$$

Now we will find angles A and B using the Law of Cosines:

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$= \frac{(10.2)^2 + (11.12)^2 - (13.8)^2}{2(10.2)(11.12)}$$

$$= 0.16423$$

$$A = 80^\circ\ 33'$$

and

$$\cos B = \frac{a^2 + c^2 - b^2}{2ac}$$

$$= \frac{(13.8)^2 + (11.12)^2 - (10.2)^2}{2(13.8)(11.12)}$$

$$= 0.68441$$

$$B = 46^\circ\ 49'$$

Case 4. All Three Sides

The Law of Cosines can also be used to find the size of the angles of a triangle when the length of all three sides are given.

EXAMPLE: Find the measure of each angle (to the nearest minute) of a triangle having sides $a = 7$, $b = 13$, and $c = 14$.

SOLUTION:

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$= \frac{(13)^2 + (14)^2 - (7)^2}{2(13)(14)}$$

$$= 0.86813$$

$$A = 29^\circ\ 45'$$

$$\cos B = \frac{a^2 + c^2 - b^2}{2ac}$$

$$= \frac{(7)^2 + (14)^2 - (13)^2}{2(7)(14)}$$

$$= 0.38776$$

$$B = 67^\circ\ 11'$$

and

$$\cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

$$= \frac{(7)^2 + (13)^2 - (14)^2}{2(7)(13)}$$

$$= 0.12088$$

$$C = 83^\circ\ 3'$$

EXAMPLE: A triangular plot of ground measures 50 meters by 70 meters by 90 meters. Find, to the nearest minute, the size of the angle, A, opposite the longest side.

SOLUTION:

$$\cos A = \frac{(50)^2 + (70)^2 - (90)^2}{2(50)(70)}$$

$$= -0.10000$$

$$= -\cos 84^\circ 16'$$

$$A = 95^\circ 44'$$

PRACTICE PROBLEMS:

Use the Law of Cosines to solve the remaining parts of triangle ABC given the following parts (give angle accuracy to the nearest minute and side accuracy to two decimal places):

1. $a = 54.2$, $c = 83.4$, and $B = 111^\circ 11'$
2. $b = 6.6$, $c = 6.6$, and $A = 60^\circ$
3. $a = 22.2$, $b = 33.3$, and $c = 44.4$
4. $a = 15.6$, $b = 16.7$, and $c = 17.8$

ANSWERS:

1. $b = 114.72$
   $A = 26^\circ 8'$
   $C = 42^\circ 41'$
2. $a = 6.6$
   $B = 60^\circ$
   $C = 60^\circ$
3. $A = 28^\circ 57'$
   $B = 46^\circ 34'$
   $C = 104^\circ 29'$
4. $A = 53^\circ 39'$
   $B = 59^\circ 34'$
   $C = 66^\circ 47'$

AREA FORMULAS

In this section two formulas for finding the area of a triangle will be developed. Recall from plane geometry that the area of a triangle is found by the formula

$$\text{area} = \frac{1}{2}bh$$

where b is any side of the triangle and h is the altitude drawn to that side. While this is a useful formula, it is not a practical one. With the help of trigonometry, we can derive more practical formulas for the area of a triangle.

Consider the triangle in figure 5-8. The length of the altitude is found to be

$$h = c \sin A$$

Substituting this value of h into the geometric area formula results in

$$\text{area} = \frac{1}{2}b(c \sin A)$$

$$= \frac{1}{2}bc \sin A$$

In general, *the area of a triangle is equal to one-half the product of the lengths of any two sides and the sine of their included angle;* that is,

$$\text{area} = \frac{1}{2}ab \sin C = \frac{1}{2}ac \sin B = \frac{1}{2}bc \sin A$$

EXAMPLE: Find the area of triangle ABC to one decimal place if $a = 13$, $b = 9$, and $C = 40°$.

SOLUTION: Since C is the angle between sides a and b, the area formula is

$$\text{area} = \frac{1}{2}ab \sin C$$

so

$$\text{area} = \frac{1}{2}(13)(9) \sin 40°$$

$$= 58.5(0.64279)$$

$$= 37.6$$

Another formula for the area of a triangle can be derived by the use of the Law of Sines and the previous formula. From the Law of Sines,

$$\frac{b}{\sin B} = \frac{c}{\sin C}$$

we find

$$b = \frac{c \sin B}{\sin C}$$

Substituting this value of b into the previous area formula

$$\text{area} = \frac{1}{2}bc \sin A$$

results in

$$\text{area} = \frac{1}{2}\left(\frac{c \sin B}{\sin C}\right)c \sin A$$

$$= \frac{c^2 \sin A \sin B}{2 \sin C}$$

Therefore, *the area of a triangle can be determined if one side and two angles are known* (since the third angle can be found directly); that is,

$$\text{area} = \frac{a^2 \sin B \sin C}{2 \sin A} = \frac{b^2 \sin A \sin C}{2 \sin B} = \frac{c^2 \sin A \sin B}{2 \sin C}$$

EXAMPLE: Find the area of triangle ABC to one decimal place if $A = 25°$, $C = 105°$, and $b = 12$.

SOLUTION: First, we find B to be

$$B = 180° - (A + C)$$

$$= 180° - (25° + 105°)$$

$$= 50°$$

The area formula for this situation would be

$$\text{area} = \frac{b^2 \sin A \sin C}{2 \sin B}$$

so,

$$\text{area} = \frac{(12)^2 \sin 25^\circ \sin 105^\circ}{2 \sin 50^\circ}$$

$$= \frac{144(0.42262)(0.96593)}{2(0.76604)}$$

$$= 38.4$$

PRACTICE PROBLEMS:

Find the area of triangle ABC to three decimal places given the following measurements:

1. $b = 20.02$, $c = 40.04$, and $A = 80^\circ\ 8'$

2. $a = 3.28$, $c = 9.18$, and $B = 42^\circ\ 21'$

3. $B = 50^\circ$, $C = 70^\circ$, and $c = 5.07$

4. $A = 103^\circ\ 48'$, $B = 34^\circ\ 6'$, and $a = 4.24$

ANSWERS:

1. 394.873

2. 10.142

3. 9.074

4. 3.479

SUMMARY

The following are the major topics covered in this chapter:

1. **Oblique triangles:** *Oblique triangles* are triangles containing no right angles. Oblique triangles are made up of either three acute angles or two acute angles and one obtuse angle.

   Acute angles have measures between 0° and 90°.

   Obtuse angles have measures between 90° and 180°.

2. **Law of Sines:** The lengths of the sides of any triangle are proportional to the sines of their opposite angles.

$$\frac{a}{\sin A} = \frac{b}{\sin B} \quad \frac{c}{\sin C}$$

3. **Standard cases for solving oblique triangles using the Law of Sines:**

   Case 1. One side and two angles

   Case 2. Two sides and an angle opposite one of them (This is referred to as the *ambiguous case* since two triangles, one triangle, or no triangle may result from the given data.)

4. **Law of Cosines:** In a triangle, the square of any side is equal to the sum of the squares of the other two sides minus twice the product of the same two sides multiplied by the cosine of the angle between them.

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$b^2 = a^2 + c^2 - 2ac \cos B$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$

5. **Standard cases for solving oblique triangles using the Law of Cosines:**

   Case 3. Two sides and the included angle

   Case 4. All three sides

6. **Area of a triangle:**

The area of a triangle is equal to one-half the product of the lengths of any two sides and the sine of their included angle.

$$\text{area} = \frac{1}{2}ab \sin C = \frac{1}{2}ac \sin B = \frac{1}{2}bc \sin A$$

The area of a triangle can be determined if one side and two angles are known.

$$\text{area} = \frac{a^2 \sin B \sin C}{2 \sin A} = \frac{b^2 \sin A \sin C}{2 \sin B} = \frac{c^2 \sin A \sin B}{2 \sin C}$$

ADDITIONAL PRACTICE PROBLEMS

Use the Law of Sines, Law of Cosines, or area formulas to solve the following problems:

1. To determine the distance from point A to point B across a canyon, Barbara lays off a distance from point C to point B as 440 yards. She then finds that $C = 30° 17'$ and $B = 104° 53'$. What is the distance, to the nearest yard, between points A and B?

2. Two buoys are 325 feet apart and a boat is 250 feet from one of them. The angle subtended by the two buoys at the boat is $65° 10'$. Find the distance, to the nearest foot, from the boat to the other buoy.

3. A triangular tract of land is to be enclosed by a fence. Side a equals 37.25 feet, side c equals 46.98 feet, and the included angle B is $100° 30'$. Find the amount of fencing, to the nearest hundredth of a foot, needed to enclose the triangular plot.

4. A 12-foot ladder is placed against an inclined support and reaches 10 feet up the side of the support. The foot of the ladder is 5 feet from the foot of the inclined support. What is the measure of the angle, to the nearest minute, the ladder makes with the support?

5. Find the area, to one decimal place, of a triangular field if two sides of the field are 127 yards and 159 yards and the included angle is $57° 18'$.

6. What is the area of a parallelogram, to one decimal place, if the length of one diagonal is 6 inches and the diagonal meets two adjacent sides of the parallelogram at angles with measures $33°$ and $44°$? HINT: Double the area of a triangle.

ANSWERS TO ADDITIONAL PRACTICE PROBLEMS

1. 315 yards
2. 338 feet
3. 149.29 feet
4. $24° 9'$
5. 8,496.3 square yards
6. 14.0 square inches

CHAPTER 6

TRIGONOMETRIC IDENTITIES AND EQUATIONS

LEARNING OBJECTIVES

Upon completion of this chapter, you should be able to do the following:

1. Apply the reciprocal, quotient, and Pythagorean identities along with identities for negative angles to problem solving.

2. Apply the sum and difference, double-angle, and half-angle formulas to problem solving.

3. Apply inverse trigonometric functions to problem solving.

4. Find solutions to trigonometric equations.

INTRODUCTION

This is the final chapter dealing directly with trigonometry and trigonometric relationships. This chapter includes the basic identities, formulas for identities involving more than one angle, and formulas for identities involving multiples of an angle.

Also included in this chapter are inverse trigonometric functions and methods for solving trigonometric equations.

FUNDAMENTAL IDENTITIES

An equality that is true for all values of an unknown is called an *identity*. Many of the identities that will be considered in this section were established in earlier chapters and will be used here to change the form of an expression.

Problems in identities are often given as equalities. The identity is established by either transforming the left side into the right side or transforming the right side into the left side. Never work across the equality sign.

We have no hard-and-fast rules to use in verifying identities. However, we do offer the following suggestions:

1. *Know the basic identities given in this section.*
2. *Attempt to transform the more complicated side into the other side.*
3. *When possible, express all trigonometric functions in the equation in terms of sine and cosine.*
4. *Perform any factoring or algebraic operations.*

RECIPROCAL IDENTITIES

The *reciprocal identities* were first introduced in chapter 3. They are as follows:

$$\cot \theta = \frac{1}{\tan \theta}$$

$$\sec \theta = \frac{1}{\cos \theta}$$

$$\csc \theta = \frac{1}{\sin \theta}$$

EXAMPLE: Use the reciprocal identities to find an equivalent expression involving only sines and cosines; then simplify for

$$\frac{\sec \theta}{\csc \theta + \sec \theta}$$

SOLUTION:

$$\frac{\sec \theta}{\csc \theta + \sec \theta} = \frac{\dfrac{1}{\cos \theta}}{\dfrac{1}{\sin \theta} + \dfrac{1}{\cos \theta}}$$

$$= \frac{\dfrac{1}{\cos \theta}}{\dfrac{\cos \theta + \sin \theta}{\sin \theta \cos \theta}}$$

$$= \left(\frac{1}{\cos \theta}\right)\left(\frac{\sin \theta \cos \theta}{\cos \theta + \sin \theta}\right)$$

$$= \frac{\sin \theta}{\cos \theta + \sin \theta}$$

QUOTIENT IDENTITIES

The *quotient identities* were also introduced in chapter 3. They are as follows:

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\cot \theta = \frac{\cos \theta}{\sin \theta}$$

EXAMPLE: Use the quotient identities to find an equivalent expression involving only sines and cosines; then simplify for

$$\frac{\tan \theta}{\cot \theta}$$

SOLUTION:

$$\frac{\tan \theta}{\cot \theta} = \frac{\dfrac{\sin \theta}{\cos \theta}}{\dfrac{\cos \theta}{\sin \theta}}$$

$$= \left(\frac{\sin \theta}{\cos \theta}\right)\left(\frac{\sin \theta}{\cos \theta}\right)$$

$$= \frac{\sin^2\theta}{\cos^2\theta}$$

PYTHAGOREAN IDENTITIES

Another group of fundamental identities, called the *Pythagorean identities*, involves the squares of the functions. These identities are so named because the Pythagorean theorem is used in their development.

Consider

$$x^2 + y^2 = r^2$$

and divide both sides by r^2 to get

$$\frac{x^2}{r^2} + \frac{y^2}{r^2} = 1$$

or

$$\left(\frac{x}{r}\right)^2 + \left(\frac{y}{r}\right)^2 = 1$$

Since $\cos \theta = x/r$ and $\sin \theta = y/r$, then

$$(\cos \theta)^2 + (\sin \theta)^2 = 1$$

or

$$\cos^2\theta + \sin^2\theta = 1$$

which is one of the Pythagorean identities.

In the same manner, dividing both sides of the equation

$$x^2 + y^2 = r^2$$

by x^2 (where $x \neq 0$) gives

$$1 + \frac{y^2}{x^2} = \frac{r^2}{x^2}$$

or

$$1 + \left(\frac{y}{x}\right)^2 = \left(\frac{r}{x}\right)^2$$

Since $\tan \theta = y/x$ and $\sec \theta = r/x$, then

$$1 + (\tan \theta)^2 = (\sec \theta)^2$$

or

$$1 + \tan^2\theta = \sec^2\theta$$

which is another one of the Pythagorean identities.

Dividing both sides of the equation

$$x^2 + y^2 = r^2$$

by y^2 (where $y \neq 0$) gives

$$\frac{x^2}{y^2} + 1 = \frac{r^2}{y^2}$$

or

$$1 + \left(\frac{x}{y}\right)^2 = \left(\frac{r}{y}\right)^2$$

Since $\cot \theta = x/y$ and $\csc \theta = r/y$, then

$$1 + (\cot \theta)^2 = (\csc \theta)^2$$

or

$$1 + \cot^2\theta = \csc^2\theta$$

which is also one of the Pythagorean identities.

EXAMPLE: Use the Pythagorean identities to find an equivalent expression involving only sines and cosines; then simplify for

$$(\csc^2\theta - 1)(\tan^2\theta + 1)$$

SOLUTION:

$$\begin{aligned}(\csc^2\theta - 1)(\tan^2\theta + 1) &= \cot^2\theta \sec^2\theta \\ &= \left(\frac{\cos^2\theta}{\sin^2\theta}\right)\left(\frac{1}{\cos^2\theta}\right) \\ &= \frac{1}{\sin^2\theta}\end{aligned}$$

IDENTITIES FOR NEGATIVE ANGLES

The following *identities for negative angles* were first introduced in chapter 4:

$$\sin(-\theta) = -\sin\theta$$

$$\cos(-\theta) = \cos\theta$$

$$\tan(-\theta) = -\tan\theta$$

EXAMPLE: Use the identities for negative angles to find an equivalent expression involving only sines and cosines with positive angles; then simplify for

$$\frac{\tan(-\theta)}{\cos(-\theta)}$$

SOLUTION:

$$\begin{aligned}\frac{\tan(-\theta)}{\cos(-\theta)} &= \frac{-\tan\theta}{\cos\theta} \\ &= \frac{-\dfrac{\sin\theta}{\cos\theta}}{\cos\theta} \\ &= -\frac{\sin\theta}{\cos^2\theta}\end{aligned}$$

VERIFYING TRIGONOMETRIC IDENTITIES

The process of verifying trigonometric identities is similar to simplifying trigonometric expressions except that we know in advance the desired result. Remember to use the suggestions we offered at the beginning of this chapter when verifying trigonometric identities.

EXAMPLE: Verify the identity

$$\frac{\sin\theta}{\csc\theta} + \frac{\cos\theta}{\sec\theta} = 1$$

SOLUTION:

$$\frac{\sin\theta}{\csc\theta} + \frac{\cos\theta}{\sec\theta} = \frac{\sin\theta}{\frac{1}{\sin\theta}} + \frac{\cos\theta}{\frac{1}{\cos\theta}}$$

$$= (\sin\theta)(\sin\theta) + (\cos\theta)(\cos\theta)$$

$$= \sin^2\theta + \cos^2\theta$$

$$= 1$$

EXAMPLE: Verify the identity

$$1 + \cot^2 2x = \frac{1}{\sin^2 2x}$$

SOLUTION:

$$1 + \cot^2 2x = \csc^2 2x$$

$$= \frac{1}{\sin^2 2x}$$

EXAMPLE: Verify the identity

$$2\sec\theta = \frac{\cos(-\theta)}{1} + \frac{\cos(-\theta)}{1}$$

SOLUTION:

$$\frac{\cos(-\theta)}{1-\sin(-\theta)}+\frac{\cos(-\theta)}{1+\sin(-\theta)}$$

$$=\frac{\cos\theta}{1+\sin\theta}+\frac{\cos\theta}{1-\sin\theta}$$

$$=\frac{(\cos\theta)(1-\sin\theta)+(\cos\theta)(1+\sin\theta)}{(1+\sin\theta)(1-\sin\theta)}$$

$$=\frac{\cos\theta-\cos\theta\sin\theta+\cos\theta+\cos\theta\sin\theta}{1+\sin\theta-\sin\theta-\sin^2\theta}$$

$$=\frac{2\cos\theta}{1-\sin^2\theta}$$

$$=\frac{2\cos\theta}{\cos^2\theta}$$

$$=\frac{2}{\cos\theta}$$

$$=2\sec\theta$$

EXAMPLE: Verify the identity

$$\frac{1+2\sin\theta+\sin^2\theta}{\cos^2\theta}=\frac{\sec\theta+\tan\theta}{\sec\theta-\tan\theta}$$

SOLUTION:

$$\frac{\sec\theta+\tan\theta}{\sec\theta-\tan\theta}=\frac{\frac{1}{\cos\theta}+\frac{\sin\theta}{\cos\theta}}{\frac{1}{\cos\theta}-\frac{\sin\theta}{\cos\theta}}$$

$$=\frac{\frac{1+\sin\theta}{\cos\theta}}{\frac{1-\sin\theta}{\cos\theta}}$$

$$=\left(\frac{1+\sin\theta}{\cos\theta}\right)\left(\frac{\cos\theta}{1-\sin\theta}\right)$$

$$=\frac{1+\sin\theta}{1-\sin\theta}$$

$$=\left(\frac{1+\sin\theta}{1-\sin\theta}\right)\left(\frac{1+\sin\theta}{1+\sin\theta}\right)$$

$$=\frac{1+\sin\theta+\sin\theta+\sin^2\theta}{1-\sin\theta+\sin\theta-\sin^2\theta}$$

$$=\frac{1+2\sin\theta+\sin^2\theta}{1-\sin^2\theta}$$

$$=\frac{1+2\sin\theta+\sin^2\theta}{\cos^2\theta}$$

PRACTICE PROBLEMS:

Verify the following identities:

1. $\dfrac{1}{\tan^2 x + 1} = \cos^2 x$

2. $\csc \theta - \sin \theta = \cos \theta \cot \theta$

3. $\dfrac{\sin^2 \theta}{1 + \cos \theta} = 1 - \cos \theta$

4. $\sin^2 \theta = [\cos (-\theta)][\sec (-\theta) - \cos (-\theta)]$

5. $1 - \cos^2 x = (\tan^2 x)(1 - \sin^2 x)$

6. $\dfrac{1 - \cos^2 \theta}{\csc \theta} = \sin^3 \theta$

7. $\dfrac{1}{2 + \cot^2(-\theta)} = \dfrac{1}{2 \csc^2(-\theta) - \cot^2(-\theta)}$

NOTE: No ANSWERS are furnished since the result is known in advance for each of the preceding PRACTICE PROBLEMS.

FORMULAS FOR IDENTITIES

In this section we will discuss the trigonometric formulas for the sum and difference of angles, for double angles, and for half angles.

SUM AND DIFFERENCE FORMULAS

The fundamental identities discussed in the previous section involved functions of a single angle. In this section we will consider identities involving functions of more than one angle.

We will start by developing a formula for cos $(\alpha - \beta)$. Refer to figure 6-1. Angles α and β are constructed in standard position, so angle KOL is equal to α and angle KOM is equal to β. We will also construct angle KON equal to $\alpha - \beta$. Since triangles KON and MOL are similar triangles, then sides LM and KN have the same length.



Figure 6-1.—Developing formula for co

Now we need to determine the coordinates of points K, L, M, and N. Recall the properties of right triangles, quadrantal angles, and reduction formulas. For the unit circle, where $r = 1$, the coordinates of point K, which lie on the positive X axis, are (1,0). According to properties of right triangles

$$\cos \theta = \frac{x}{r}$$

and

$$\sin \theta = \frac{y}{r}$$

So the coordinates of point N are [cos $(\alpha - \beta)$,sin $(\alpha - \beta)$].

Recall from chapter 4 that

$$\cos (180° - \theta) = -\cos \theta$$

and

$$\sin (180° - \theta) = \sin \theta$$

where θ is a positive acute angle. If we apply these formulas to angles α and β and note that the coordinates of a point in the second quadrant are $(-x,y)$, then the coordinates of point L are (cos α,sin α) and the coordinates of point M are (cos β,sin β).

Using the coordinates of these points and the distance formula, we can determine the lengths of sides LM and KN. Hence,

$$\begin{aligned}(LM)^2 &= (\cos\alpha - \cos\beta)^2 + (\sin\alpha - \sin\beta)^2 \\ &= \cos^2\alpha - 2\cos\alpha\cos\beta + \cos^2\beta \\ &\quad + \sin^2\alpha - 2\sin\alpha\sin\beta + \sin^2\beta \\ &= 2 - 2\cos\alpha\cos\beta - 2\sin\alpha\sin\beta\end{aligned}$$

and

$$\begin{aligned}(KN)^2 &= [1 - \cos(\alpha - \beta)]^2 + [0 - \sin(\alpha - \beta)]^2 \\ &= 1 - 2\cos(\alpha - \beta) + \cos^2(\alpha - \beta) + \sin^2(\alpha - \beta) \\ &= 2 - 2\cos(\alpha - \beta)\end{aligned}$$

Since sides LM and KN have the same length, we can equate the distances and simplify as follows:

$$\begin{aligned}2 - 2\cos\alpha\cos\beta - 2\sin\alpha\sin\beta &= 2 - 2\cos(\alpha - \beta) \\ -2(\cos\alpha\cos\beta + \sin\alpha\sin\beta) &= -2\cos(\alpha - \beta) \\ \cos\alpha\cos\beta + \sin\alpha\sin\beta &= \cos(\alpha - \beta)\end{aligned}$$

Therefore, *the cosine of the difference of two angles is equal to the cosine of the first angle times the cosine of the second angle plus the sine of the first angle times the sine of the second angle*; that is,

$$\cos(\alpha - \beta) = \cos\alpha\cos\beta + \sin\alpha\sin\beta$$

EXAMPLE: Simplify $\cos(90° - \beta)$.

SOLUTION: If

$$\cos(\alpha - \beta) = \cos\alpha\cos\beta + \sin\alpha\sin\beta$$

then

$$\begin{aligned}\cos(90° - \beta) &= \cos 90°\cos\beta + \sin 90°\sin\beta \\ &= (0)(\cos\beta) + (1)(\sin\beta) \\ &= \sin\beta\end{aligned}$$

which is the same result shown in chapter 4.

EXAMPLE: Determine $\cos 15°$ using the cosine of the difference of two angles.

SOLUTION:

$$\begin{aligned}\cos 15° &= \cos (45° - 30°) \\ &= \cos 45° \cos 30° + \sin 45° \sin 30° \\ &= \left(\frac{\sqrt{2}}{2}\right)\left(\frac{\sqrt{3}}{2}\right) + \left(\frac{\sqrt{2}}{2}\right)\left(\frac{1}{2}\right) \\ &= \frac{\sqrt{6} + \sqrt{2}}{4}\end{aligned}$$

To develop a formula for $\cos (\alpha + \beta)$, we will substitute $(-\beta)$ for β into the formula for $\cos (\alpha - \beta)$ as follows:

$$\begin{aligned}\cos (\alpha + \beta) &= \cos [\alpha - (-\beta)] \\ &= \cos \alpha \cos (-\beta) + \sin \alpha \sin (-\beta) \\ &= \cos \alpha \cos \beta - \sin \alpha \sin \beta\end{aligned}$$

Therefore, *the cosine of the sum of two angles is equal to the cosine of the first angle times the cosine of the second angle minus the sine of the first angle times the sine of the second angle;* that is,

$$\cos (\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

EXAMPLE: Determine $\cos 105°$ using the cosine of the sum of two angles.

SOLUTION:

$$\begin{aligned}\cos 105° &= \cos (45° + 60°) \\ &= \cos 45° \cos 60° - \sin 45° \sin 60° \\ &= \left(\frac{\sqrt{2}}{2}\right)\left(\frac{1}{2}\right) - \left(\frac{\sqrt{2}}{2}\right)\left(\frac{\sqrt{3}}{2}\right) \\ &= \frac{\sqrt{2} - \sqrt{6}}{4}\end{aligned}$$

We will now use the identities

$$\cos \theta = \sin (90° - \theta)$$

and

$$\sin \theta = \cos (90° - \theta)$$

and the formula

$$\cos (\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

to develop a formula for $\sin (\alpha + \beta)$ as follows:

$$\begin{aligned}
\sin (\alpha + \beta) &= \cos [90° - (\alpha + \beta)] \\
&= \cos [(90° - \alpha) - \beta] \\
&= \cos (90° - \alpha) \cos \beta + \sin (90° - \alpha) \sin \beta \\
&= \sin \alpha \cos \beta + \cos \alpha \sin \beta
\end{aligned}$$

Therefore, *the sine of the sum of two angles is equal to the sine of the first angle times the cosine of the second angle plus the cosine of the first angle times the sine of the second angle;* that is,

$$\sin (\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

EXAMPLE: Verify that

$$\sin (\alpha + 45°) = \frac{\sqrt{2}}{2}(\sin \alpha + \cos \alpha)$$

SOLUTION:

$$\begin{aligned}
\sin (\alpha + 45°) &= \sin \alpha \cos 45° + \cos \alpha \sin 45° \\
&= (\sin \alpha)\left(\frac{\sqrt{2}}{2}\right) + (\cos \alpha)\left(\frac{\sqrt{2}}{2}\right) \\
&= \frac{\sqrt{2}}{2}(\sin \alpha + \cos \alpha)
\end{aligned}$$

Substituting $(-\beta)$ for β into the formula for $\sin (\alpha + \beta)$ produces

$$\begin{aligned}
\sin (\alpha - \beta) &= \sin [\alpha + (-\beta)] \\
&= \sin \alpha \cos (-\beta) + \cos \alpha \sin (-\beta) \\
&= \sin \alpha \cos \beta - \cos \alpha \sin \beta.
\end{aligned}$$

Therefore, *the sine of the difference of two angles is equal to the sine of the first angle times the cosine of the second angle minus the cosine of the first angle times the sine of the second angle*; that is,

$$\sin(\alpha - \beta) = \sin\alpha\cos\beta - \cos\alpha\sin\beta$$

EXAMPLE: Use the formula for the sine of the difference of two angles to determine the value of

$$\sin 40^\circ \cos 10^\circ - \cos 40^\circ \sin 10^\circ$$

SOLUTION:

$$\begin{aligned}\sin 40^\circ \cos 10^\circ - \cos 40^\circ \sin 10^\circ &= \sin(40^\circ - 10^\circ)\\ &= \sin 30^\circ\\ &= 1/2\end{aligned}$$

Now, using the identity

$$\tan\theta = \frac{\sin\theta}{\cos\theta}$$

and the formulas for $\sin(\alpha + \beta)$ and $\cos(\alpha + \beta)$, we can develop a formula for $\tan(\alpha + \beta)$ as follows:

$$\begin{aligned}\tan(\alpha + \beta) &= \frac{\sin(\alpha + \beta)}{\cos(\alpha + \beta)}\\ &= \frac{\sin\alpha\cos\beta + \cos\alpha\sin\beta}{\cos\alpha\cos\beta - \sin\alpha\sin\beta}\end{aligned}$$

Dividing both the numerator and denominator by $\cos\alpha\cos\beta$ gives

$$\begin{aligned}\tan(\alpha + \beta) &= \frac{\dfrac{\sin\alpha\cos\beta}{\cos\alpha\cos\beta} + \dfrac{\cos\alpha\sin\beta}{\cos\alpha\cos\beta}}{\dfrac{\cos\alpha\cos\beta}{\cos\alpha\cos\beta} - \dfrac{\sin\alpha\sin\beta}{\cos\alpha\cos\beta}}\\ &= \frac{\tan\alpha + \tan\beta}{1 - \tan\alpha\tan\beta}\end{aligned}$$

Therefore, *the tangent of the sum of two angles is equal to the quantity of the tangent of the first angle plus the tangent of the*

second angle divided by the quantity of 1 minus the tangent of the first angle times the tangent of the second angle; that is,

$$\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$$

EXAMPLE: If $\sin \alpha = -4/5$ and $\cos \beta = 12/13$, where α is in quadrant III and β is in quadrant IV, find $\tan(\alpha + \beta)$.

SOLUTION: Refer to figure 6-2. If $\sin \alpha = -4/5$ and α is in quadrant III, then $\tan \alpha = 4/3$. Likewise, if $\cos \beta = 12/13$ and β is in quadrant IV, then $\tan \beta = -5/12$. Therefore,

$$\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$$

$$= \frac{(4/3) + (-5/12)}{1 - (4/3)(-5/12)}$$

$$= \frac{11/12}{1 + 20/36}$$

$$= \left(\frac{11}{12}\right)\left(\frac{36}{56}\right)$$

$$= \frac{33}{56}$$



Figure 6-2.—Triangles in quadra

As before, to develop a formula for $\tan(\alpha - \beta)$, we will substitute $(-\beta)$ for β into the formula for $\tan(\alpha + \beta)$ as follows:

$$\tan(\alpha - \beta) = \tan[\alpha + (-\beta)]$$

$$= \frac{\tan \alpha + \tan(-\beta)}{1 - \tan \alpha \tan(-\beta)}$$

$$= \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \tan \beta}$$

Therefore, *the tangent of the difference of two angles is equal to the quantity of the tangent of the first angle minus the tangent of the second angle divided by the quantity of 1 plus the tangent of the first angle times the tangent of the second angle*; that is,

$$\tan(\alpha - \beta) = \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \tan \beta}$$

EXAMPLE: If $\csc \alpha = 29/21$, $\sin \beta = -8/17$, $\cos \alpha$ is negative, and $\sec \beta$ is positive, find $\cot (\alpha - \beta)$.

SOLUTION: Note that

$$\cot (\alpha - \beta) = \frac{1}{\tan (\alpha - \beta)}$$

So we determine the value of $\cot (\alpha - \beta)$ using the formula for $\tan (\alpha - \beta)$. Since $\csc \alpha$ is positive and $\cos \alpha$ is negative in quadrant II, then $\tan \alpha = -21/20$. Likewise, since $\sin \beta$ is negative and $\sec \beta$ is positive in quadrant IV, then $\tan \beta = -8/15$. Hence,

$$\tan (\alpha - \beta) = \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \tan \beta}$$

$$= \frac{(-21/20) - (-8/15)}{1 + (-21/20)(-8/15)}$$

$$= \frac{-31/60}{1 + 14/25}$$

$$= \left(\frac{-31}{60}\right)\left(\frac{25}{39}\right)$$

$$= -\frac{155}{468}$$

Therefore,

$$\cot (\alpha - \beta) = -\frac{468}{155}$$

PRACTICE PROBLEMS:

Use sum and difference formulas to find the values of the following:

1. $\sin \frac{13\pi}{12}$

2. $\cot 165°$

Verify the following using sum and difference formulas:

3. $\dfrac{\cos(\alpha - \beta)}{\cos\alpha \sin\beta} = \cot\beta + \tan\alpha$

4. $\tan\left(\alpha - \dfrac{\pi}{4}\right) = \dfrac{\tan\alpha - 1}{\tan\alpha + 1}$

5. If $\sin\alpha = -1/4$ and $\cos\beta = -4/5$, where α and β are both in quadrant III, find $\cos(\alpha + \beta)$.

ANSWERS:

1. $\dfrac{-\sqrt{6} + \sqrt{2}}{4}$

2. $\dfrac{1 + \sqrt{3}}{1 - \sqrt{3}}$ or $-2 - \sqrt{3}$

3. Result is known

4. Result is known

5. $\dfrac{4\sqrt{15} - 3}{20}$

DOUBLE-ANGLE FORMULAS

Formulas for the functions of twice an angle may be derived from the functions of the sum of two angles. Setting $\beta = \alpha$ in the formulas for $\sin(\alpha + \beta)$, $\cos(\alpha + \beta)$, and $\tan(\alpha + \beta)$ gives the following results:

$$\sin(\alpha + \alpha) = \sin\alpha\cos\alpha + \cos\alpha\sin\alpha$$

$$\cos(\alpha + \alpha) = \cos\alpha\cos\alpha - \sin\alpha\sin\alpha$$

$$\tan(\alpha + \alpha) = \frac{\tan\alpha + \tan\alpha}{1 - \tan\alpha\tan\alpha}$$

Hence,

$$\sin 2\alpha = 2 \sin \alpha \cos \alpha$$

$$\begin{aligned}\cos 2\alpha &= \cos^2\alpha - \sin^2\alpha \\ &= 1 - 2 \sin^2\alpha \\ &= 2 \cos^2\alpha - 1\end{aligned}$$

$$\tan 2\alpha = \frac{2 \tan \alpha}{1 - \tan^2\alpha}$$

The previous formulas are known as the *double-angle formulas.*

EXAMPLE: Find the values for $\sin 2\theta$, $\cos 2\theta$, and $\tan 2\theta$, if $\tan \theta = -12/5$ and θ is in the second quadrant.

SOLUTION: Since θ is in the second quadrant, then $\sin \theta = 12/13$ and $\cos \theta = -5/13$; so,

$$\begin{aligned}\sin 2\theta &= 2 \sin \theta \cos \theta \\ &= 2\left(\frac{12}{13}\right)\left(\frac{-5}{13}\right) \\ &= -\frac{120}{169}\end{aligned}$$

$$\begin{aligned}\cos 2\theta &= \cos^2\theta - \sin^2\theta \\ &= \left(\frac{-5}{13}\right)^2 - \left(\frac{12}{13}\right)^2 \\ &= \frac{25}{169} - \frac{144}{169} \\ &= -\frac{119}{169}\end{aligned}$$

$$\begin{aligned}\tan 2\theta &= \frac{2 \tan \theta}{1 - \tan^2\theta} \\ &= \frac{2(-12/5)}{1 - (-12/5)^2} \\ &= \frac{-24/5}{1 - 144/25} \\ &= \frac{-24/5}{-119/25} \\ &= \left(\frac{24}{5}\right)\left(\frac{25}{119}\right) \\ &= \frac{120}{119}\end{aligned}$$

EXAMPLE: Verify that

$$\csc 2x = \frac{1}{2} \csc x \sec x$$

SOLUTION:

$$\frac{1}{2} \csc x \sec x = \frac{1}{2}\left(\frac{1}{\sin x}\right)\left(\frac{1}{\cos x}\right)$$

$$= \frac{1}{2 \sin x \cos x}$$

$$= \frac{1}{\sin 2x}$$

$$= \csc 2x$$

HALF-ANGLE FORMULAS

From the double-angle formulas we can derive the *half-angle formulas*. Since

$$\cos 2\alpha = 1 - 2 \sin^2 \alpha$$

then solving for $\sin \alpha$ results in

$$\sin \alpha = \pm \sqrt{\frac{1 - \cos 2\alpha}{2}}$$

Now, if $2\alpha = \theta$, so that $\alpha = \theta/2$, then

$$\sin \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{2}}$$

which is the half-angle formula for $\sin \theta/2$.

The half-angle formula for $\cos \theta/2$ can be obtained by solving

$$\cos 2\alpha = 2 \cos^2 \alpha - 1$$

for $\cos \alpha$ or

$$\cos \alpha = \pm \sqrt{\frac{1 + \cos 2\alpha}{2}}$$

As before, if $2\alpha = \theta$, so that $\alpha = \theta/2$, then

$$\cos \frac{\theta}{2} = \pm \sqrt{\frac{1 + \cos \theta}{2}}$$

The half-angle formula for tan $\theta/2$ is derived from the half-angle formulas for sine and cosine as follows:

$$\tan\frac{\theta}{2} = \frac{\sin\frac{\theta}{2}}{\cos\frac{\theta}{2}}$$

$$= \frac{\pm\sqrt{\frac{1-\cos\theta}{2}}}{\pm\sqrt{\frac{1+\cos\theta}{2}}}$$

$$= \pm\sqrt{\frac{1-\cos\theta}{1+\cos\theta}}$$

NOTE: For the half-angle formulas, the positive or negative sign is selected according to the quadrant in which $\theta/2$ lies.

EXAMPLE: Use the half-angle formulas to find the cosine, sine, and tangent of 112.5°.

SOLUTION: Since 112.5° lies in quadrant II, the cosine and tangent will be negative and the sine will be positive; so,

$$\cos 112.5° = -\sqrt{\frac{1+\cos 225°}{2}}$$

$$= -\sqrt{\frac{1+(-\sqrt{2}/2)}{2}}$$

$$= -\sqrt{\frac{2-\sqrt{2}}{4}}$$

$$= -\frac{\sqrt{2-\sqrt{2}}}{2}$$

$$\sin 112.5° = \sqrt{\frac{1-\cos 225°}{2}}$$

$$= \sqrt{\frac{1-(-\sqrt{2}/2)}{2}}$$

$$= \sqrt{\frac{2+\sqrt{2}}{4}}$$

$$= \frac{\sqrt{2+\sqrt{2}}}{2}$$

$$\tan 112.5^\circ = -\sqrt{\frac{1 - \cos 225^\circ}{1 + \cos 225^\circ}}$$

$$= -\sqrt{\frac{1 - (-\sqrt{2}/2)}{1 + (-\sqrt{2}/2)}}$$

$$= -\sqrt{\frac{1 + \sqrt{2}/2}{1 - \sqrt{2}/2}}$$

$$= -\sqrt{\frac{2 + \sqrt{2}}{2 - \sqrt{2}}}$$

$$= -\sqrt{\left(\frac{2 + \sqrt{2}}{2 - \sqrt{2}}\right)\left(\frac{2 + \sqrt{2}}{2 + \sqrt{2}}\right)}$$

$$= -\sqrt{\frac{6 + 4\sqrt{2}}{2}}$$

$$= -\sqrt{3 + 2\sqrt{2}}$$

EXAMPLE: Verify that

$$\sin^2\left(\frac{\theta}{2}\right) = \frac{\sec \theta - 1}{2 \sec \theta}$$

SOLUTION:

$$\frac{\sec \theta - 1}{2 \sec \theta} = \frac{\dfrac{1}{\cos \theta} - 1}{2\left(\dfrac{1}{\cos \theta}\right)}$$

$$= \frac{\dfrac{1 - \cos \theta}{\cos \theta}}{\dfrac{2}{\cos \theta}}$$

$$= \frac{1 - \cos \theta}{2}$$

$$= \sin^2\left(\frac{\theta}{2}\right)$$

PRACTICE PROBLEMS:

1. Find the values for $\sin 2\theta$, $\cos 2\theta$, and $\tan 2\theta$, if $\theta = \pi$.

2. Find the values for $\sin \theta/2$, $\cos \theta/2$, and $\tan \theta/2$ if $\sec \theta = 17/8$, $\tan \theta$ is positive, and $0 \leq \theta \leq 360°$.

Verify the following using double-angle and half-angle formulas:

3. $(1 + \tan x)(\tan 2x) = \dfrac{2 \tan x}{1 - \tan x}$

4. $\dfrac{2}{1 + \cos \theta} - \tan^2\left(\dfrac{\theta}{2}\right) = 1$

ANSWERS:

1. $\sin 2\theta = 0$

   $\cos 2\theta = 1$

   $\tan 2\theta = 0$

2. $\sin \theta/2 = 3\sqrt{34}/34$

   $\cos \theta/2 = 5\sqrt{34}/34$

   $\tan \theta/2 = 3/5$

3. Result is known

4. Result is known

INVERSE TRIGONOMETRIC FUNCTIONS

In this section we will discuss the notations that apply to the inverse trigonometric functions along with the principal values of the inverse functions.

NOTATION

Let us consider the inverse of the sine function, $y = \sin x$. The inverse of the sine function may be denoted as

$$x = \sin y$$

or

$$y = \sin^{-1}x$$

which can be read "the inverse sine of x." Note that $\sin^{-1}x$ does not mean $1/\sin x$.

The inverse of the sine function may also be denoted by

$$y = \arcsin x$$

which can be read "the arc sine of x." The notation arcsin x arises because it is the length of an arc on the unit circle for which the sine is x.

Similar notation occurs for the inverses of the other trigonometric functions; that is, $\cos^{-1}x$ or arccos x, $\tan^{-1}x$ or arctan x, etc.

PRINCIPAL VALUES

For any angle, one, and only one, value of a trigonometric function corresponds to the angle; but for any value of a trigonometric function, numerous angles satisfy the value. Hence, the inverses of the trigonometric functions are not themselves functions. However, if we restrict the ranges of these relationships, we can obtain functions. The values of the trigonometric functions in the restricted ranges are called *principal values*. To indicate this restriction, we will capitalize the first letter in the name of the inverse trigonometric function; that is,

$$y = \text{Sin}^{-1}x$$

or

$$y = \text{Arcsin } x$$

Table 6-1.—Inverse Trigonometric Functions

FUNCTION	DOMAIN	RANGE
$y = \text{Sin}^{-1}x$	$-1 \leq x \leq 1$	$-\pi/2 \leq y \leq \pi/2$
$y = \text{Cos}^{-1}x$	$-1 \leq x \leq 1$	$0 \leq y \leq \pi$
$y = \text{Tan}^{-1}x$	Any real number	$-\pi/2 < y < \pi/2$
$y = \text{Cot}^{-1}x$	Any real number	$0 < y < \pi$
$y = \text{Sec}^{-1}x$	$x \leq -1$ or $x \geq 1$	$0 \leq y \leq \pi$, $y \neq \pi/2$
$y = \text{Csc}^{-1}x$	$x \leq -1$ or $x \geq 1$	$-\pi/2 \leq y \leq \pi/2$, $y \neq 0$

and so on for all the trigonometric functions. Table 6-1 shows the six inverse trigonometric functions, their domains, and their ranges.

EXAMPLE: Find all values of arctan 1.

SOLUTION: The tangent of many angles is 1, such as $\pi/4$, $5\pi/4$, $9\pi/4$, and $13\pi/4$. Thus the values of arctan 1 are

$$\frac{\pi}{4} + n\pi$$

where n is any integer.

EXAMPLE: Find Arctan 1.

SOLUTION: In the restricted range, as shown in table 6-1, the only number whose tangent is 1 is $\pi/4$. Hence,

$$\text{Arctan } 1 = \pi/4$$

EXAMPLE: Find Arcsec 2.236 in degrees.

SOLUTION: As previously determined,

$$\sec \theta = \frac{1}{\cos \theta}$$

If

$$x = \sec \theta$$

then

$$x = \frac{1}{\cos \theta}$$

$$\cos \theta = \frac{1}{x}$$

If we solve for θ in the above equations, then

$$\theta = \text{arcsec } x$$

and

$$\theta = \arccos \frac{1}{x}$$

so,

$$\text{arcsec } x = \arccos \frac{1}{x}$$

Hence, for the given problem

$$\text{Arcsec } 2.236 = \text{Arccos} \left(\frac{1}{2.236} \right)$$

$$= \text{Arccos } 0.44723$$

According to appendix II, the angle whose cosine is 0.44723 is $63^\circ 26'$ to the nearest minute, which is in the range $0 \leq y \leq 180^\circ$. Therefore,

$$\text{Arcsec } 2.236 = 63^\circ 26'$$

EXAMPLE: Find $\text{Cos}^{-1}(-0.50000)$ in degrees.

SOLUTION: According to appendix II, the angle whose cosine is 0.50000 is 60°. However, we want the angle whose cosine is a negative number so that the angle is in the range of $0 \leq y \leq 180^\circ$. Since the cosine of a number is negative in the second quadrant where the reference angle of 60° corresponds to 120°, then

$$\text{Cos}^{-1}(-0.50000) = 120^\circ$$

EXAMPLE: Find $\text{Cot}^{-1}(-\sqrt{3})$ in degrees.

SOLUTION: The expression $\text{Cot}^{-1}(-\sqrt{3})$ can be interpreted as "the angle between 0 and π, whose cotangent is $-\sqrt{3}$." Recall that

$$\cot \theta = \frac{\text{adjacent}}{\text{opposite}}$$

For the given problem,

$$\theta = \text{Cot}^{-1}(-\sqrt{3})$$

or

$$\text{Cot } \theta = -\sqrt{3}$$

From our previous discussion of special angles, you should recognize the reference angle of θ to be 30°. Since the cotangent of an angle is negative in the second quadrant for the range from 0° to 180°, then θ is 150°; that is,

$$\text{Cot}^{-1}(-\sqrt{3}) = 150°$$

EXAMPLE: Evaluate $\cos\left(\text{Arcsin } \frac{5}{13}\right)$.

SOLUTION: Let

$$u = \text{Arcsin } \frac{5}{13}$$

so

$$\text{Sin } u = \frac{5}{13}$$

Since Arcsin is defined only in quadrants I and IV and since 5/13 is positive, then u is in quadrant I. Figure 6-3 shows a triangle in quadrant I whose sine of angle u is 5/13. Using the Pythagorean theorem, we find that the side adjacent to angle u is 12. Therefore,



Figure 6-3.—Triangl

$$\cos u = \frac{12}{13}$$

or

$$\cos\left(\text{Arcsin}\ \frac{5}{13}\right) = \frac{12}{13}$$

EXAMPLE: Evaluate $\sin\left(\text{Arccos}\ \frac{12}{13} - \text{Arcsin}\ \frac{4}{5}\right)$.

SOLUTION: Let

$$u = \text{Arccos}\ \frac{12}{13}$$

or

$$\text{Cos}\ u = \frac{12}{13}$$

and let

$$v = \text{Arcsin}\ \frac{4}{5}$$

or

$$\text{Sin}\ v = \frac{4}{5}$$

Angles u and v would both be in quadrant I according to previous conditions. Figure 6-4, view A, shows a triangle in quadrant I where cos u = 12/13. Figure 6-4, view B, shows a triangle in quadrant I where sin v = 4/5.



Figure 6-4.—Triangles in quadrant I.

Our given equation is in the form of the difference formula

$$\sin(\alpha - \beta) = \sin\alpha\cos\beta - \cos\alpha\sin\beta$$

where

$$\sin\left(\text{Arccos}\,\frac{12}{13} - \text{Arcsin}\,\frac{4}{5}\right)$$

$$= \sin\left(\text{Arccos}\,\frac{12}{13}\right)\cos\left(\text{Arcsin}\,\frac{4}{5}\right)$$

$$- \cos\left(\text{Arccos}\,\frac{12}{13}\right)\sin\left(\text{Arcsin}\,\frac{4}{5}\right)$$

Referring to figure 6-4, view A, we find that

$$\sin\left(\text{Arccos}\,\frac{12}{13}\right) = \frac{5}{13}$$

and

$$\cos\left(\text{Arccos}\,\frac{12}{13}\right) = \frac{12}{13}$$

Referring to figure 6-4, view B, we find that

$$\sin\left(\text{Arcsin}\,\frac{4}{5}\right) = \frac{4}{5}$$

and

$$\cos\left(\text{Arcsin}\,\frac{4}{5}\right) = \frac{3}{5}$$

Hence,

$$\sin\left(\text{Arccos}\,\frac{12}{13} - \text{Arcsin}\,\frac{4}{5}\right) = \left(\frac{5}{13}\right)\left(\frac{3}{5}\right) - \left(\frac{12}{13}\right)\left(\frac{4}{5}\right)$$

$$= \frac{15 - 48}{65}$$

$$= -\frac{33}{65}$$

PRACTICE PROBLEMS:

1. Find $\text{Cos}^{-1}(1/2)$.

2. Find Arcsin 0.88295 in degrees.

3. Find $\text{Csc}^{-1}(-1.57208)$ in degrees.

4. Find Arccot $(-\sqrt{3}/3)$.

5. Evaluate tan [Arcsin $(-1/2)$].

6. Evaluate cot $[\text{Cos}^{-1}(-0.19994)]$.

7. Evaluate cos (Arctan 5/12 − Arccot 4/3).

8. Evaluate sin $[\text{Sec}^{-1}(-25/24) - \text{Csc}^{-1}(-17/8)]$.

ANSWERS:

1. $\pi/3$

2. $62°$

3. $-39° 30'$

4. $2\pi/3$

5. $-1/\sqrt{3}$ or $-\sqrt{3}/3$

6. -0.20406

7. $63/65$

8. $-87/425$

TRIGONOMETRIC EQUATIONS

A *trigonometric equation* is an equality that is true for some values but may not be true for all values of the variable. The principles and processes used to solve algebraic equations may be

used to solve trigonometric equations. The identities and formulas previously studied may also be used in solving trigonometric equations.

The following suggestions may be helpful to you in solving trigonometric equations:

1. *If only one trigonometric function is present, solve the equation for that function.*
2. *If more than one function is present, rearrange the equation so that one side equals 0. Then try to factor and set each factor equal to zero to solve. You may find it helpful to use identities and formulas to change the form of the equation or to square both sides of the equation.*
3. *If the equation is quadratic in form, but not factorable, use the quadratic formula.*
4. *All possible solutions should be tested in the given equation.*

EXAMPLE: Solve $\tan \theta - 1 = 0$ for $0° \leq \theta < 360°$.

SOLUTION: We can rewrite

$$\tan \theta - 1 = 0$$

to read

$$\tan \theta = 1$$

or

$$\theta = \arctan 1$$

and solve the equation. Therefore, the solutions in the given interval are

$$\theta = 45° \text{ and } 225°$$

EXAMPLE: Solve $\sin 2\theta = 2 \cos \theta$ for $0° \leq \theta < 360°$.

SOLUTION: We will rearrange the equation so that one side equals 0; hence,

$$\sin 2\theta - 2 \cos \theta = 0$$

Since,

$$\sin 2\theta = 2 \sin \theta \cos \theta$$

we will substitute this formula to change the form of our equation; that is,

$$2 \sin \theta \cos \theta - 2 \cos \theta = 0$$

We will now factor $2 \cos \theta$ from each term, so

$$2 \cos \theta (\sin \theta - 1) = 0$$

and set each factor equal to zero where

$$2 \cos \theta = 0$$

and

$$\sin \theta - 1 = 0$$

Solving each term gives

$$2 \cos \theta = 0$$

or

$$\theta = \arccos 0$$

where

$$\theta = 90° \text{ and } 270°$$

and

$$\sin \theta - 1 = 0$$

$$\sin \theta = 1$$

or

$$\theta = \arcsin 1$$

where

$$\theta = 90°$$

Substituting the value of $90°$ into the given equation gives

$$\sin 2(90°) = 2 \cos 90°$$

$$\sin 180° = (2)(0)$$

$$0 = 0$$

Substituting the value of $270°$ into the given equation gives

$$\sin 2(270°) = 2 \cos 270°$$

$$\sin 540° = 2(0)$$

$$0 = 0$$

Therefore, $\theta = 90°$ and $270°$ are the solutions to the equation.

EXAMPLE: Solve $\tan x - \sec x + 1 = 0$ for $0 \leq x < 2\pi$.

SOLUTION: Rewrite the given equation as

$$\tan x + 1 = \sec x$$

Square both sides of the equation to get

$$(\tan x + 1)^2 = (\sec x)^2$$

$$\tan^2 x + 2 \tan x + 1 = \sec^2 x$$

$$(\tan^2 x + 1) + 2 \tan x = \sec^2 x$$

Note that $\tan^2 x + 1 = \sec^2 x$, so

$$2 \tan x = 0$$

$$\tan x = 0$$

or

$$x = \arctan 0$$

Hence, the possible solutions are 0 and π. Substituting 0 into the original equation gives

$$\tan 0 + 1 = \sec 0$$

$$0 + 1 = 1$$

$$1 = 1$$

Substituting π into the original equation gives

$$\tan \pi + 1 = \sec \pi$$

$$0 + 1 = -1$$

but

$$1 \neq -1$$

Therefore, the only solution to the given equation is 0.

EXAMPLE: Solve $\cot^2\theta - 3 \cot \theta - 2 = 0$ for $0° \leq \theta < 360°$.

SOLUTION: Since this equation cannot be factored, we will use the quadratic formula, introduced in *Mathematics*, Volume 1,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

where the values for x are possible solutions to the equation

$$ax^2 + bx + c = 0$$

For our equation

$$x = \cot \theta$$

$$a = 1$$

$$b = -3$$

$$c = -2$$

Hence,

$$\cot \theta = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(1)(-2)}}{2(1)}$$

$$= \frac{3 \pm \sqrt{17}}{2}$$

So,

$$\cot \theta = 3.56155$$

or

$$\theta = \text{arccot } 3.56155$$

where

$$\theta = 15° 41' \text{ and } 195° 41'$$

to the nearest minute in quadrants I and III, respectively. And

$$\cot \theta = -0.56155$$

or

$$\theta = \text{arccot } (-0.56155)$$

where

$$\theta = 119° 19' \text{ and } 299° 19'$$

to the nearest minute in quadrants II and IV, respectively. Substituting all four of the values of

$$\theta = 15° 41', 119° 19', 195° 41', 299° 19'$$

into the original equation shows that they are solutions.

NOTE: When substituting a possible solution into the original equation, we may not be able to equate the sides exactly because of rounding error.

PRACTICE PROBLEMS:

1. Solve $\sin \theta = -\sqrt{3}/2$ for $0° \leq \theta < 360°$.
2. Solve $\tan x \cos^2 x = \sin^2 x$ for $0 \leq x < 2\pi$.
3. Solve $\cot \theta - \csc \theta - \sqrt{3} = 0$ for $0° \leq \theta < 360°$.
4. Solve $7 \sin^2 \theta - 3 \sin \theta - 4 = 0$ for $0° \leq \theta < 360°$.

ANSWERS:

1. $\theta = 240°$ and $300°$
2. $x = 0, \pi/4, \pi, 5\pi/4$
3. $\theta = 240°$
4. $\theta = 90°, 214° 51', \text{ and } 325° 9'$

SUMMARY

The following are the major topics covered in this chapter:

1. **Suggestions in solving identities:**

   1. Know the basic identities.
   2. Attempt to transform the more complicated side into the other side.
   3. When possible, express all trigonometric functions in the equation in terms of sine and cosine.
   4. Perform any factoring or algebraic operations.

2. **Reciprocal identities:**

$$\cot \theta = \frac{1}{\tan \theta}$$

$$\sec \theta = \frac{1}{\cos \theta}$$

$$\csc \theta = \frac{1}{\sin \theta}$$

3. **Quotient identities:**

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\cot \theta = \frac{\cos \theta}{\sin \theta}$$

4. **Pythagorean identities:**

$$\cos^2\theta + \sin^2\theta = 1$$

$$1 + \tan^2\theta = \sec^2\theta$$

$$1 + \cot^2\theta = \csc^2\theta$$

5. **Identities for negative angles:**

$$\sin (-\theta) = -\sin \theta$$

$$\cos (-\theta) = \cos \theta$$

$$\tan (-\theta) = -\tan \theta$$

6. **Sum and difference formulas:**

The cosine of the difference of two angles is equal to the cosine of the first angle times the cosine of the second angle plus the sine of the first angle times the sine of the second angle; that is,

$$\cos(\alpha - \beta) = \cos\alpha \cos\beta + \sin\alpha \sin\beta$$

The cosine of the sum of two angles is equal to the cosine of the first angle times the cosine of the second angle minus the sine of the first angle times the sine of the second angle; that is,

$$\cos(\alpha + \beta) = \cos\alpha \cos\beta - \sin\alpha \sin\beta$$

The sine of the sum of two angles is equal to the sine of the first angle times the cosine of the second angle plus the cosine of the first angle times the sine of the second angle; that is,

$$\sin(\alpha + \beta) = \sin\alpha \cos\beta + \cos\alpha \sin\beta$$

The sine of the difference of two angles is equal to the sine of the first angle times the cosine of the second angle minus the cosine of the first angle times the sine of the second angle; that is,

$$\sin(\alpha - \beta) = \sin\alpha \cos\beta - \cos\alpha \sin\beta$$

The tangent of the sum of two angles is equal to the quantity of the tangent of the first angle plus the tangent of the second angle divided by the quantity of 1 minus the tangent of the first angle times the tangent of the second angle; that is,

$$\tan(\alpha + \beta) = \frac{\tan\alpha + \tan\beta}{1 - \tan\alpha \tan\beta}$$

The tangent of the difference of two angles is equal to the quantity of the tangent of the first angle minus the tangent of the second angle divided by the quantity of 1 plus the tangent of the first angle times the tangent of the second angle; that is

$$\tan(\alpha - \beta) = \frac{\tan\alpha - \tan\beta}{1 + \tan\alpha \tan\beta}$$

7. **Double-angle formulas:**

$$\sin 2\alpha = 2 \sin \alpha \cos \alpha$$

$$\cos 2\alpha = \cos^2\alpha - \sin^2\alpha$$

$$= 1 - 2 \sin^2\alpha$$

$$= 2 \cos^2\alpha - 1$$

$$\tan 2\alpha = \frac{2 \tan \alpha}{1 - \tan^2\alpha}$$

8. **Half-angle formulas:**

$$\sin \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{2}}$$

$$\cos \frac{\theta}{2} = \pm \sqrt{\frac{1 + \cos \theta}{2}}$$

$$\tan \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{1 + \cos \theta}}$$

9. **Inverse trigonometric functions:**

$$x = \sin y \text{ or } y = \sin^{-1}x \text{ or } y = \arcsin x$$

$$x = \cos y \text{ or } y = \cos^{-1}x \text{ or } y = \arccos x$$

$$x = \tan y \text{ or } y = \tan^{-1}x \text{ or } y = \arctan x$$

$$x = \cot y \text{ or } y = \cot^{-1}x \text{ or } y = \operatorname{arccot} x$$

$$x = \sec y \text{ or } y = \sec^{-1}x \text{ or } y = \operatorname{arcsec} x$$

$$x = \csc y \text{ or } y = \csc^{-1}x \text{ or } y = \operatorname{arccsc} x$$

10. **Principal values:** The values of the trigonometric functions in the restricted ranges are called *principal values*. This restriction is indicated by the capitalization of the first letter of the name of the inverse trigonometric function.

11. **Trigonometric equations:** A *trigonometric equation* is an equality that is true for some values but may not be true for all values of the variable.

12. **Suggestions in solving trigonometric equations:**

1. If only one trigonometric function is present, solve the equation for that function.
2. If more than one function is present, rearrange the equation so that one side equals 0. Then try to factor and set each factor equal to zero to solve. You may find it helpful to use identities and formulas to change the form of the equation or to square both sides of the equation.
3. If the equation is quadratic in form, but not factorable, use the quadratic formula.
4. All possible solutions should be tested in the given equation.

ADDITIONAL PRACTICE PROBLEMS

1. Verify that $\frac{1}{1 - \cos x} + \frac{1}{1 + \cos x} = 2 \csc^2 x$.

2. Verify that $\frac{\sin (x + y)}{\cos (x - y)} = \frac{\cot x + \cot y}{1 + \cot x \cot y}$ using sum and difference formulas.

3. If $\tan \alpha = 8/15$ with α in quadrant I and $\cos \beta = 7/25$ with β in quadrant IV, find $\sec (\alpha - \beta)$.

4. Verify that $\tan x = \frac{1 - \cos 2x}{\sin 2x}$ using double-angle formulas.

5. Verify that $8 \sin^2\left(\frac{x}{2}\right) \cos^2\left(\frac{x}{2}\right) = 1 - \cos 2x$ using half-angle formulas.

6. Find Arcsec (-2).

7. Find $\text{Tan}^{-1}(-0.12278)$ in degrees.

8. Evaluate $\sin [2 \text{ Tan}^{-1}(12/5)]$. HINT: $\sin 2\theta = 2 \sin \theta \cos \theta$.

9. Evaluate $\tan \left[\text{Arccos} \frac{\sqrt{3}}{2} - \text{Arcsin} \left(\frac{-3}{5} \right) \right]$.

10. Solve $\sin x + \cos x = \sqrt{2}$ if $0 \leq x < 2\pi$.

ANSWERS TO ADDITIONAL PRACTICE PROBLEMS

1. Result is known
2. Result is known
3. $-425/87$
4. Result is known
5. Result is known
6. $2\pi/3$
7. $-7°$
8. $120/169$
9. $\dfrac{4 + 3\sqrt{3}}{4\sqrt{3} - 3}$ or $\dfrac{25\sqrt{3} + 48}{39}$
10. $\pi/4$

CHAPTER 7

VECTORS AND FORCES

LEARNING OBJECTIVES

Upon completion of this chapter, you should be able to do the following:

1. Add, subtract, and determine the components of vectors.
2. Solve problems involving forces.
3. Solve problems involving translational and rotational equilibrium.

INTRODUCTION

The last chapter in this course deals with vectors and forces. Any study of vectors and forces requires a knowledge of trigonometry.

VECTORS

A *scalar quantity* is one that has magnitude only; that is, 10 watts, 4 miles, 17 acres, and 28.2 pounds per square inch. A *vector quantity* is one that has both magnitude and direction; that is, 6 miles due north, 250 knots at 30°, and 400 miles per hour to the west. Scalar quantities are represented by italicized letters. Vector quantities are represented by placing arrows over the italicized letters, for instance, $\vec{A}$; and the magnitude of the vector quantity is represented by the italicized letter of the vector quantity. Vectors are geometrically represented by arrows. The arrowhead represents the terminal end of a vector and indicates the vector's direction. The other end of the vector is called the initial end. The magnitude of the vector is the vector's length.

Two vectors are said to be equal if they are of the same length, are parallel, and point in the same direction. In figure 7-1, view A, $\vec{A} = \vec{B}$.

If two vectors have the same length, are parallel, but point in opposite directions, they are said to be negatives of each other. In figure 7-1, view B, $\vec{C} = -\vec{D}$ and $\vec{D} = -\vec{C}$.



Figure 7-1.—Equal and nega

Hence, a vector can be moved from one position to another without being changed if its direction and magnitude are kept unchanged.

VECTOR ADDITION

The general rule for adding vectors is illustrated in figure 7-2. To add $\vec{B}$ to $\vec{A}$, shift $\vec{B}$ until its initial end coincides with the terminal end of $\vec{A}$. In its new position, $\vec{B}$ will be parallel, the same length, and in the same direction it was in the old position of $\vec{B}$. To find the vector sum of $\vec{A} + \vec{B}$, draw a vector, $\vec{R}$, with its initial end at the initial end of $\vec{A}$ and its terminal end at the terminal end of $\vec{B}$. Hence, $\vec{R}$ is called the *resultant vector* of $\vec{A}$ and $\vec{B}$, which is written

$$\vec{R} = \vec{A} + \vec{B}$$

If we reverse the process to add $\vec{A}$ to $\vec{B}$, we would move $\vec{A}$ until its initial end coincides with the terminal end of $\vec{B}$ so that



Figure 7-2.—Addition of $\vec{B}$ to $\vec{A}$.

is also the resultant vector of $\vec{B} + \vec{A}$. (See fig. 7-3.) us

$$\vec{R} = \vec{A} + \vec{B} = \vec{B} + \vec{A}$$

To find the resultant vector of any number of vectors, place the tial end of each vector to the terminal end of the previous vector. sure the new vector position is parallel, the same length, and in e same direction as the old position. Draw a vector $\vec{R}$, from the tial end of the first vector to the terminal end of the last vector so at $\vec{R}$ is the resultant vector. Figure 7-4 shows how four vectors e added together. We recognize again that the order in which e vectors are added does not affect the result. (See fig. 7-5.) Thus,

$$\vec{R} = \vec{A} + \vec{B} + \vec{C} + \vec{D} = \vec{D} + \vec{C} + \vec{B} + \vec{A}$$



Figure 7-3.—Ad $\vec{A}$ to $\vec{B}$.

ECTOR SUBTRACTION

We can also subtract one vector from another vector. Refer figure 7-6. To subtract $\vec{B}$ from $\vec{A}$, we need to determine



Figure 7-4.—Addition of $\vec{A} + \vec{B} + \vec{C} + \vec{D}$.



Figure 7-5.—Addition of $\vec{D} + \vec{C} + \vec{B} + \vec{A}$.



Figure 7-6.—Vector subtraction.

the negative of $\vec{B}$, denoted by $-\vec{B}$. The negative of a vector is a vector that is parallel, the same length, and points in the opposite direction. Hence, we add $-\vec{B}$ to $\vec{A}$ as previously discussed. This process may be summarized as

$$\vec{A} - \vec{B} = \vec{A} + (-\vec{B})$$

COMPONENTS OF VECTORS

The projections of a vector onto the X and Y axes of the rectangular coordinate system are called the *components* of a vector. We say that a vector is resolved into its x and y components, called the *horizontal* and *vertical components* of a vector, respectively. In figure 7-7, view A,

$$\vec{V}_x = \text{horizontal component of } \vec{V}$$

and

$$\vec{V}_y = \text{vertical component of } \vec{V}$$

Figure 7-7, view B, shows the magnitudes of $\vec{V}$, $\vec{V}_x$, and $\vec{V}_y$. Using properties of right triangles, we see that

$$\cos \theta = \frac{V_x}{V}$$

or

$$V_x = V \cos \theta$$

and

$$\sin \theta = \frac{V_y}{V}$$



Figure 7-7.—Components of a vector.

or

$$V_y = V \sin \theta$$

where V_x, V_y, and V are the magnitudes of $\vec{V}_x$, $\vec{V}_y$, and $\vec{V}$, respectively. Notice also that (by use of the Pythagorean theorem) *the magnitude of V can be found to be*

$$V^2 = V_x^2 + V_y^2$$

or

$$V = \sqrt{V_x^2 + V_y^2}$$

The direction of $\vec{V}$ is the angle, θ, the vector makes with the horizontal. This direction can be determined by

$$\tan \theta = \frac{V_y}{V_x}$$

or

$$\theta = \arctan \frac{V_y}{V_x}$$

The direction of $\vec{V}_x$ is 0° and the direction of $\vec{V}_y$ is 90°.

EXAMPLE: Find the magnitude of the horizontal and vertical components of a vector having a magnitude of 50 pounds acting at an angle of 30° to the horizontal.

SOLUTION:

$$\begin{aligned} V_x &= V \cos \theta \\ &= 50 \cos 30° \\ &= 50\left(\frac{\sqrt{3}}{2}\right) \\ &= 25\sqrt{3} \\ &= 43.3 \text{ pounds (rounded)} \end{aligned}$$

and

$$\begin{aligned} V_y &= V \sin \theta \\ &= 50 \sin 30° \\ &= 50\left(\frac{1}{2}\right) \\ &= 25 \text{ pounds} \end{aligned}$$

EXAMPLE: Find the magnitude and direction of a vector whose horizontal and vertical components have a magnitude of 90 newtons and 60 newtons, respectively.

SOLUTION: The magnitude of the resultant vector is

$$\begin{aligned} V &= \sqrt{V_x^2 + V_y^2} \\ &= \sqrt{(90)^2 + (60)^2} \\ &= \sqrt{8{,}100 + 3{,}600} \\ &= \sqrt{11{,}700} \\ &= 108.2 \text{ newtons (rounded)} \end{aligned}$$

The direction of the resultant vector is

$$\begin{aligned} \theta &= \arctan \frac{60}{90} \\ &= \arctan 0.66667 \\ &= 33^\circ\, 41' \text{ (to the nearest minute)} \end{aligned}$$

VECTOR ADDITION BY COMPONENTS

We can add vectors that lie in the same plane by working in terms of their components. This procedure is as follows:

1. *Resolve the given vectors into their x and y components.*
2. *Add the magnitudes of the x components to give R_x (the magnitude of the x component of $\vec{R}$), and add the magnitudes of the y components to give R_y (the magnitude the y component of $\vec{R}$); that is,*

$$R_x = A_x + B_x + C_x + \ldots$$

and

$$R_y = A_y + B_y + C_y + \ldots$$

3. *Find the magnitude and direction of $\vec{R}$ from R_x and R_y. The magnitude can be determined by the use of the Pythagorean theorem; that is,*

$$R = \sqrt{R_x^2 + R_y^2}$$

The direction of $\vec{R}$ can be found from the values of the components by trigonometry; that is,

$$\theta = \arctan \frac{R_y}{R_x}$$

EXAMPLE: A girl walks 110 feet south, 100 feet east, 120 feet northeast, and then 90 feet northwest. What is the magnitude and direction from her starting point?

SOLUTION: For convenience we will call north the positive y direction, south the negative y direction, east the positive x direction, and west the negative x direction on a rectangular coordinate system. Refer to figure 7-8, view A, for the path the girl walks. Figure 7-8, view B, shows the magnitude and direction of each vector from the origin according to the rectangular coordinate system. Hence, the magnitudes of the components of $\vec{A}$ are

$$\begin{aligned} A_x &= A \cos \theta \\ &= 110 \cos 270° \\ &= 110(0) \\ &= 0 \text{ feet} \end{aligned}$$

and

$$\begin{aligned} A_y &= A \sin \theta \\ &= 110 \sin 270° \\ &= 110(-1) \\ &= -110 \text{ feet} \end{aligned}$$



Figure 7-8.—Vector addition by components.

The magnitudes of the components of $\vec{B}$ are

$$\begin{aligned} B_x &= B \cos \theta \\ &= 100 \cos 0^\circ \\ &= 100(1) \\ &= 100 \text{ feet} \end{aligned}$$

and

$$\begin{aligned} B_y &= B \sin \theta \\ &= 100 \sin 0^\circ \\ &= 100(0) \\ &= 0 \text{ feet} \end{aligned}$$

The magnitudes of the components of $\vec{C}$ are

$$\begin{aligned} C_x &= C \cos \theta \\ &= 120 \cos 45^\circ \\ &= 120\left(\frac{\sqrt{2}}{2}\right) \\ &= 60\sqrt{2} \text{ feet} \end{aligned}$$

and

$$\begin{aligned} C_y &= C \sin \theta \\ &= 120 \sin 45^\circ \\ &= 120\left(\frac{\sqrt{2}}{2}\right) \\ &= 60\sqrt{2} \text{ feet} \end{aligned}$$

And the magnitudes of the components of $\vec{D}$ are

$$\begin{aligned} D_x &= D \cos \theta \\ &= 90 \cos 135^\circ \\ &= -90 \cos 45^\circ \\ &= -90\left(\frac{\sqrt{2}}{2}\right) \\ &= -45\sqrt{2} \text{ feet} \end{aligned}$$

and

$$\begin{aligned} D_y &= D \sin \theta \\ &= 90 \sin 135° \\ &= 90 \sin 45° \\ &= 90\left(\frac{\sqrt{2}}{2}\right) \\ &= 45\sqrt{2} \text{ feet} \end{aligned}$$

Now we will add the magnitudes of the x components to get R_x and add the magnitudes of the y components to get R_y:

$$\begin{aligned} R_x &= A_x + B_x + C_x + D_x \\ &= 0 + 100 + 60\sqrt{2} - 45\sqrt{2} \\ &= 121.21 \text{ feet (rounded)} \end{aligned}$$

and

$$\begin{aligned} R_y &= A_y + B_y + C_y + D_y \\ &= -110 + 0 + 60\sqrt{2} + 45\sqrt{2} \\ &= 38.49 \text{ feet (rounded)} \end{aligned}$$

Therefore, the magnitude of the resultant vector from the girl's starting point is

$$\begin{aligned} R &= \sqrt{R_x^2 + R_y^2} \\ &= \sqrt{(121.21)^2 + (38.49)^2} \\ &= \sqrt{16{,}173.34} \\ &= 127.17 \text{ feet (rounded)} \end{aligned}$$

and the direction from her starting point is

$$\begin{aligned} \theta &= \arctan \frac{R_y}{R_x} \\ &= \arctan \frac{38.49}{121.21} \\ &= \arctan 0.31755 \\ &= 17° 37' \text{ north of east} \end{aligned}$$

PRACTICE PROBLEMS:

Give magnitude accuracy to one decimal place and angle accuracy to the nearest minute for the following problems:

1. Find V_x and V_y of $\vec{V}$ having a magnitude of 325 pounds making an angle of $78^\circ 20'$ with the horizontal component vector.

2. Two vectors, having magnitudes of 150 newtons and 220 newtons, act at right angles to each other. Find the magnitude of their resultant vector and the angle it makes with the larger vector.

3. An airplane is heading due east with an airspeed (speed relative to the air) of 350 mph. A wind is blowing from due south at 58 mph.

   a. Find the airplane's angle of drift (the angle between its heading and its actual course).

   b. Find the ground speed (actual speed along its course).

4. Given three vectors with magnitudes and directions of 40 feet, 60°; 60 feet, 150°; and 80 feet, 225°, find the magnitude and the direction of the resultant vector.

ANSWERS:

1. $V_x = 65.7$ pounds

   $V_y = 318.3$ pounds

2. $V = 266.3$ newtons

   $\theta = 34^\circ 17'$

3. a. $9^\circ 25'$ north of east

   b. 354.8 mph

4. $V = 88.9$ feet

   $\theta = 174^\circ 50'$

FORCES

A *force* produces or prevents motion or has the tendency to do so. The effect of a force upon a body depends upon the magnitude and direction of the force. Therefore, *a force can be represented by a vector quantity*. The *resolution of a force*, then, is the separation of a single force into two or more component forces acting in given directions on the same point. Moreover, when two or more forces act on the same body, the *resultant force* is the single force whose effect upon the body is equal in magnitude and direction to the combined effects of all the forces acting on the body.

EXAMPLE: Two dogs on leashes held by a person are trying to move in directions perpendicular to each other, one pulling with a force of 64 pounds, the other with a 52-pound force. Find the magnitude of the resultant force and the angle it makes with the larger force vector.

SOLUTION: Refer to figure 7-9. If we let $\vec{F}$ be the force vector, then $\vec{F}_x$ and $\vec{F}_y$ are the two components at right angles to each other. If point A is the person holding the two dogs, $\overrightarrow{AB}$ the larger force vector, and $\overrightarrow{AD}$ the smaller force vector, then



Figure 7-9

$$F = \sqrt{F_x^2 + F_y^2}$$

$$= \sqrt{(AB)^2 + (\mathrm{AD})^2}$$

$$= \sqrt{(64)^2 + (52)^2}$$

$$= \sqrt{6{,}800}$$

$$= 82.5 \text{ pounds (rounded)}$$

and

$$\theta = \arctan \frac{F_y}{F_x}$$

$$= \arctan \frac{52}{64}$$

$$= \arctan 0.81250$$

$$= 39^\circ\, 6'$$

EXAMPLE: An automobile weighing 3,200 pounds is parked on a driveway that makes a 17° angle with the horizontal. Find the components of the car's weight parallel and perpendicular to the driveway.

SOLUTION: The weight of an object is the gravitational force the earth exerts on it, which is a force that acts vertically downward. (See fig. 7-10.) Since $\vec{F} = \vec{W}$ is vertical and $\vec{F}_x$ is perpendicular to the driveway, then the angle between $\vec{F}$ and $\vec{F}_x$ is also $\theta = 17°$. Hence,

$$\begin{aligned} F_x &= F \cos \theta \\ &= W \cos \theta \\ &= 3{,}200 \cos 17° \\ &= 3{,}200(0.95630) \\ &= 3{,}060.2 \text{ pounds (rounded)} \end{aligned}$$



Figure 7-10.—G

which is the car's weight perpendicular to the driveway, and

$$\begin{aligned} F_y &= F \sin \theta \\ &= W \sin \theta \\ &= 3{,}200 \sin 17° \\ &= 3{,}200(0.29237) \\ &= 935.6 \text{ pounds (rounded)} \end{aligned}$$

which is the car's weight parallel to the driveway.

PRACTICE PROBLEMS:

1. A force of 235 pounds makes an angle of 60° 40′ with the vertical. Resolve the force into its horizontal and vertical components. Give component accuracy to one decimal place.

2. Two forces, 15 pounds at 335° and 25 pounds at 14°, act on the same point. Determine the magnitude (to one decimal place) and direction (to the nearest minute) of the resultant force.

3. A force of 53.5 pounds is acting on an object at a 25° angle to the horizontal. Find the horizontal and vertical components of the force. Give component accuracy to two decimal places.

ANSWERS:

1. $\vec{F}_x = 204.9$ pounds

   $\vec{F}_y = 115.1$ pounds

2. $F = 37.9$ pounds

   $\theta = 359° 34'$

3. $\vec{F}_x = 48.49$ pounds

   $\vec{F}_y = 22.61$ pounds

EQUILIBRIUM

If a body undergoes no change in its motion, it is said to be in a state of *equilibrium*. Two conditions are required for a body at rest to be in equilibrium. The body must have neither translatory (straight line) motion nor rotary (spinning) motion.

When two or more forces act together at a point, the *equilibriant force* is that single force applied at the same point which produces equilibrium. *The equilibriant force has a magnitude equal to that of the resultant of the separate forces, but it acts in the opposite direction.*

EXAMPLE: A force of 17 newtons at 123° and a force of 33 newtons at 333° act on the same point. Determine the magnitudes and directions of both the resultant and the equilibriant.

SOLUTION: For the force of 17 newtons at 123°, the magnitudes of the horizontal and vertical components are

$$\begin{aligned} A_x &= 17 \cos 123° \\ &= -17 \cos 57° \\ &= -9.3 \text{ newtons (rounded)} \end{aligned}$$

and

$$\begin{aligned} A_y &= 17 \sin 123° \\ &= 17 \sin 57° \\ &= 14.3 \text{ newtons (rounded)} \end{aligned}$$

For the force of 33 newtons at 333°, the magnitudes of the horizontal and vertical components are

$$\begin{aligned} B_x &= 33 \cos 333° \\ &= 33 \cos 27° \\ &= 29.4 \text{ newtons (rounded)} \end{aligned}$$

and

$$\begin{aligned} B_y &= 33 \sin 333° \\ &= -33 \sin 27° \\ &= -15.0 \text{ newtons (rounded)} \end{aligned}$$

Hence, the magnitudes of the horizontal and vertical components of the resultant vector are

$$\begin{aligned} F_x &= A_x + B_x \\ &= -9.3 + 29.4 \\ &= 20.1 \text{ newtons} \end{aligned}$$

and

$$\begin{aligned} F_y &= A_y + B_y \\ &= 14.3 + -15.0 \\ &= -0.7 \text{ newtons} \end{aligned}$$

Since the resultant and the equilibriant have equal magnitudes, then

$$F = \sqrt{F_x^2 + F_y^2}$$

$$= \sqrt{(20.1)^2 + (-0.7)^2}$$

$$= 20.1 \text{ newtons (rounded)}$$

The direction of the resultant is

$$\theta = \arctan \frac{F_y}{F_x}$$

$$= \arctan \frac{-0.7}{20.1}$$

$$= -2°$$

$$= 358°$$

(since our resultant force is in the fourth quadrant) and the direction of the equilibriant is

$$\theta' = \theta \pm 180°$$

$$= 358° - 180°$$

$$= 178°$$

TRANSLATIONAL EQUILIBRIUM

The first condition for equilibrium, no translatory motion, is met when no unbalanced forces act on a body. Therefore, *the sum of the forces acting on a body in any direction must be equal to the sum of the forces acting on a body in the opposite direction.*

Since the sum of all forces acting on a body must equal zero, then the sum of all the magnitudes of the horizontal components

ıust equal zero and the sum of all the magnitudes of the vertical omponents must equal zero; that is,

$$F_x = 0$$

nd

$$F_y = 0$$

EXAMPLE: Find the tension (a force acting against the esistance of a body) of a weightless rope supporting a 50-pound lock.

SOLUTION: Refer to figure 7-11. Since there are no horizontal omponents in this problem, we only need to find the magnitude f the vertical component of the tension, $\vec{T}_y$, and the mag-itude of the vertical component of the weight of the block, $\vec{W}_y$, such that

$$F_y = T_y + W_y = 0$$

r

$$T_y = -W_y$$

Since

$$W_y = W \sin \theta$$

$$= 50 \sin 270°$$

$$= -50 \text{ pounds}$$

ıen

$$T_y = -W_y$$

$$= -(-50)$$

$$= 50 \text{ pounds}$$

Therefore, the tension in the rope is equal to the weight being ıpported.

EXAMPLE: A weight of 10 newtons is supported by two ords. One cord makes an angle of 30° with the horizontal while ıe other makes an angle of 60° with the horizontal. Find the ension in each cord.



Figure 7-11 supported b



Figure 7-12.—Weight supported by two cords.

SOLUTION: Refer to figure 7-12. We begin by determining the magnitudes of the horizontal and vertical components of each vector. For tension $\vec{A}$, the reference angle (in the second quadrant) of sin θ is positive and the reference angle of cos θ is negative; so

$$A_x = -A \cos \theta$$

$$= -A \cos 30°$$

$$= -\frac{\sqrt{3}}{2}A \text{ newtons}$$

and

$$A_y = A \sin \theta$$

$$= A \sin 30°$$

$$= \frac{1}{2}A \text{ newtons}$$

For tension $\vec{B}$, the reference angles of $\sin\theta$ and $\cos\theta$ are both positive in quadrant I; so

$$\begin{aligned} B_x &= B\cos\theta \\ &= B\cos 60^\circ \\ &= \frac{1}{2}B \text{ newtons} \end{aligned}$$

and

$$\begin{aligned} B_y &= B\sin\theta \\ &= B\sin 60^\circ \\ &= \frac{\sqrt{3}}{2}B \text{ newtons} \end{aligned}$$

For the weight of 10 newtons,

$$\begin{aligned} W_x &= W\cos\theta \\ &= 10\cos 270^\circ \\ &= 0 \text{ newtons} \end{aligned}$$

and

$$\begin{aligned} W_y &= W\sin\theta \\ &= 10\sin 270^\circ \\ &= -10 \text{ newtons} \end{aligned}$$

Now, since the sum of the magnitudes of the horizontal components must equal zero and the sum of the magnitudes of the vertical components must equal zero, then

$$A_x + B_x + W_x = 0$$

or

$$-\frac{\sqrt{3}}{2}A + \frac{1}{2}B = 0$$

$$-\frac{\sqrt{3}}{2}A = -\frac{1}{2}B$$

$$\sqrt{3}A = B$$

and

$$A_y + B_y + W_y = 0$$

or

$$\frac{1}{2}A + \frac{\sqrt{3}}{2}B + -10 = 0$$

$$\frac{1}{2}A + \frac{\sqrt{3}}{2}B = 10$$

Substituting $\sqrt{3}A$ for B in the last equation, we obtain

$$\frac{1}{2}A + \frac{\sqrt{3}}{2}(\sqrt{3}A) = 10$$

$$\frac{1}{2}A + \frac{3}{2}A = 10$$

$$2A = 10$$

$$A = 5 \text{ newtons}$$

and

$$B = \sqrt{3}A$$

$$= \sqrt{3}(5)$$

$$= 8.7 \text{ newtons (rounded)}$$

ROTATIONAL EQUILIBRIUM

The second condition for equilibrium, no rotary motion, is met when the sum of the torques acting upon a body about a point equals zero; that is,

$$\tau_R = \tau_1 + \tau_2 + \tau_3 + \ldots = 0$$

Hence, *the sum of all the clockwise torques equals the sum of all the counterclockwise torques about an axis of rotation. Torque* is the product of the magnitude of a force, F, and the length of its torque or lever arm, L, where L is measured perpendicular to the line of action of the force. Hence,

$$\tau = -F_1L_1 + F_2L_2 + F_3L_3 + \ldots = 0$$

If a force tends to produce a counterclockwise rotation about an axis, the torque will be considered positive. If a force tends to produce a clockwise rotation about an axis, the torque will be considered negative.

EXAMPLE: A person exerts a 15-pound force at the end of an 8-inch wrench. (See fig. 7-13.) If this force makes an angle of 45° with the handle, what is the torque produced on the nut?



Figure 7-13.—Torque.

SOLUTION: First we need to find the length of the torque arm. Since L is measured perpendicular to the line of action, then to solve for L, we will use

$$\sin \theta = \frac{\text{opposite}}{\text{hypotenuse}}$$

or

$$\sin 45° = \frac{L}{8}$$

$$L = 8 \sin 45°$$

$$= 8\left(\frac{\sqrt{2}}{2}\right)$$

$$= 4\sqrt{2} \text{ inches}$$

Since the force tends to produce a counterclockwise rotation about the axis, then the torque produced on the nut is positive and

$$\tau = FL$$

$$= 15(4\sqrt{2})$$

$$= 60\sqrt{2}$$

$$= 84.9 \text{ pound} \cdot \text{inches (rounded)}$$

EXAMPLE: A rod 12 meters long has weights of 5 newtons and 15 newtons at its ends. (Assume that the weight of the rod is negligible.) At what point should the rod be picked up if it is to have no tendency to rotate (where is the balance point of the rod)?

SOLUTION: Refer to figure 7-14. First, compute the torques about the unknown balance point. If x is the distance of the 15-newton weight from this point, then the 5-newton weight is $(12 - x)$ meters from the point on the other side. Note that x is the length of the torque arm of the 15-newton weight and $(12 - x)$ is the length of the torque arm of the 5-newton weight. Since the 15-newton weight tends to produce a counterclockwise rotation about the balance point of the rod, then the torque will be considered positive. Likewise, since the 5-newton weight tends to produce a clockwise rotation about the balance point, then the torque will be considered negative. Hence,



Figure 7-14.—Balanced rod.

$$\tau_1 = F_1L_1$$
$$= 15(x)$$
$$= 15x$$

and

$$\tau_2 = F_2L_2$$
$$= -5(12 - x)$$
$$= -60 + 5x$$

Since no rotary motion occurs when

$$\tau_R = \tau_1 + \tau_2 = 0$$

then

$$F_1L_1 + F_2L_2 = 0$$
$$15x - 60 + 5x = 0$$
$$20x = 60$$
$$x = 3 \text{ meters}$$

Therefore, when the rod is picked up 3 meters from the 15-newton weight end, the two weights exert opposite torques of the same magnitude [$15x = 45 = 5(12 - x)$] about this point, where the rod is balanced.

PRACTICE PROBLEMS:

Give magnitude accuracy to one decimal place and angl accuracy to the nearest minute for the following:

1. Find the magnitudes and directions of the resultant an equilibriant forces of a force of 21 newtons due nort and a second force of 32 newtons southeast.

2. A 20-pound ball is suspended by a rope, A, attached to wall. Rope A is pulled away from the wall by horizontal rope, B, and is held so that rope A forms a angle of 30° with the vertical wall. Find the tension in ropes A and B.

3. A 150-newton force is applied to a pole 6 meters abov its base at an angle of 45° above the horizontal. Fin the torque about the base of the pole.

4. A uniform horizontal bar is 550 millimeters long and i of negligible weight. A 32-newton weight is hung fron the left end of the bar, and a 70-newton weight is hun from the right end. Where should a single upward suppor be positioned to balance the system?

ANSWERS:

1. $R = 22.7$ newtons

   $E = 22.7$ newtons

   $\theta_R = 355° 57'$

   $\theta_E = 175° 57'$

2. $A = 23.1$ pounds

   $B = 11.5$ pounds

3. 636.4 newton · meters

4. 172.5 millimeters from the right end

SUMMARY

The following are the major topics covered in this chapter:

1. **Definitions:**

   A *scalar quantity* is one that has magnitude only.

   A *vector quantity* is one that has both magnitude and direction.

2. **Vectors:**

   Two vectors are said to be equal if they are of the same length, are parallel, and point in the same direction.

   If two vectors have the same length, are parallel, but point in opposite directions, they are said to be negatives of each other.

3. **Resultant vectors:** To find the resultant vector of any number of vectors, place the initial end of each vector to the terminal end of the previous vector. Be sure the new vector position is parallel, the same length, and in the same direction as the old vector position. Draw a vector from the initial end of the first vector to the terminal end of the last vector. This newly formed vector is the *resultant vector*.

4. **Vector addition:**

$$\vec{R} = \vec{A} + \vec{B} = \vec{B} + \vec{A}$$

   where $\vec{A}$ and $\vec{B}$ are added and $\vec{R}$ is their resultant vector.

5. **Vector subtraction:**

$$\vec{A} - \vec{B} = \vec{A} + (-\vec{B})$$

   where $\vec{B}$ is subtracted from $\vec{A}$, which is the same as adding the negative of $\vec{B}$ to $\vec{A}$.

6. **Components of vectors:** The projections of a vector onto the X and Y axes of the rectangular coordinate system are called the *components* of a vector. The *horizontal component* of $\vec{V}$ is $\vec{V}_x$ and the *vertical component* of $\vec{V}$ is $\vec{V}_y$.

7. **Magnitudes of vectors:** The magnitudes of V_x, V_y, and V, respectively, are

$$V_x = V \cos \theta$$

$$V_y = V \sin \theta$$

$$V = \sqrt{V_x^2 + V_y^2}$$

8. **Directions of vectors:** The direction of $\vec{V}$ is the angle, θ, the vector makes with the horizontal.

$$\theta = \arctan \frac{V_y}{V_x}$$

The direction of $\vec{V}_x$ is 0° and the direction of $\vec{V}_y$ is 90°.

9. **Vector addition by components:**

1. Resolve the given vectors into their x and y components.

2. Add the magnitudes of the x components to give R_x (the magnitude of the x component of $\vec{R}$), and add the magnitudes of the y components to give R_y (the magnitude of the y component of $\vec{R}$); that is,

$$R_x = A_x + B_x + C_x + \ldots$$

and

$$R_y = A_y + B_y + C_y + \ldots$$

3. Find the magnitude and direction of $\vec{R}$ from R_x and R_y. The magnitude can be determined by the use of the Pythagorean theorem; that is,

$$R = \sqrt{R_x^2 + R_y^2}$$

The direction of $\vec{R}$ can be found from the values of the components by trigonometry; that is,

$$\theta = \arctan \frac{R_y}{R_x}$$

10. **Force:** A *force* produces or prevents motion or has the tendency to do so.

A force can be represented by a vector quantity.

The *resolution of a force* is the separation of a single force into two or more component forces acting in given directions on the same point.

When two or more forces act on the same body, the *resultant force* is the single force whose effect upon the body is equal in magnitude and direction to the combined effects of all the forces acting on the body.

11. **Equilibrium:** If a body undergoes no change in its motion, it is said to be in a state of *equilibrium*.

    For a body at rest to be in equilibrium, it must have neither translatory motion nor rotary motion.

12. **Equilibriant force:** When two or more forces act together at a point, the *equilibriant force* is that single force applied at the same point which produces equilibrium.

    The equilibriant force has a magnitude equal to that of the resultant of the separate forces, but it acts in the opposite direction.

13. **Translational equilibrium:** The sum of the forces acting on a body in any direction must be equal to the sum of the forces acting on a body in the opposite direction, such that

$$F_x = 0$$

    and

$$F_y = 0$$

14. **Rotational equilibrium:** The sum of all the clockwise torques equals the sum of all the counterclockwise torques about an axis of rotation, such that,

$$\tau_R = \tau_1 + \tau_2 + \tau_3 + \ldots = F_1L_1 + F_2L_2 + F_3L_3 + \ldots = 0$$

    Torque is the product of the magnitude of force, F, and the length of its torque or lever arm, L, where L is measured perpendicular to the line of action of the force.

    If a force tends to produce a counterclockwise rotation about an axis, the torque will be considered positive. If a force tends to produce a clockwise rotation about an axis, the torque will be considered negative.

ADDITIONAL PRACTICE PROBLEMS

Give magnitude accuracy to one decimal place and angle accuracy to the nearest minute for the following:

1. An airplane is heading due south at 220 mph and the wind is blowing from due east at 65 mph. Find the airplane's angle of drift and its ground speed.

2. An airplane flies 100 miles west from city A to city B, then 100 miles north from city B to city C, and finally 50 miles southeast to city D. How far is it from city A to city D?

3. Three forces act simultaneously on a point. One force is 15 newtons at 0°; the second is 20 newtons at 210°; and the third is 30 newtons at 60°. Determine the magnitude and direction of the resultant force.

4. A force of 11 pounds at 111° acts on a point. A second force of 22 pounds at 222° and a third force of 33 pounds at 333° also act on the same point. Determine the magnitude and direction of the equilibriant force.

5. A 100-pound tightrope walker stands at the center of a rope that is 200 feet in length. If the rope sags 20 feet at the center, find the tension in each side of the rope.

6. A 40-pound child and a 60-pound child sit at opposite ends of a 12-foot seesaw pivoted at its center. Where should a third child who weighs 50 pounds sit in order to balance the seesaw?

ANSWERS TO ADDITIONAL PRACTICE PROBLEMS

1. $\theta = 16° 28'$ west of south

   $S = 229.4$ mph

2. 91.4 miles

3. $F = 20.4$ newtons

   $\theta = 51° 34'$

4. $E = 21.5$ pounds

   $\theta = 115° 22'$

5. 255 pounds

6. 2.4 feet from the pivot on the same side as the 40-pound child

APPENDIX I

COMMON LOGARITHMS OF NUMBE

No.	0	1	2	3	4	5	6	7	
1.0	.0000	.0043	.0086	.0128	.0170	.0212	.0253	.0294	.
1.1	.0414	.0453	.0492	.0531	.0569	.0607	.0645	.0682	.
1.2	.0792	.0828	.0864	.0899	.0934	.0969	.1004	.1038	.
1.3	.1139	.1173	.1206	.1239	.1271	.1303	.1335	.1367	.
1.4	.1461	.1492	.1523	.1553	.1584	.1614	.1644	.1673	.
1.5	.1761	.1790	.1818	.1847	.1875	.1903	.1931	.1959	.
1.6	.2041	.2068	.2095	.2122	.2148	.2175	.2201	.2227	.
1.7	.2304	.2330	.2355	.2380	.2405	.2430	.2455	.2480	.
1.8	.2553	.2577	.2601	.2625	.2648	.2672	.2695	.2718	.
1.9	.2788	.2810	.2833	.2856	.2878	.2900	.2923	.2945	.
2.0	.3010	.3032	.3054	.3075	.3096	.3118	.3139	.3160	.
2.1	.3222	.3243	.3263	.3284	.3304	.3324	.3345	.3365	.
2.2	.3424	.3444	.3464	.3483	.3502	.3522	.3541	.3560	.
2.3	.3617	.3636	.3655	.3674	.3692	.3711	.3729	.3747	.
2.4	.3802	.3820	.3838	.3856	.3874	.3892	.3909	.3927	.
2.5	.3979	.3997	.4014	.4031	.4048	.4065	.4082	.4099	.
2.6	.4150	.4166	.4183	.4200	.4216	.4232	.4249	.4265	.
2.7	.4314	.4330	.4346	.4362	.4378	.4393	.4409	.4425	.
2.8	.4472	.4487	.4502	.4518	.4533	.4548	.4564	.4579	.
2.9	.4624	.4639	.4654	.4669	.4683	.4698	.4713	.4728	.
3.0	.4771	.4786	.4800	.4814	.4829	.4843	.4857	.4871	.
3.1	.4914	.4928	.4942	.4955	.4969	.4983	.4997	.5011	.
3.2	.5051	.5065	.5079	.5092	.5105	.5119	.5132	.5145	.
3.3	.5185	.5198	.5211	.5224	.5237	.5250	.5263	.5276	.
3.4	.5315	.5328	.5340	.5353	.5366	.5378	.5391	.5403	.
3.5	.5441	.5453	.5465	.5478	.5490	.5502	.5514	.5527	.
3.6	.5563	.5575	.5587	.5599	.5611	.5623	.5635	.5647	.
3.7	.5682	.5694	.5705	.5717	.5729	.5740	.5752	.5763	.
3.8	.5798	.5809	.5821	.5832	.5843	.5855	.5866	.5877	.
3.9	.5911	.5922	.5933	.5944	.5955	.5966	.5977	.5988	.
4.0	.6021	.6031	.6042	.6053	.6064	.6075	.6085	.6096	.
4.1	.6128	.6138	.6149	.6160	.6170	.6180	.6191	.6201	.
4.2	.6232	.6243	.6253	.6263	.6274	.6284	.6294	.6304	.
4.3	.6335	.6345	.6355	.6365	.6375	.6385	.6395	.6405	.
4.4	.6435	.6444	.6454	.6464	.6474	.6484	.6493	.6503	.
4.5	.6532	.6542	.6551	.6561	.6571	.6580	.6590	.6599	.
4.6	.6628	.6637	.6646	.6656	.6665	.6675	.6684	.6693	.
4.7	.6721	.6730	.6739	.6749	.6758	.6767	.6776	.6785	.
4.8	.6812	.6821	.6830	.6839	.6848	.6857	.6866	.6875	.
4.9	.6902	.6911	.6920	.6928	.6937	.6946	.6955	.6964	.
5.0	.6990	.6998	.7007	.7016	.7024	.7033	.7042	.7050	.
5.1	.7076	.7084	.7093	.7101	.7110	.7118	.7126	.7135	.
5.2	.7160	.7168	.7177	.7185	.7193	.7202	.7210	.7218	.
5.3	.7243	.7251	.7259	.7267	.7275	.7284	.7292	.7300	.
5.4	.7324	.7332	.7340	.7348	.7356	.7364	.7372	.7380	.

No.	0	1	2	3	4	5	6	7	

No.	0	1	2	3	4	5	6	7	
5.5	.7404	.7412	.7419	.7427	.7435	.7443	.7451	.7459	.7
5.6	.7482	.7490	.7497	.7505	.7513	.7520	.7528	.7536	.7
5.7	.7559	.7566	.7574	.7582	.7589	.7597	.7604	.7612	.7
5.8	.7634	.7642	.7649	.7657	.7664	.7672	.7679	.7686	.7
5.9	.7709	.7716	.7723	.7731	.7738	.7745	.7752	.7760	.7
6.0	.7782	.7789	.7796	.7803	.7810	.7818	.7825	.7832	.7
6.1	.7853	.7860	.7868	.7875	.7882	.7889	.7896	.7903	.7
6.2	.7924	.7931	.7938	.7945	.7952	.7959	.7966	.7973	.7
6.3	.7993	.8000	.8007	.8014	.8021	.8028	.8035	.8041	.8
6.4	.8062	.8069	.8075	.8082	.8089	.8096	.8102	.8109	.8
6.5	.8129	.8136	.8142	.8149	.8156	.8162	.8169	.8176	.8
6.6	.8195	.8202	.8209	.8215	.8222	.8228	.8235	.8241	.8
6.7	.8261	.8267	.8274	.8280	.8287	.8293	.8299	.8306	.8
6.8	.8325	.8331	.8338	.8344	.8351	.8357	.8363	.8370	.8
6.9	.8388	.8395	.8401	.8407	.8414	.8420	.8426	.8432	.8
7.0	.8451	.8457	.8463	.8470	.8476	.8482	.8488	.8494	.8
7.1	.8513	.8519	.8525	.8531	.8537	.8543	.8549	.8555	.8
7.2	.8573	.8579	.8585	.8591	.8597	.8603	.8609	.8615	.8
7.3	.8633	.8639	.8645	.8651	.8657	.8663	.8669	.8675	.8
7.4	.8692	.8698	.8704	.8710	.8716	.8722	.8727	.8733	.8
7.5	.8751	.8756	.8762	.8768	.8774	.8779	.8785	.8791	.8
7.6	.8808	.8814	.8820	.8825	.8831	.8837	.8842	.8848	.8
7.7	.8865	.8871	.8876	.8882	.8887	.8893	.8899	.8904	.8
7.8	.8921	.8927	.8932	.8938	.8943	.8949	.8954	.8960	.8
7.9	.8976	.8982	.8987	.8993	.8998	.9004	.9009	.9015	.9
8.0	.9031	.9036	.9042	.9047	.9053	.9058	.9063	.9069	.9
8.1	.9085	.9090	.9096	.9101	.9106	.9112	.9117	.9122	.9
8.2	.9138	.9143	.9149	.9154	.9159	.9165	.9170	.9175	.9
8.3	.9191	.9196	.9201	.9206	.9212	.9217	.9222	.9227	.9
8.4	.9243	.9248	.9253	.9258	.9263	.9269	.9274	.9279	.9
8.5	.9294	.9299	.9304	.9309	.9315	.9320	.9325	.9330	.9
8.6	.9345	.9350	.9355	.9360	.9365	.9370	.9375	.9380	.9
8.7	.9395	.9400	.9405	.9410	.9415	.9420	.9425	.9430	.9
8.8	.9445	.9450	.9455	.9460	.9465	.9469	.9474	.9479	.9
8.9	.9494	.9499	.9504	.9509	.9513	.9518	.9523	.9528	.9
9.0	.9542	.9547	.9552	.9557	.9562	.9566	.9571	.9576	.9
9.1	.9590	.9595	.9600	.9605	.9609	.9614	.9619	.9624	.9
9.2	.9638	.9643	.9647	.9652	.9657	.9661	.9666	.9671	.9
9.3	.9685	.9689	.9694	.9699	.9703	.9708	.9713	.9717	.9
9.4	.9731	.9736	.9741	.9745	.9750	.9754	.9759	.9763	.9
9.5	.9777	.9782	.9786	.9791	.9795	.9800	.9805	.9809	.9
9.6	.9823	.9827	.9832	.9836	.9841	.9845	.9850	.9854	.9
9.7	.9868	.9872	.9877	.9881	.9886	.9890	.9894	.9899	.9
9.8	.9912	.9917	.9921	.9926	.9930	.9934	.9939	.9943	.9
9.9	.9956	.9961	.9965	.9969	.9974	.9978	.9983	.9987	.9
No.	0	1	2	3	4	5	6	7	

APPENDIX II

NATURAL SINES AND COSINES

0°		1°		2°		3°		4°	
SIN	COS	SIN	COS	SIN	COS	SIN	COS	SIN	COS
0000000	1.00000	0.01745	0.99985	0.03490	0.99939	0.05234	0.99863	0.06976	0.99756
0.00029	1.00000	0.01774	0.99984	0.03519	0.99938	0.05263	0.99861	0.07005	0.99754
0.00058	1.00000	0.01803	0.99984	0.03548	0.99937	0.05292	0.99860	0.07034	0.99752
0.00087	1.00000	0.01832	0.99983	0.03577	0.99936	0.05321	0.99858	0.07063	0.99750
0.00116	1.00000	0.01862	0.99983	0.03606	0.99935	0.05350	0.99857	0.07092	0.99748
0.00145	1.00000	0.01891	0.99982	0.03635	0.99934	0.05379	0.99855	0.07121	0.99746
0.00175	1.00000	0.01920	0.99982	0.03664	0.99933	0.05408	0.99854	0.07150	0.99744
0.00204	1.00000	0.01949	0.99981	0.03693	0.99932	0.05437	0.99852	0.07179	0.99742
0.00233	1.00000	0.01978	0.99980	0.03723	0.99931	0.05466	0.99851	0.07208	0.99740
0.00262	1.00000	0.02007	0.99980	0.03752	0.99930	0.05495	0.99849	0.07237	0.99738
0.00291	1.00000	0.02036	0.99979	0.03781	0.99929	0.05524	0.99847	0.07266	0.99736
0.00320	0.99999	0.02065	0.99979	0.03810	0.99927	0.05553	0.99846	0.07295	0.99734
0.00349	0.99999	0.02094	0.99978	0.03839	0.99926	0.05582	0.99844	0.07324	0.99731
0.00378	0.99999	0.02123	0.99977	0.03868	0.99925	0.05611	0.99842	0.07353	0.99729
0.00407	0.99999	0.02152	0.99977	0.03897	0.99924	0.05640	0.99841	0.07382	0.99727
0.00436	0.99999	0.02181	0.99976	0.03926	0.99923	0.05669	0.99839	0.07411	0.99725
0.00465	0.99999	0.02211	0.99976	0.03955	0.99922	0.05698	0.99838	0.07440	0.99723
0.00495	0.99999	0.02240	0.99975	0.03984	0.99921	0.05727	0.99836	0.07469	0.99721
0.00524	0.99999	0.02269	0.99974	0.04013	0.99919	0.05756	0.99834	0.07498	0.99719
0.00553	0.99998	0.02298	0.99974	0.04042	0.99918	0.05785	0.99833	0.07527	0.99716
0.00582	0.99998	0.02327	0.99973	0.04071	0.99917	0.05814	0.99831	0.07556	0.99714
0.00611	0.99998	0.02356	0.99972	0.04100	0.99916	0.05844	0.99829	0.07585	0.99712
0.00640	0.99998	0.02385	0.99972	0.04129	0.99915	0.05873	0.99827	0.07614	0.99710
0.00669	0.99998	0.02414	0.99971	0.04159	0.99913	0.05902	0.99826	0.07643	0.99708
0.00698	0.99998	0.02443	0.99970	0.04188	0.99912	0.05931	0.99824	0.07672	0.99705
0.00727	0.99997	0.02472	0.99969	0.04217	0.99911	0.05960	0.99822	0.07701	0.99703
0.00756	0.99997	0.02501	0.99969	0.04246	0.99910	0.05989	0.99821	0.07730	0.99701
0.00785	0.99997	0.02530	0.99968	0.04275	0.99909	0.06018	0.99819	0.07759	0.99699
0.00814	0.99997	0.02560	0.99967	0.04304	0.99907	0.06047	0.99817	0.07788	0.99696
0.00844	0.99996	0.02589	0.99966	0.04333	0.99906	0.06076	0.99815	0.07817	0.99694
0.00873	0.99996	0.02618	0.99966	0.04362	0.99905	0.06105	0.99813	0.07846	0.99692
0.00902	0.99996	0.02647	0.99965	0.04391	0.99904	0.06134	0.99812	0.07875	0.99689
0.00931	0.99996	0.02676	0.99964	0.04420	0.99902	0.06163	0.99810	0.07904	0.99687
0.00960	0.99995	0.02705	0.99963	0.04449	0.99901	0.06192	0.99808	0.07933	0.99685
0.00989	0.99995	0.02734	0.99963	0.04478	0.99900	0.06221	0.99806	0.07962	0.99683
0.01018	0.99995	0.02763	0.99962	0.04507	0.99898	0.06250	0.99804	0.07991	0.99680
0.01047	0.99995	0.02792	0.99961	0.04536	0.99897	0.06279	0.99803	0.08020	0.99678
0.01076	0.99994	0.02821	0.99960	0.04565	0.99896	0.06308	0.99801	0.08049	0.99676
0.01105	0.99994	0.02850	0.99959	0.04594	0.99894	0.06337	0.99799	0.08078	0.99673
0.01134	0.99994	0.02879	0.99959	0.04623	0.99893	0.06366	0.99797	0.08107	0.99671
0.01164	0.99993	0.02908	0.99958	0.04653	0.99892	0.06395	0.99795	0.08136	0.99668
0.01193	0.99993	0.02938	0.99957	0.04682	0.99890	0.06424	0.99793	0.08165	0.99666
0.01222	0.99993	0.02967	0.99956	0.04711	0.99889	0.06453	0.99792	0.08194	0.99664
0.01251	0.99992	0.02996	0.99955	0.04740	0.99888	0.06482	0.99790	0.08223	0.99661
0.01280	0.99992	0.03025	0.99954	0.04769	0.99886	0.06511	0.99788	0.08252	0.99659
0.01309	0.99991	0.03054	0.99953	0.04798	0.99885	0.06540	0.99786	0.08281	0.99657
0.01338	0.99991	0.03083	0.99952	0.04827	0.99883	0.06569	0.99784	0.08310	0.99654
0.01367	0.99991	0.03112	0.99952	0.04856	0.99882	0.06598	0.99782	0.08339	0.99652
0.01396	0.99990	0.03141	0.99951	0.04885	0.99881	0.06627	0.99780	0.08368	0.99649
0.01425	0.99990	0.03170	0.99950	0.04914	0.99879	0.06656	0.99778	0.08397	0.99647
0.01454	0.99989	0.03199	0.99949	0.04943	0.99878	0.06685	0.99776	0.08426	0.99644
0.01483	0.99989	0.03228	0.99948	0.04972	0.99876	0.06714	0.99774	0.08455	0.99642
0.01513	0.99989	0.03257	0.99947	0.05001	0.99875	0.06743	0.99772	0.08484	0.99639
0.01542	0.99988	0.03286	0.99946	0.05030	0.99873	0.06773	0.99770	0.08513	0.99637
0.01571	0.99988	0.03316	0.99945	0.05059	0.99872	0.06802	0.99768	0.08542	0.99635
0.01600	0.99987	0.03345	0.99944	0.05088	0.99870	0.06831	0.99766	0.08571	0.99632
0.01629	0.99987	0.03374	0.99943	0.05117	0.99869	0.06860	0.99764	0.08600	0.99630
0.01658	0.99986	0.03403	0.99942	0.05146	0.99867	0.06889	0.99762	0.08629	0.99627
0.01687	0.99986	0.03432	0.99941	0.05175	0.99866	0.06918	0.99760	0.08658	0.99625
0.01716	0.99985	0.03461	0.99940	0.05205	0.99864	0.06947	0.99758	0.08687	0.99622
0.01745	0.99985	0.03490	0.99939	0.05234	0.99863	0.06976	0.99756	0.08716	0.99619
COS	SIN	COS	SIN	COS	SIN	COS	SIN	COS	SIN
89°		88°		87°		86°		85°	

MIN	5°		6°		7°		8°		
	SIN	COS	SIN	COS	SIN	COS	SIN	COS	SIN
0	0.08716	0.99619	0.10453	0.99452	0.12187	0.99255	0.13917	0.99027	0.15643
1	0.08745	0.99617	0.10482	0.99449	0.12216	0.99251	0.13946	0.99023	0.15672
2	0.08774	0.99614	0.10511	0.99446	0.12245	0.99248	0.13975	0.99019	0.15701
3	0.08803	0.99612	0.10540	0.99443	0.12274	0.99244	0.14004	0.99015	0.15730
4	0.08831	0.99609	0.10569	0.99440	0.12302	0.99240	0.14033	0.99011	0.15758
5	0.08860	0.99607	0.10597	0.99437	0.12331	0.99237	0.14061	0.99006	0.15787
6	0.08889	0.99604	0.10626	0.99434	0.12360	0.99233	0.14090	0.99002	0.15816
7	0.08918	0.99602	0.10655	0.99431	0.12389	0.99230	0.14119	0.98998	0.15845
8	0.08947	0.99599	0.10684	0.99428	0.12418	0.99226	0.14148	0.98994	0.15873
9	0.08976	0.99596	0.10713	0.99424	0.12447	0.99222	0.14177	0.98990	0.15902
10	0.09005	0.99594	0.10742	0.99421	0.12476	0.99219	0.14205	0.98986	0.15931
11	0.09034	0.99591	0.10771	0.99418	0.12504	0.99215	0.14234	0.98982	0.15959
12	0.09063	0.99588	0.10800	0.99415	0.12533	0.99211	0.14263	0.98978	0.15988
13	0.09092	0.99586	0.10829	0.99412	0.12562	0.99208	0.14292	0.98973	0.16017
14	0.09121	0.99583	0.10858	0.99409	0.12591	0.99204	0.14320	0.98969	0.16046
15	0.09150	0.99580	0.10887	0.99406	0.12620	0.99200	0.14349	0.98965	0.16074
16	0.09179	0.99578	0.10916	0.99402	0.12649	0.99197	0.14378	0.98961	0.16103
17	0.09208	0.99575	0.10945	0.99399	0.12678	0.99193	0.14407	0.98957	0.16132
18	0.09237	0.99572	0.10973	0.99396	0.12706	0.99189	0.14436	0.98953	0.16160
19	0.09266	0.99570	0.11002	0.99393	0.12735	0.99186	0.14464	0.98948	0.16189
20	0.09295	0.99567	0.11031	0.99390	0.12764	0.99182	0.14493	0.98944	0.16218
21	0.09324	0.99564	0.11060	0.99386	0.12793	0.99178	0.14522	0.98940	0.16246
22	0.09353	0.99562	0.11089	0.99383	0.12822	0.99175	0.14551	0.98936	0.16275
23	0.09382	0.99559	0.11118	0.99380	0.12851	0.99171	0.14580	0.98931	0.16304
24	0.09411	0.99556	0.11147	0.99377	0.12880	0.99167	0.14608	0.98927	0.16333
25	0.09440	0.99553	0.11176	0.99374	0.12908	0.99163	0.14637	0.98923	0.1636
26	0.09469	0.99551	0.11205	0.99370	0.12937	0.99160	0.14666	0.98919	0.16390
27	0.09498	0.99548	0.11234	0.99367	0.12966	0.99156	0.14695	0.98914	0.16419
28	0.09527	0.99545	0.11263	0.99364	0.12995	0.99152	0.14723	0.98910	0.16447
29	0.09556	0.99542	0.11291	0.99360	0.13024	0.99148	0.14752	0.98906	0.16476
30	0.09585	0.99540	0.11320	0.99357	0.13053	0.99144	0.14781	0.98902	0.16505
31	0.09614	0.99537	0.11349	0.99354	0.13081	0.99141	0.14810	0.98897	0.16533
32	0.09642	0.99534	0.11378	0.99351	0.13110	0.99137	0.14838	0.98893	0.16562
33	0.09671	0.99531	0.11407	0.99347	0.13139	0.99133	0.14867	0.98889	0.1659
34	0.09700	0.99528	0.11436	0.99344	0.13168	0.99129	0.14896	0.98884	0.16620
35	0.09729	0.99526	0.11465	0.99341	0.13197	0.99125	0.14925	0.98880	0.16648
36	0.09758	0.99523	0.11494	0.99337	0.13226	0.99122	0.14954	0.98876	0.16677
37	0.09787	0.99520	0.11523	0.99334	0.13254	0.99118	0.14982	0.98871	0.16706
38	0.09816	0.99517	0.11552	0.99331	0.13283	0.99114	0.15011	0.98867	0.16735
39	0.09845	0.99514	0.11580	0.99327	0.13312	0.99110	0.15040	0.98863	0.16764
40	0.09874	0.99511	0.11609	0.99324	0.13341	0.99106	0.15069	0.98858	0.16793
41	0.09903	0.99508	0.11638	0.99320	0.13370	0.99102	0.15097	0.98854	0.16822
42	0.09932	0.99506	0.11667	0.99317	0.13399	0.99098	0.15126	0.98849	0.16851
43	0.09961	0.99503	0.11696	0.99314	0.13427	0.99094	0.15155	0.98845	0.16879
44	0.09990	0.99500	0.11725	0.99310	0.13456	0.99091	0.15184	0.98841	0.16908
45	0.10019	0.99497	0.11754	0.99307	0.13485	0.99087	0.15212	0.98836	0.16937
46	0.10048	0.99494	0.11783	0.99303	0.13514	0.99083	0.15241	0.98832	0.16966
47	0.10077	0.99491	0.11812	0.99300	0.13543	0.99079	0.15270	0.98827	0.16995
48	0.10106	0.99488	0.11840	0.99297	0.13572	0.99075	0.15299	0.98823	0.17024
49	0.10135	0.99485	0.11869	0.99293	0.13600	0.99071	0.15327	0.98818	0.17053
50	0.10164	0.99482	0.11898	0.99290	0.13629	0.99067	0.15356	0.98814	0.17081
51	0.10192	0.99479	0.11927	0.99286	0.13658	0.99063	0.15385	0.98809	0.17110
52	0.10221	0.99476	0.11956	0.99283	0.13687	0.99059	0.15414	0.98805	0.17139
53	0.10250	0.99473	0.11985	0.99279	0.13716	0.99055	0.15442	0.98800	0.17168
54	0.10279	0.99470	0.12014	0.99276	0.13744	0.99051	0.15471	0.98796	0.17197
55	0.10308	0.99467	0.12043	0.99272	0.13773	0.99047	0.15500	0.98791	0.17226
56	0.10337	0.99464	0.12071	0.99269	0.13802	0.99043	0.15529	0.98787	0.17255
57	0.10366	0.99461	0.12100	0.99265	0.13831	0.99039	0.15557	0.98782	0.17284
58	0.10395	0.99458	0.12129	0.99262	0.13860	0.99035	0.15586	0.98778	0.17313
59	0.10424	0.99455	0.12158	0.99258	0.13889	0.99031	0.15615	0.98773	0.17342
60	0.10453	0.99452	0.12187	0.99255	0.13917	0.99027	0.15643	0.98769	0.17371

	COS	SIN	COS	SIN	COS	SIN	COS	SIN	COS
	84°		83°		82°		81°		

MIN	10°		11°		12°		13°		1
	SIN	COS	SIN	COS	SIN	COS	SIN	COS	SIN
0	0.17365	0.98481	0.19081	0.98163	0.20791	0.97815	0.22495	0.97437	0.24192
1	0.17393	0.98476	0.19109	0.98157	0.20820	0.97809	0.22523	0.97430	0.24220
2	0.17422	0.98471	0.19138	0.98152	0.20848	0.97803	0.22552	0.97424	0.24249
3	0.17451	0.98466	0.19167	0.98146	0.20877	0.97797	0.22580	0.97417	0.24277
4	0.17479	0.98461	0.19195	0.98140	0.20905	0.97791	0.22608	0.97411	0.24305
5	0.17508	0.98455	0.19224	0.98135	0.20933	0.97784	0.22637	0.97404	0.24333
6	0.17537	0.98450	0.19252	0.98129	0.20962	0.97778	0.22665	0.97398	0.24362
7	0.17565	0.98445	0.19281	0.98124	0.20990	0.97772	0.22693	0.97391	0.24390
8	0.17594	0.98440	0.19309	0.98118	0.21019	0.97766	0.22722	0.97384	0.24418
9	0.17623	0.98435	0.19338	0.98112	0.21047	0.97760	0.22750	0.97378	0.24446
10	0.17651	0.98430	0.19366	0.98107	0.21076	0.97754	0.22778	0.97371	0.24474
11	0.17680	0.98425	0.19395	0.98101	0.21104	0.97748	0.22807	0.97365	0.24503
12	0.17708	0.98420	0.19423	0.98096	0.21132	0.97742	0.22835	0.97358	0.24531
13	0.17737	0.98414	0.19452	0.98090	0.21161	0.97735	0.22863	0.97351	0.24559
14	0.17766	0.98409	0.19481	0.98084	0.21189	0.97729	0.22892	0.97345	0.24587
15	0.17794	0.98404	0.19509	0.98079	0.21218	0.97723	0.22920	0.97338	0.24615
16	0.17823	0.98399	0.19538	0.98073	0.21246	0.97717	0.22948	0.97331	0.24644
17	0.17852	0.98394	0.19566	0.98067	0.21275	0.97711	0.22977	0.97325	0.24672
18	0.17880	0.98389	0.19595	0.98061	0.21303	0.97705	0.23005	0.97318	0.24700
19	0.17909	0.98383	0.19623	0.98056	0.21331	0.97698	0.23033	0.97311	0.24728
20	0.17937	0.98378	0.19652	0.98050	0.21360	0.97692	0.23062	0.97304	0.24756
21	0.17966	0.98373	0.19680	0.98044	0.21388	0.97686	0.23090	0.97298	0.24784
22	0.17995	0.98368	0.19709	0.98039	0.21417	0.97680	0.23118	0.97291	0.24813
23	0.18023	0.98362	0.19737	0.98033	0.21445	0.97673	0.23146	0.97284	0.24841
24	0.18052	0.98357	0.19766	0.98027	0.21474	0.97667	0.23175	0.97278	0.24869
25	0.18081	0.98352	0.19794	0.98021	0.21502	0.97661	0.23203	0.97271	0.24897
26	0.18109	0.98347	0.19823	0.98016	0.21530	0.97655	0.23231	0.97264	0.24925
27	0.18138	0.98341	0.19851	0.98010	0.21559	0.97648	0.23260	0.97257	0.24954
28	0.18166	0.98336	0.19880	0.98004	0.21587	0.97642	0.23288	0.97251	0.24982
29	0.18195	0.98331	0.19908	0.97998	0.21616	0.97636	0.23316	0.97244	0.25010
30	0.18224	0.98325	0.19937	0.97992	0.21644	0.97630	0.23345	0.97237	0.25038
31	0.18252	0.98320	0.19965	0.97987	0.21672	0.97623	0.23373	0.97230	0.25066
32	0.18281	0.98315	0.19994	0.97981	0.21701	0.97617	0.23401	0.97223	0.25094
33	0.18309	0.98310	0.20022	0.97975	0.21729	0.97611	0.23429	0.97217	0.25122
34	0.18338	0.98304	0.20051	0.97969	0.21758	0.97604	0.23458	0.97210	0.25151
35	0.18367	0.98299	0.20079	0.97963	0.21786	0.97598	0.23486	0.97203	0.25179
36	0.18395	0.98294	0.20108	0.97958	0.21814	0.97592	0.23514	0.97196	0.25207
37	0.18424	0.98288	0.20136	0.97952	0.21843	0.97585	0.23542	0.97189	0.25235
38	0.18452	0.98283	0.20165	0.97946	0.21871	0.97579	0.23571	0.97182	0.25263
39	0.18481	0.98277	0.20193	0.97940	0.21899	0.97573	0.23599	0.97176	0.25291
40	0.18509	0.98272	0.20222	0.97934	0.21928	0.97566	0.23627	0.97169	0.25320
41	0.18538	0.98267	0.20250	0.97928	0.21956	0.97560	0.23656	0.97162	0.25348
42	0.18567	0.98261	0.20279	0.97922	0.21985	0.97553	0.23684	0.97155	0.25376
43	0.18595	0.98256	0.20307	0.97916	0.22013	0.97547	0.23712	0.97148	0.25404
44	0.18624	0.98250	0.20336	0.97910	0.22041	0.97541	0.23740	0.97141	0.25432
45	0.18652	0.98245	0.20364	0.97905	0.22070	0.97534	0.23769	0.97134	0.25460
46	0.18681	0.98240	0.20393	0.97899	0.22098	0.97528	0.23797	0.97127	0.25488
47	0.18710	0.98234	0.20421	0.97893	0.22126	0.97521	0.23825	0.97120	0.25516
48	0.18738	0.98229	0.20450	0.97887	0.22155	0.97515	0.23853	0.97113	0.25545
49	0.18767	0.98223	0.20478	0.97881	0.22183	0.97508	0.23882	0.97106	0.25573
50	0.18795	0.98218	0.20507	0.97875	0.22212	0.97502	0.23910	0.97100	0.25601
51	0.18824	0.98212	0.20535	0.97869	0.22240	0.97496	0.23938	0.97093	0.25629
52	0.18852	0.98207	0.20563	0.97863	0.22268	0.97489	0.23966	0.97086	0.25657
53	0.18881	0.98201	0.20592	0.97857	0.22297	0.97483	0.23995	0.97079	0.25685
54	0.18910	0.98196	0.20620	0.97851	0.22325	0.97476	0.24023	0.97072	0.25713
55	0.18938	0.98190	0.20649	0.97845	0.22353	0.97470	0.24051	0.97065	0.2574
56	0.18967	0.98185	0.20677	0.97839	0.22382	0.97463	0.24079	0.97058	0.2576
57	0.18995	0.98179	0.20706	0.97833	0.22410	0.97457	0.24108	0.97051	0.2579
58	0.19024	0.98174	0.20734	0.97827	0.22438	0.97450	0.24136	0.97044	0.2582
59	0.19052	0.98168	0.20763	0.97821	0.22467	0.97444	0.24164	0.97037	0.2585
60	0.19081	0.98163	0.20791	0.97815	0.22495	0.97437	0.24192	0.97030	0.2588

	COS	SIN	COS	SIN	COS	SIN	COS	SIN	COS
	79°		78°		77°		76°		7

MIN	15°		16°		17°		18°		1°
	SIN	COS	SIN	COS	SIN	COS	SIN	COS	SIN
0	0.25882	0.96593	0.27564	0.96126	0.29237	0.95630	0.30902	0.95106	0.32557
1	0.25910	0.96585	0.27592	0.96118	0.29265	0.95622	0.30929	0.95097	0.32584
2	0.25938	0.96578	0.27620	0.96110	0.29293	0.95613	0.30957	0.95088	0.32612
3	0.25966	0.96570	0.27648	0.96102	0.29321	0.95605	0.30985	0.95079	0.32639
4	0.25994	0.96562	0.27676	0.96094	0.29348	0.95596	0.31012	0.95070	0.32667
5	0.26022	0.96555	0.27704	0.96086	0.29376	0.95588	0.31040	0.95061	0.32694
6	0.26050	0.96547	0.27731	0.96078	0.29404	0.95579	0.31068	0.95052	0.32722
7	0.26079	0.96540	0.27759	0.96070	0.29432	0.95571	0.31095	0.95043	0.32749
8	0.26107	0.96532	0.27787	0.96062	0.29460	0.95562	0.31123	0.95033	0.32777
9	0.26135	0.96524	0.27815	0.96054	0.29487	0.95554	0.31151	0.95024	0.32804
10	0.26163	0.96517	0.27843	0.96046	0.29515	0.95545	0.31178	0.95015	0.32832
11	0.26191	0.96509	0.27871	0.96037	0.29543	0.95536	0.31206	0.95006	0.32859
12	0.26219	0.96502	0.27899	0.96029	0.29571	0.95528	0.31233	0.94997	0.32887
13	0.26247	0.96494	0.27927	0.96021	0.29599	0.95519	0.31261	0.94988	0.32914
14	0.26275	0.96486	0.27955	0.96013	0.29626	0.95511	0.31289	0.94979	0.32942
15	0.26303	0.96479	0.27983	0.96005	0.29654	0.95502	0.31316	0.94970	0.32969
16	0.26331	0.96471	0.28011	0.95997	0.29682	0.95493	0.31344	0.94961	0.32997
17	0.26359	0.96463	0.28039	0.95989	0.29710	0.95485	0.31372	0.94952	0.33024
18	0.26387	0.96456	0.28067	0.95981	0.29737	0.95476	0.31399	0.94943	0.33051
19	0.26415	0.96448	0.28095	0.95972	0.29765	0.95467	0.31427	0.94933	0.33079
20	0.26443	0.96440	0.28123	0.95964	0.29793	0.95459	0.31454	0.94924	0.33106
21	0.26471	0.96433	0.28150	0.95956	0.29821	0.95450	0.31482	0.94915	0.33134
22	0.26500	0.96425	0.28178	0.95948	0.29849	0.95441	0.31510	0.94906	0.33161
23	0.26528	0.96417	0.28206	0.95940	0.29876	0.95433	0.31537	0.94897	0.33189
24	0.26556	0.96410	0.28234	0.95931	0.29904	0.95424	0.31565	0.94888	0.33216
25	0.26584	0.96402	0.28262	0.95923	0.29932	0.95415	0.31593	0.94878	0.33244
26	0.26612	0.96394	0.28290	0.95915	0.29960	0.95407	0.31620	0.94869	0.33271
27	0.26640	0.96386	0.28318	0.95907	0.29987	0.95398	0.31648	0.94860	0.33298
28	0.26668	0.96379	0.28346	0.95898	0.30015	0.95389	0.31675	0.94851	0.33326
29	0.26696	0.96371	0.28374	0.95890	0.30043	0.95380	0.31703	0.94842	0.33353
30	0.26724	0.96363	0.28402	0.95882	0.30071	0.95372	0.31730	0.94832	0.33381
31	0.26752	0.96355	0.28429	0.95874	0.30098	0.95363	0.31758	0.94823	0.33408
32	0.26780	0.96347	0.28457	0.95865	0.30126	0.95354	0.31786	0.94814	0.33436
33	0.26808	0.96340	0.28485	0.95857	0.30154	0.95345	0.31813	0.94805	0.33463
34	0.26836	0.96332	0.28513	0.95849	0.30182	0.95337	0.31841	0.94795	0.33490
35	0.26864	0.96324	0.28541	0.95841	0.30209	0.95328	0.31868	0.94786	0.33518
36	0.26892	0.96316	0.28569	0.95832	0.30237	0.95319	0.31896	0.94777	0.33545
37	0.26920	0.96308	0.28597	0.95824	0.30265	0.95310	0.31923	0.94768	0.33573
38	0.26948	0.96301	0.28625	0.95816	0.30292	0.95301	0.31951	0.94758	0.33600
39	0.26976	0.96293	0.28652	0.95807	0.30320	0.95293	0.31979	0.94749	0.33627
40	0.27004	0.96285	0.28680	0.95799	0.30348	0.95284	0.32006	0.94740	0.33655
41	0.27032	0.96277	0.28708	0.95791	0.30376	0.95275	0.32034	0.94730	0.33682
42	0.27060	0.96269	0.28736	0.95782	0.30403	0.95266	0.32061	0.94721	0.33710
43	0.27088	0.96261	0.28764	0.95774	0.30431	0.95257	0.32089	0.94712	0.33737
44	0.27116	0.96253	0.28792	0.95766	0.30459	0.95248	0.32116	0.94702	0.33764
45	0.27144	0.96246	0.28820	0.95757	0.30486	0.95240	0.32144	0.94693	0.33792
46	0.27172	0.96238	0.28847	0.95749	0.30514	0.95231	0.32171	0.94684	0.33819
47	0.27200	0.96230	0.28875	0.95740	0.30542	0.95222	0.32199	0.94674	0.33846
48	0.27228	0.96222	0.28903	0.95732	0.30570	0.95213	0.32227	0.94665	0.33874
49	0.27256	0.96214	0.28931	0.95724	0.30597	0.95204	0.32254	0.94656	0.33901
50	0.27284	0.96206	0.28959	0.95715	0.30625	0.95195	0.32282	0.94646	0.33929
51	0.27312	0.96198	0.28987	0.95707	0.30653	0.95186	0.32309	0.94637	0.33956
52	0.27340	0.96190	0.29015	0.95698	0.30680	0.95177	0.32337	0.94627	0.33983
53	0.27368	0.96182	0.29042	0.95690	0.30708	0.95168	0.32364	0.94618	0.34011
54	0.27396	0.96174	0.29070	0.95681	0.30736	0.95159	0.32392	0.94609	0.34038
55	0.27424	0.96166	0.29098	0.95673	0.30763	0.95150	0.32419	0.94599	0.34065
56	0.27452	0.96158	0.29126	0.95664	0.30791	0.95142	0.32447	0.94590	0.34093
57	0.27480	0.96150	0.29154	0.95656	0.30819	0.95133	0.32474	0.94580	0.34120
58	0.27508	0.96142	0.29182	0.95647	0.30846	0.95124	0.32502	0.94571	0.34147
59	0.27536	0.96134	0.29209	0.95639	0.30874	0.95115	0.32529	0.94561	0.34175
60	0.27564	0.96126	0.29237	0.95630	0.30902	0.95106	0.32557	0.94552	0.34202
	COS	SIN	COS	SIN	COS	SIN	COS	SIN	COS
	74°		73°		72°		71°		7

MIN	20°		21°		22°		23°		24
	SIN	COS	SIN	COS	SIN	COS	SIN	COS	SIN
0	0.34202	0.93969	0.35837	0.93358	0.37461	0.92718	0.39073	0.92050	0.40674
1	0.34229	0.93959	0.35864	0.93348	0.37488	0.92707	0.39100	0.92039	0.40700
2	0.34257	0.93949	0.35891	0.93337	0.37515	0.92697	0.39127	0.92028	0.40727
3	0.34284	0.93939	0.35918	0.93327	0.37542	0.92686	0.39153	0.92016	0.40753
4	0.34311	0.93929	0.35945	0.93316	0.37569	0.92675	0.39180	0.92005	0.40780
5	0.34339	0.93919	0.35973	0.93306	0.37595	0.92664	0.39207	0.91994	0.40806
6	0.34366	0.93909	0.36000	0.93295	0.37622	0.92653	0.39234	0.91982	0.40833
7	0.34393	0.93899	0.36027	0.93285	0.37649	0.92642	0.39260	0.91971	0.40860
8	0.34421	0.93889	0.36054	0.93274	0.37676	0.92631	0.39287	0.91959	0.40886
9	0.34448	0.93879	0.36081	0.93264	0.37703	0.92620	0.39314	0.91948	0.40913
10	0.34475	0.93869	0.36108	0.93253	0.37730	0.92609	0.39341	0.91936	0.40939
11	0.34503	0.93859	0.36135	0.93243	0.37757	0.92598	0.39367	0.91925	0.40966
12	0.34530	0.93849	0.36162	0.93232	0.37784	0.92587	0.39394	0.91914	0.40992
13	0.34557	0.93839	0.36190	0.93222	0.37811	0.92576	0.39421	0.91902	0.41019
14	0.34584	0.93829	0.36217	0.93211	0.37838	0.92565	0.39448	0.91891	0.41045
15	0.34612	0.93819	0.36244	0.93201	0.37865	0.92554	0.39474	0.91879	0.41072
16	0.34639	0.93809	0.36271	0.93190	0.37892	0.92543	0.39501	0.91868	0.41098
17	0.34666	0.93799	0.36298	0.93180	0.37919	0.92532	0.39528	0.91856	0.41125
18	0.34694	0.93789	0.36325	0.93169	0.37946	0.92521	0.39555	0.91845	0.41151
19	0.34721	0.93779	0.36352	0.93159	0.37973	0.92510	0.39581	0.91833	0.41178
20	0.34748	0.93769	0.36379	0.93148	0.37999	0.92499	0.39608	0.91822	0.41204
21	0.34775	0.93759	0.36406	0.93137	0.38026	0.92488	0.39635	0.91810	0.41231
22	0.34803	0.93748	0.36434	0.93127	0.38053	0.92477	0.39661	0.91799	0.41257
23	0.34830	0.93738	0.36461	0.93116	0.38080	0.92466	0.39688	0.91787	0.41284
24	0.34857	0.93728	0.36488	0.93106	0.38107	0.92455	0.39715	0.91775	0.41310
25	0.34884	0.93718	0.36515	0.93095	0.38134	0.92444	0.39741	0.91764	0.41337
26	0.34912	0.93708	0.36542	0.93084	0.38161	0.92432	0.39768	0.91752	0.41363
27	0.34939	0.93698	0.36569	0.93074	0.38188	0.92421	0.39795	0.91741	0.41390
28	0.34966	0.93688	0.36596	0.93063	0.38215	0.92410	0.39822	0.91729	0.41416
29	0.34993	0.93677	0.36623	0.93052	0.38241	0.92399	0.39848	0.91718	0.41443
30	0.35021	0.93667	0.36650	0.93042	0.38268	0.92388	0.39875	0.91706	0.41469
31	0.35048	0.93657	0.36677	0.93031	0.38295	0.92377	0.39902	0.91694	0.41496
32	0.35075	0.93647	0.36704	0.93020	0.38322	0.92366	0.39928	0.91683	0.41522
33	0.35102	0.93637	0.36731	0.93010	0.38349	0.92355	0.39955	0.91671	0.41549
34	0.35130	0.93626	0.36758	0.92999	0.38376	0.92343	0.39982	0.91660	0.41575
35	0.35157	0.93616	0.36785	0.92988	0.38403	0.92332	0.40008	0.91648	0.41602
36	0.35184	0.93606	0.36812	0.92978	0.38430	0.92321	0.40035	0.91636	0.41628
37	0.35211	0.93596	0.36839	0.92967	0.38456	0.92310	0.40062	0.91625	0.41655
38	0.35239	0.93585	0.36867	0.92956	0.38483	0.92299	0.40088	0.91613	0.41681
39	0.35266	0.93575	0.36894	0.92945	0.38510	0.92287	0.40115	0.91601	0.41707
40	0.35293	0.93565	0.36921	0.92935	0.38537	0.92276	0.40141	0.91590	0.41734
41	0.35320	0.93555	0.36948	0.92924	0.38564	0.92265	0.40168	0.91578	0.41760
42	0.35347	0.93544	0.36975	0.92913	0.38591	0.92254	0.40195	0.91566	0.41787
43	0.35375	0.93534	0.37002	0.92902	0.38617	0.92243	0.40221	0.91555	0.41813
44	0.35402	0.93524	0.37029	0.92892	0.38644	0.92231	0.40248	0.91543	0.41840
45	0.35429	0.93514	0.37056	0.92881	0.38671	0.92220	0.40275	0.91531	0.41866
46	0.35456	0.93503	0.37083	0.92870	0.38698	0.92209	0.40301	0.91519	0.41892
47	0.35484	0.93493	0.37110	0.92859	0.38725	0.92198	0.40328	0.91508	0.41919
48	0.35511	0.93483	0.37137	0.92849	0.38752	0.92186	0.40355	0.91496	0.41945
49	0.35538	0.93472	0.37164	0.92838	0.38778	0.92175	0.40381	0.91484	0.41972
50	0.35565	0.93462	0.37191	0.92827	0.38805	0.92164	0.40408	0.91472	0.41998
51	0.35592	0.93452	0.37218	0.92816	0.38832	0.92152	0.40434	0.91461	0.42024
52	0.35619	0.93441	0.37245	0.92805	0.38859	0.92141	0.40461	0.91449	0.42051
53	0.35647	0.93431	0.37272	0.92794	0.38886	0.92130	0.40488	0.91437	0.42077
54	0.35674	0.93420	0.37299	0.92784	0.38912	0.92119	0.40514	0.91425	0.42104
55	0.35701	0.93410	0.37326	0.92773	0.38939	0.92107	0.40541	0.91414	0.42130
56	0.35728	0.93400	0.37353	0.92762	0.38966	0.92096	0.40567	0.91402	0.42156
57	0.35755	0.93389	0.37380	0.92751	0.38993	0.92085	0.40594	0.91390	0.42183
58	0.35782	0.93379	0.37407	0.92740	0.39020	0.92073	0.40621	0.91378	0.42209
59	0.35810	0.93368	0.37434	0.92729	0.39046	0.92062	0.40647	0.91366	0.42235
60	0.35837	0.93358	0.37461	0.92718	0.39073	0.92050	0.40674	0.91355	0.42262

	COS	SIN	COS	SIN	COS	SIN	COS	SIN	COS
	69°		68°		67°		66°		6

MIN	25°		26°		27°		28°		29
	SIN	COS	SIN	COS	SIN	COS	SIN	COS	SIN
0	0.42262	0.90631	0.43837	0.89879	0.45399	0.89101	0.46947	0.88295	0.48481
1	0.42288	0.90618	0.43863	0.89867	0.45425	0.89087	0.46973	0.88281	0.48506
2	0.42315	0.90606	0.43889	0.89854	0.45451	0.89074	0.46999	0.88267	0.48532
3	0.42341	0.90594	0.43916	0.89841	0.45477	0.89061	0.47024	0.88254	0.48557
4	0.42367	0.90582	0.43942	0.89828	0.45503	0.89048	0.47050	0.88240	0.48583
5	0.42394	0.90569	0.43968	0.89816	0.45529	0.89035	0.47076	0.88226	0.48608
6	0.42420	0.90557	0.43994	0.89803	0.45554	0.89021	0.47101	0.88213	0.48634
7	0.42446	0.90545	0.44020	0.89790	0.45580	0.89008	0.47127	0.88199	0.48659
8	0.42473	0.90532	0.44046	0.89777	0.45606	0.88995	0.47153	0.88185	0.48684
9	0.42499	0.90520	0.44072	0.89764	0.45632	0.88981	0.47178	0.88172	0.48710
10	0.42525	0.90507	0.44098	0.89752	0.45658	0.88968	0.47204	0.88158	0.48735
11	0.42552	0.90495	0.44124	0.89739	0.45684	0.88955	0.47229	0.88144	0.48761
12	0.42578	0.90483	0.44151	0.89726	0.45710	0.88942	0.47255	0.88130	0.48786
13	0.42604	0.90470	0.44177	0.89713	0.45736	0.88928	0.47281	0.88117	0.48811
14	0.42631	0.90458	0.44203	0.89700	0.45762	0.88915	0.47306	0.88103	0.48837
15	0.42657	0.90446	0.44229	0.89687	0.45787	0.88902	0.47332	0.88089	0.48862
16	0.42683	0.90433	0.44255	0.89674	0.45813	0.88888	0.47358	0.88075	0.48888
17	0.42709	0.90421	0.44281	0.89662	0.45839	0.88875	0.47383	0.88062	0.48913
18	0.42736	0.90408	0.44307	0.89649	0.45865	0.88862	0.47409	0.88048	0.48938
19	0.42762	0.90396	0.44333	0.89636	0.45891	0.88848	0.47434	0.88034	0.48964
20	0.42788	0.90383	0.44359	0.89623	0.45917	0.88835	0.47460	0.88020	0.48989
21	0.42815	0.90371	0.44385	0.89610	0.45942	0.88822	0.47486	0.88006	0.49014
22	0.42841	0.90358	0.44411	0.89597	0.45968	0.88808	0.47511	0.87993	0.49040
23	0.42867	0.90346	0.44437	0.89584	0.45994	0.88795	0.47537	0.87979	0.49065
24	0.42894	0.90334	0.44464	0.89571	0.46020	0.88782	0.47562	0.87965	0.49090
25	0.42920	0.90321	0.44490	0.89558	0.46046	0.88768	0.47588	0.87951	0.49116
26	0.42946	0.90309	0.44516	0.89545	0.46072	0.88755	0.47614	0.87937	0.49141
27	0.42972	0.90296	0.44542	0.89532	0.46097	0.88741	0.47639	0.87923	0.49166
28	0.42999	0.90284	0.44568	0.89519	0.46123	0.88728	0.47665	0.87909	0.49192
29	0.43025	0.90271	0.44594	0.89506	0.46149	0.88715	0.47690	0.87896	0.49217
30	0.43051	0.90259	0.44620	0.89493	0.46175	0.88701	0.47716	0.87882	0.49242
31	0.43077	0.90246	0.44646	0.89480	0.46201	0.88688	0.47741	0.87868	0.49268
32	0.43104	0.90233	0.44672	0.89467	0.46226	0.88674	0.47767	0.87854	0.49293
33	0.43130	0.90221	0.44698	0.89454	0.46252	0.88661	0.47793	0.87840	0.49318
34	0.43156	0.90208	0.44724	0.89441	0.46278	0.88647	0.47818	0.87826	0.49344
35	0.43182	0.90196	0.44750	0.89428	0.46304	0.88634	0.47844	0.87812	0.49369
36	0.43209	0.90183	0.44776	0.89415	0.46330	0.88620	0.47869	0.87798	0.49394
37	0.43235	0.90171	0.44802	0.89402	0.46355	0.88607	0.47895	0.87784	0.49419
38	0.43261	0.90158	0.44828	0.89389	0.46381	0.88593	0.47920	0.87770	0.49445
39	0.43287	0.90146	0.44854	0.89376	0.46407	0.88580	0.47946	0.87756	0.49470
40	0.43313	0.90133	0.44880	0.89363	0.46433	0.88566	0.47971	0.87743	0.49495
41	0.43340	0.90120	0.44906	0.89350	0.46458	0.88553	0.47997	0.87729	0.49521
42	0.43366	0.90108	0.44932	0.89337	0.46484	0.88539	0.48022	0.87715	0.49546
43	0.43392	0.90095	0.44958	0.89324	0.46510	0.88526	0.48048	0.87701	0.49571
44	0.43418	0.90082	0.44984	0.89311	0.46536	0.88512	0.48073	0.87687	0.49596
45	0.43445	0.90070	0.45010	0.89298	0.46561	0.88499	0.48099	0.87673	0.49622
46	0.43471	0.90057	0.45036	0.89285	0.46587	0.88485	0.48124	0.87659	0.49647
47	0.43497	0.90045	0.45062	0.89272	0.46613	0.88472	0.48150	0.87645	0.49672
48	0.43523	0.90032	0.45088	0.89259	0.46639	0.88458	0.48175	0.87631	0.49697
49	0.43549	0.90019	0.45114	0.89245	0.46664	0.88445	0.48201	0.87617	0.49723
50	0.43575	0.90007	0.45140	0.89232	0.46690	0.88431	0.48226	0.87603	0.49748
51	0.43602	0.89994	0.45166	0.89219	0.46716	0.88417	0.48252	0.87589	0.49773
52	0.43628	0.89981	0.45192	0.89206	0.46742	0.88404	0.48277	0.87575	0.49798
53	0.43654	0.89968	0.45218	0.89193	0.46767	0.88390	0.48303	0.87561	0.49824
54	0.43680	0.89956	0.45243	0.89180	0.46793	0.88377	0.48328	0.87546	0.49849
55	0.43706	0.89943	0.45269	0.89167	0.46819	0.88363	0.48354	0.87532	0.49874
56	0.43733	0.89930	0.45295	0.89153	0.46844	0.88349	0.48379	0.87518	0.49899
57	0.43759	0.89918	0.45321	0.89140	0.46870	0.88336	0.48405	0.87504	0.49924
58	0.43785	0.89905	0.45347	0.89127	0.46896	0.88322	0.48430	0.87490	0.49950
59	0.43811	0.89892	0.45373	0.89114	0.46921	0.88308	0.48456	0.87476	0.49975
60	0.43837	0.89879	0.45399	0.89101	0.46947	0.88295	0.48481	0.87462	0.50000
	COS	SIN	COS	SIN	COS	SIN	COS	SIN	COS
	64°		63°		62°		61°		6

30°		31°		32°		33°		34°	
SIN	COS	SIN	COS	SIN	COS	SIN	COS	SIN	COS
0.50000	0.86603	0.51504	0.85717	0.52992	0.84805	0.54464	0.83867	0.55919	0.82904
0.50025	0.86588	0.51529	0.85702	0.53017	0.84789	0.54488	0.83851	0.55943	0.82887
0.50050	0.86573	0.51554	0.85687	0.53041	0.84774	0.54513	0.83835	0.55968	0.82871
0.50076	0.86559	0.51579	0.85672	0.53066	0.84759	0.54537	0.83819	0.55992	0.82855
0.50101	0.86544	0.51604	0.85657	0.53091	0.84743	0.54561	0.83804	0.56016	0.82839
0.50126	0.86530	0.51628	0.85642	0.53115	0.84728	0.54586	0.83788	0.56040	0.82822
0.50151	0.86515	0.51653	0.85627	0.53140	0.84712	0.54610	0.83772	0.56064	0.82806
0.50176	0.86501	0.51678	0.85612	0.53164	0.84697	0.54635	0.83756	0.56088	0.82790
0.50201	0.86486	0.51703	0.85597	0.53189	0.84681	0.54659	0.83740	0.56112	0.82773
0.50227	0.86471	0.51728	0.85582	0.53214	0.84666	0.54683	0.83724	0.56136	0.82757
0.50252	0.86457	0.51753	0.85567	0.53238	0.84650	0.54708	0.83708	0.56160	0.82741
0.50277	0.86442	0.51778	0.85551	0.53263	0.84635	0.54732	0.83692	0.56184	0.82724
0.50302	0.86427	0.51803	0.85536	0.53288	0.84619	0.54756	0.83676	0.56208	0.82708
0.50327	0.86413	0.51828	0.85521	0.53312	0.84604	0.54781	0.83660	0.56232	0.82692
0.50352	0.86398	0.51852	0.85506	0.53337	0.84588	0.54805	0.83645	0.56256	0.82675
0.50377	0.86384	0.51877	0.85491	0.53361	0.84573	0.54829	0.83629	0.56280	0.82659
0.50403	0.86369	0.51902	0.85476	0.53386	0.84557	0.54854	0.83613	0.56305	0.82643
0.50428	0.86354	0.51927	0.85461	0.53411	0.84542	0.54878	0.83597	0.56329	0.82626
0.50453	0.86340	0.51952	0.85446	0.53435	0.84526	0.54902	0.83581	0.56353	0.82610
0.50478	0.86325	0.51977	0.85431	0.53460	0.84511	0.54927	0.83565	0.56377	0.82593
0.50503	0.86310	0.52002	0.85416	0.53484	0.84495	0.54951	0.83549	0.56401	0.82577
0.50528	0.86295	0.52026	0.85401	0.53509	0.84480	0.54975	0.83533	0.56425	0.82561
0.50553	0.86281	0.52051	0.85385	0.53534	0.84464	0.54999	0.83517	0.56449	0.82544
0.50578	0.86266	0.52076	0.85370	0.53558	0.84448	0.55024	0.83501	0.56473	0.82528
0.50603	0.86251	0.52101	0.85355	0.53583	0.84433	0.55048	0.83485	0.56497	0.82511
0.50628	0.86237	0.52126	0.85340	0.53607	0.84417	0.55072	0.83469	0.56521	0.82495
0.50654	0.86222	0.52151	0.85325	0.53632	0.84402	0.55097	0.83453	0.56545	0.82478
0.50679	0.86207	0.52175	0.85310	0.53656	0.84386	0.55121	0.83437	0.56569	0.82462
0.50704	0.86192	0.52200	0.85294	0.53681	0.84370	0.55145	0.83421	0.56593	0.82446
0.50729	0.86178	0.52225	0.85279	0.53705	0.84355	0.55169	0.83405	0.56617	0.82429
0.50754	0.86163	0.52250	0.85264	0.53730	0.84339	0.55194	0.83389	0.56641	0.82413
0.50779	0.86148	0.52275	0.85249	0.53754	0.84324	0.55218	0.83373	0.56665	0.82396
0.50804	0.86133	0.52299	0.85234	0.53779	0.84308	0.55242	0.83356	0.56689	0.82380
0.50829	0.86119	0.52324	0.85218	0.53804	0.84292	0.55266	0.83340	0.56713	0.82363
0.50854	0.86104	0.52349	0.85203	0.53828	0.84277	0.55291	0.83324	0.56736	0.82347
0.50879	0.86089	0.52374	0.85188	0.53853	0.84261	0.55315	0.83308	0.56760	0.82330
0.50904	0.86074	0.52399	0.85173	0.53877	0.84245	0.55339	0.83292	0.56784	0.82314
0.50929	0.86059	0.52423	0.85157	0.53902	0.84230	0.55363	0.83276	0.56808	0.82297
0.50954	0.86045	0.52448	0.85142	0.53926	0.84214	0.55388	0.83260	0.56832	0.82281
0.50979	0.86030	0.52473	0.85127	0.53951	0.84198	0.55412	0.83244	0.56856	0.82264
0.51004	0.86015	0.52498	0.85112	0.53975	0.84182	0.55436	0.83228	0.56880	0.82248
0.51029	0.86000	0.52522	0.85096	0.54000	0.84167	0.55460	0.83212	0.56904	0.82231
0.51054	0.85985	0.52547	0.85081	0.54024	0.84151	0.55484	0.83195	0.56928	0.82214
0.51079	0.85970	0.52572	0.85066	0.54049	0.84135	0.55509	0.83179	0.56952	0.82198
0.51104	0.85956	0.52597	0.85051	0.54073	0.84120	0.55533	0.83163	0.56976	0.82181
0.51129	0.85941	0.52621	0.85035	0.54097	0.84104	0.55557	0.83147	0.57000	0.82165
0.51154	0.85926	0.52646	0.85020	0.54122	0.84088	0.55581	0.83131	0.57024	0.82148
0.51179	0.85911	0.52671	0.85005	0.54146	0.84072	0.55605	0.83115	0.57047	0.82132
0.51204	0.85896	0.52696	0.84989	0.54171	0.84057	0.55630	0.83098	0.57071	0.82115
0.51229	0.85881	0.52720	0.84974	0.54195	0.84041	0.55654	0.83082	0.57095	0.82098
0.51254	0.85866	0.52745	0.84959	0.54220	0.84025	0.55678	0.83066	0.57119	0.82082
0.51279	0.85851	0.52770	0.84943	0.54244	0.84009	0.55702	0.83050	0.57143	0.82065
0.51304	0.85836	0.52794	0.84928	0.54269	0.83994	0.55726	0.83034	0.57167	0.82048
0.51329	0.85821	0.52819	0.84913	0.54293	0.83978	0.55750	0.83017	0.57191	0.82032
0.51354	0.85806	0.52844	0.84897	0.54317	0.83962	0.55775	0.83001	0.57215	0.82015
0.51379	0.85792	0.52869	0.84882	0.54342	0.83946	0.55799	0.82985	0.57238	0.81999
0.51404	0.85777	0.52893	0.84866	0.54366	0.83930	0.55823	0.82969	0.57262	0.81982
0.51429	0.85762	0.52918	0.84851	0.54391	0.83915	0.55847	0.82953	0.57286	0.81965
0.51454	0.85747	0.52943	0.84836	0.54415	0.83899	0.55871	0.82936	0.57310	0.81949
0.51479	0.85732	0.52967	0.84820	0.54440	0.83883	0.55895	0.82920	0.57334	0.81932
0.51504	0.85717	0.52992	0.84805	0.54464	0.83867	0.55919	0.82904	0.57358	0.81915

COS	SIN	COS	SIN	COS	SIN	COS	SIN	COS	SIN
59°		58°		57°		56°		55°	

M I N	35°		36°		37°		38°		39
	SIN	COS	SIN	COS	SIN	COS	SIN	COS	SIN
0	0.57358	0.81915	0.58779	0.80902	0.60182	0.79864	0.61566	0.78801	0.62932
1	0.57381	0.81899	0.58802	0.80885	0.60205	0.79846	0.61589	0.78783	0.62955
2	0.57405	0.81882	0.58826	0.80867	0.60228	0.79829	0.61612	0.78765	0.62977
3	0.57429	0.81865	0.58849	0.80850	0.60251	0.79811	0.61635	0.78747	0.63000
4	0.57453	0.81848	0.58873	0.80833	0.60274	0.79793	0.61658	0.78729	0.63022
5	0.57477	0.81832	0.58896	0.80816	0.60298	0.79776	0.61681	0.78711	0.63045
6	0.57501	0.81815	0.58920	0.80799	0.60321	0.79758	0.61704	0.78694	0.63068
7	0.57524	0.81798	0.58943	0.80782	0.60344	0.79741	0.61726	0.78676	0.63090
8	0.57548	0.81782	0.58967	0.80765	0.60367	0.79723	0.61749	0.78658	0.63113
9	0.57572	0.81765	0.58990	0.80748	0.60390	0.79706	0.61772	0.78640	0.63135
10	0.57596	0.81748	0.59014	0.80730	0.60414	0.79688	0.61795	0.78622	0.63158
11	0.57619	0.81731	0.59037	0.80713	0.60437	0.79671	0.61818	0.78604	0.63180
12	0.57643	0.81714	0.59061	0.80696	0.60460	0.79653	0.61841	0.78586	0.63203
13	0.57667	0.81698	0.59084	0.80679	0.60483	0.79635	0.61864	0.78568	0.63225
14	0.57691	0.81681	0.59108	0.80662	0.60506	0.79618	0.61887	0.78550	0.63248
15	0.57715	0.81664	0.59131	0.80644	0.60529	0.79600	0.61909	0.78532	0.63271
16	0.57738	0.81647	0.59154	0.80627	0.60553	0.79583	0.61932	0.78514	0.63293
17	0.57762	0.81631	0.59178	0.80610	0.60576	0.79565	0.61955	0.78496	0.63316
18	0.57786	0.81614	0.59201	0.80593	0.60599	0.79547	0.61978	0.78478	0.63338
19	0.57810	0.81597	0.59225	0.80576	0.60622	0.79530	0.62001	0.78460	0.63361
20	0.57833	0.81580	0.59248	0.80558	0.60645	0.79512	0.62024	0.78442	0.63383
21	0.57857	0.81563	0.59272	0.80541	0.60668	0.79494	0.62046	0.78424	0.63406
22	0.57881	0.81546	0.59295	0.80524	0.60691	0.79477	0.62069	0.78405	0.63428
23	0.57904	0.81530	0.59318	0.80507	0.60714	0.79459	0.62092	0.78387	0.63451
24	0.57928	0.81513	0.59342	0.80489	0.60738	0.79441	0.62115	0.78369	0.63473
25	0.57952	0.81496	0.59365	0.80472	0.60761	0.79424	0.62138	0.78351	0.63496
26	0.57976	0.81479	0.59389	0.80455	0.60784	0.79406	0.62160	0.78333	0.63518
27	0.57999	0.81462	0.59412	0.80438	0.60807	0.79388	0.62183	0.78315	0.63540
28	0.58023	0.81445	0.59436	0.80420	0.60830	0.79371	0.62206	0.78297	0.63563
29	0.58047	0.81428	0.59459	0.80403	0.60853	0.79353	0.62229	0.78279	0.63585
30	0.58070	0.81412	0.59482	0.80386	0.60876	0.79335	0.62251	0.78261	0.63608
31	0.58094	0.81395	0.59506	0.80368	0.60899	0.79318	0.62274	0.78243	0.63630
32	0.58118	0.81378	0.59529	0.80351	0.60922	0.79300	0.62297	0.78225	0.63653
33	0.58141	0.81361	0.59552	0.80334	0.60945	0.79282	0.62320	0.78206	0.63675
34	0.58165	0.81344	0.59576	0.80316	0.60968	0.79264	0.62342	0.78188	0.63698
35	0.58189	0.81327	0.59599	0.80299	0.60991	0.79247	0.62365	0.78170	0.63720
36	0.58212	0.81310	0.59622	0.80282	0.61015	0.79229	0.62388	0.78152	0.63742
37	0.58236	0.81293	0.59646	0.80264	0.61038	0.79211	0.62411	0.78134	0.63765
38	0.58260	0.81276	0.59669	0.80247	0.61061	0.79193	0.62433	0.78116	0.63787
39	0.58283	0.81259	0.59693	0.80230	0.61084	0.79176	0.62456	0.78098	0.63810
40	0.58307	0.81242	0.59716	0.80212	0.61107	0.79158	0.62479	0.78079	0.63832
41	0.58330	0.81225	0.59739	0.80195	0.61130	0.79140	0.62502	0.78061	0.63854
42	0.58354	0.81208	0.59763	0.80178	0.61153	0.79122	0.62524	0.78043	0.63877
43	0.58378	0.81191	0.59786	0.80160	0.61176	0.79105	0.62547	0.78025	0.63899
44	0.58401	0.81174	0.59809	0.80143	0.61199	0.79087	0.62570	0.78007	0.63922
45	0.58425	0.81157	0.59832	0.80125	0.61222	0.79069	0.62592	0.77988	0.63944
46	0.58449	0.81140	0.59856	0.80108	0.61245	0.79051	0.62615	0.77970	0.63966
47	0.58472	0.81123	0.59879	0.80091	0.61268	0.79033	0.62638	0.77952	0.63989
48	0.58496	0.81106	0.59902	0.80073	0.61291	0.79016	0.62660	0.77934	0.64011
49	0.58519	0.81089	0.59926	0.80056	0.61314	0.78998	0.62683	0.77916	0.64033
50	0.58543	0.81072	0.59949	0.80038	0.61337	0.78980	0.62706	0.77897	0.64056
51	0.58567	0.81055	0.59972	0.80021	0.61360	0.78962	0.62728	0.77879	0.64078
52	0.58590	0.81038	0.59995	0.80003	0.61383	0.78944	0.62751	0.77861	0.64100
53	0.58614	0.81021	0.60019	0.79986	0.61406	0.78926	0.62774	0.77843	0.64123
54	0.58637	0.81004	0.60042	0.79968	0.61429	0.78908	0.62796	0.77824	0.64145
55	0.58661	0.80987	0.60065	0.79951	0.61451	0.78891	0.62819	0.77806	0.64167
56	0.58684	0.80970	0.60089	0.79934	0.61474	0.78873	0.62842	0.77788	0.64190
57	0.58708	0.80953	0.60112	0.79916	0.61497	0.78855	0.62864	0.77769	0.64212
58	0.58731	0.80936	0.60135	0.79899	0.61520	0.78837	0.62887	0.77751	0.64234
59	0.58755	0.80919	0.60158	0.79881	0.61543	0.78819	0.62909	0.77733	0.64256
60	0.58779	0.80902	0.60182	0.79864	0.61566	0.78801	0.62932	0.77715	0.64279
	COS	SIN	COS	SIN	COS	SIN	COS	SIN	COS
	54°		53°		52°		51°		S

40°		41°		42°		43°		44°	
SIN	COS	SIN	COS	SIN	COS	SIN	COS	SIN	COS
0.64279	0.76604	0.65606	0.75471	0.66913	0.74314	0.68200	0.73135	0.69466	0.71934
0.64301	0.76586	0.65628	0.75452	0.66935	0.74295	0.68221	0.73116	0.69487	0.71914
0.64323	0.76567	0.65650	0.75433	0.66956	0.74276	0.68242	0.73096	0.69508	0.71894
0.64346	0.76548	0.65672	0.75414	0.66978	0.74256	0.68264	0.73076	0.69529	0.71873
0.64368	0.76530	0.65694	0.75395	0.66999	0.74237	0.68285	0.73056	0.69549	0.71853
0.64390	0.76511	0.65716	0.75375	0.67021	0.74217	0.68306	0.73036	0.69570	0.71833
0.64412	0.76492	0.65738	0.75356	0.67043	0.74198	0.68327	0.73016	0.69591	0.71813
0.64435	0.76473	0.65759	0.75337	0.67064	0.74178	0.68349	0.72996	0.69612	0.71792
0.64457	0.76455	0.65781	0.75318	0.67086	0.74159	0.68370	0.72976	0.69633	0.71772
0.64479	0.76436	0.65803	0.75299	0.67107	0.74139	0.68391	0.72957	0.69654	0.71752
0.64501	0.76417	0.65825	0.75280	0.67129	0.74120	0.68412	0.72937	0.69675	0.71732
0.64524	0.76398	0.65847	0.75261	0.67151	0.74100	0.68434	0.72917	0.69696	0.71711
0.64546	0.76380	0.65869	0.75241	0.67172	0.74080	0.68455	0.72897	0.69717	0.71691
0.64568	0.76361	0.65891	0.75222	0.67194	0.74061	0.68476	0.72877	0.69737	0.71671
0.64590	0.76342	0.65913	0.75203	0.67215	0.74041	0.68497	0.72857	0.69758	0.71650
0.64612	0.76323	0.65935	0.75184	0.67237	0.74022	0.68518	0.72837	0.69779	0.71630
0.64635	0.76304	0.65956	0.75165	0.67258	0.74002	0.68539	0.72817	0.69800	0.71610
0.64657	0.76286	0.65978	0.75146	0.67280	0.73983	0.68561	0.72797	0.69821	0.71590
0.64679	0.76267	0.66000	0.75126	0.67301	0.73963	0.68582	0.72777	0.69842	0.71569
0.64701	0.76248	0.66022	0.75107	0.67323	0.73944	0.68603	0.72757	0.69862	0.71549
0.64723	0.76229	0.66044	0.75088	0.67344	0.73924	0.68624	0.72737	0.69883	0.71529
0.64746	0.76210	0.66066	0.75069	0.67366	0.73904	0.68645	0.72717	0.69904	0.71508
0.64768	0.76192	0.66088	0.75050	0.67387	0.73885	0.68666	0.72697	0.69925	0.71488
0.64790	0.76173	0.66109	0.75030	0.67409	0.73865	0.68688	0.72677	0.69946	0.71468
0.64812	0.76154	0.66131	0.75011	0.67430	0.73846	0.68709	0.72657	0.69966	0.71447
0.64834	0.76135	0.66153	0.74992	0.67452	0.73826	0.68730	0.72637	0.69987	0.71427
0.64856	0.76116	0.66175	0.74973	0.67473	0.73806	0.68751	0.72617	0.70008	0.71407
0.64878	0.76097	0.66197	0.74953	0.67495	0.73787	0.68772	0.72597	0.70029	0.71386
0.64901	0.76078	0.66218	0.74934	0.67516	0.73767	0.68793	0.72577	0.70049	0.71366
0.64923	0.76059	0.66240	0.74915	0.67538	0.73747	0.68814	0.72557	0.70070	0.71345
0.64945	0.76041	0.66262	0.74896	0.67559	0.73728	0.68835	0.72537	0.70091	0.71325
0.64967	0.76022	0.66284	0.74876	0.67580	0.73708	0.68857	0.72517	0.70112	0.71305
0.64989	0.76003	0.66306	0.74857	0.67602	0.73688	0.68878	0.72497	0.70132	0.71284
0.65011	0.75984	0.66327	0.74838	0.67623	0.73669	0.68899	0.72477	0.70153	0.71264
0.65033	0.75965	0.66349	0.74818	0.67645	0.73649	0.68920	0.72457	0.70174	0.71243
0.65055	0.75946	0.66371	0.74799	0.67666	0.73629	0.68941	0.72437	0.70195	0.71223
0.65077	0.75927	0.66393	0.74780	0.67688	0.73610	0.68962	0.72417	0.70215	0.71203
0.65100	0.75908	0.66414	0.74760	0.67709	0.73590	0.68983	0.72397	0.70236	0.71182
0.65122	0.75889	0.66436	0.74741	0.67730	0.73570	0.69004	0.72377	0.70257	0.71162
0.65144	0.75870	0.66458	0.74722	0.67752	0.73551	0.69025	0.72357	0.70277	0.71141
0.65166	0.75851	0.66480	0.74703	0.67773	0.73531	0.69046	0.72337	0.70298	0.71121
0.65188	0.75832	0.66501	0.74683	0.67795	0.73511	0.69067	0.72317	0.70319	0.71100
0.65210	0.75813	0.66523	0.74664	0.67816	0.73491	0.69088	0.72297	0.70339	0.71080
0.65232	0.75794	0.66545	0.74644	0.67837	0.73472	0.69109	0.72277	0.70360	0.71059
0.65254	0.75775	0.66566	0.74625	0.67859	0.73452	0.69130	0.72257	0.70381	0.71039
0.65276	0.75756	0.66588	0.74606	0.67880	0.73432	0.69151	0.72236	0.70401	0.71019
0.65298	0.75738	0.66610	0.74586	0.67901	0.73413	0.69172	0.72216	0.70422	0.70998
0.65320	0.75719	0.66632	0.74567	0.67923	0.73393	0.69193	0.72196	0.70443	0.70978
0.65342	0.75700	0.66653	0.74548	0.67944	0.73373	0.69214	0.72176	0.70463	0.70957
0.65364	0.75680	0.66675	0.74528	0.67965	0.73353	0.69235	0.72156	0.70484	0.70937
0.65386	0.75661	0.66697	0.74509	0.67987	0.73333	0.69256	0.72136	0.70505	0.70916
0.65408	0.75642	0.66718	0.74489	0.68008	0.73314	0.69277	0.72116	0.70525	0.70896
0.65430	0.75623	0.66740	0.74470	0.68029	0.73294	0.69298	0.72095	0.70546	0.70875
0.65452	0.75604	0.66762	0.74451	0.68051	0.73274	0.69319	0.72075	0.70567	0.70855
0.65474	0.75585	0.66783	0.74431	0.68072	0.73254	0.69340	0.72055	0.70587	0.70834
0.65496	0.75566	0.66805	0.74412	0.68093	0.73234	0.69361	0.72035	0.70608	0.70813
0.65518	0.75547	0.66827	0.74392	0.68115	0.73215	0.69382	0.72015	0.70628	0.70793
0.65540	0.75528	0.66848	0.74373	0.68136	0.73195	0.69403	0.71995	0.70649	0.70772
0.65562	0.75509	0.66870	0.74353	0.68157	0.73175	0.69424	0.71974	0.70670	0.70752
0.65584	0.75490	0.66891	0.74334	0.68179	0.73155	0.69445	0.71954	0.70690	0.70731
0.65606	0.75471	0.66913	0.74314	0.68200	0.73135	0.69466	0.71934	0.70711	0.70711
COS	SIN	COS	SIN	COS	SIN	COS	SIN	COS	SIN
49°		48°		47°		46°		45°	

APPENDIX III

NATURAL TANGENTS AND COTANGENTS

0°		1°		2°		3°		4°	
TAN	COT	TAN	COT	TAN	COT	TAN	COT	TAN	COT
0000000	0000000	0.01746	57.2900	0.03492	28.6363	0.05241	19.0811	0.06993	14.3007
0.00029	3437.75	0.01775	56.3506	0.03521	28.3994	0.05270	18.9755	0.07022	14.2411
0.00058	1718.87	0.01804	55.4415	0.03550	28.1664	0.05299	18.8711	0.07051	14.1821
0.00087	1145.92	0.01833	54.5613	0.03579	27.9372	0.05328	18.7678	0.07080	14.1235
0.00116	859.436	0.01862	53.7086	0.03609	27.7117	0.05357	18.6656	0.07110	14.0655
0.00145	687.549	0.01891	52.8821	0.03638	27.4899	0.05387	18.5645	0.07139	14.0079
0.00175	572.957	0.01920	52.0807	0.03667	27.2715	0.05416	18.4645	0.07168	13.9507
0.00204	491.106	0.01949	51.3032	0.03696	27.0566	0.05445	18.3655	0.07197	13.8940
0.00233	429.718	0.01978	50.5485	0.03725	26.8450	0.05474	18.2677	0.07227	13.8378
0.00262	381.971	0.02007	49.8157	0.03754	26.6367	0.05503	18.1708	0.07256	13.7821
0.00291	343.774	0.02036	49.1039	0.03783	26.4316	0.05533	18.0750	0.07285	13.7267
0.00320	312.521	0.02066	48.4121	0.03812	26.2296	0.05562	17.9802	0.07314	13.6719
0.00349	286.478	0.02095	47.7395	0.03842	26.0307	0.05591	17.8863	0.07344	13.6174
0.00378	264.441	0.02124	47.0853	0.03871	25.8348	0.05620	17.7934	0.07373	13.5634
0.00407	245.552	0.02153	46.4489	0.03900	25.6418	0.05649	17.7015	0.07402	13.5098
0.00436	229.182	0.02182	45.8294	0.03929	25.4517	0.05678	17.6106	0.07431	13.4566
0.00465	214.858	0.02211	45.2261	0.03958	25.2644	0.05708	17.5205	0.07461	13.4039
0.00495	202.219	0.02240	44.6386	0.03987	25.0798	0.05737	17.4314	0.07490	13.3515
0.00524	190.984	0.02269	44.0661	0.04016	24.8978	0.05766	17.3432	0.07519	13.2996
0.00553	180.932	0.02298	43.5081	0.04046	24.7185	0.05795	17.2558	0.07548	13.2480
0.00582	171.885	0.02328	42.9641	0.04075	24.5418	0.05824	17.1693	0.07578	13.1969
0.00611	163.700	0.02357	42.4335	0.04104	24.3675	0.05854	17.0837	0.07607	13.1461
0.00640	156.259	0.02386	41.9158	0.04133	24.1957	0.05883	16.9990	0.07636	13.0958
0.00669	149.465	0.02415	41.4106	0.04162	24.0263	0.05912	16.9150	0.07665	13.0458
0.00698	143.237	0.02444	40.9174	0.04191	23.8593	0.05941	16.8319	0.07695	12.9962
0.00727	137.507	0.02473	40.4358	0.04220	23.6945	0.05970	16.7496	0.07724	12.9469
0.00756	132.219	0.02502	39.9655	0.04250	23.5321	0.05999	16.6681	0.07753	12.8981
0.00785	127.321	0.02531	39.5059	0.04279	23.3718	0.06029	16.5874	0.07782	12.8496
0.00815	122.774	0.02560	39.0568	0.04308	23.2137	0.06058	16.5075	0.07812	12.8014
0.00844	118.540	0.02589	38.6177	0.04337	23.0577	0.06087	16.4283	0.07841	12.7536
0.00873	114.589	0.02619	38.1885	0.04366	22.9038	0.06116	16.3499	0.07870	12.7062
0.00902	110.892	0.02648	37.7686	0.04395	22.7519	0.06145	16.2722	0.07899	12.6591
0.00931	107.426	0.02677	37.3579	0.04424	22.6020	0.06175	16.1952	0.07929	12.6124
0.00960	104.171	0.02706	36.9560	0.04454	22.4541	0.06204	16.1190	0.07958	12.5660
0.00989	101.107	0.02735	36.5627	0.04483	22.3081	0.06233	16.0435	0.07987	12.5199
0.01018	98.2179	0.02764	36.1776	0.04512	22.1640	0.06262	15.9687	0.08017	12.4742
0.01047	95.4895	0.02793	35.8006	0.04541	22.0217	0.06291	15.8945	0.08046	12.4288
0.01076	92.9085	0.02822	35.4313	0.04570	21.8813	0.06321	15.8211	0.08075	12.3838
0.01105	90.4633	0.02851	35.0695	0.04599	21.7426	0.06350	15.7483	0.08104	12.3390
0.01135	88.1436	0.02881	34.7151	0.04628	21.6056	0.06379	15.6762	0.08134	12.2946
0.01164	85.9398	0.02910	34.3678	0.04658	21.4704	0.06408	15.6048	0.08163	12.2505
0.01193	83.8435	0.02939	34.0273	0.04687	21.3369	0.06438	15.5340	0.08192	12.2067
0.01222	81.8470	0.02968	33.6935	0.04716	21.2049	0.06467	15.4638	0.08221	12.1632
0.01251	79.9434	0.02997	33.3662	0.04745	21.0747	0.06496	15.3943	0.08251	12.1201
0.01280	78.1263	0.03026	33.0452	0.04774	20.9460	0.06525	15.3254	0.08280	12.0772
0.01309	76.3900	0.03055	32.7303	0.04803	20.8188	0.06554	15.2571	0.08309	12.0346
0.01338	74.7292	0.03084	32.4213	0.04833	20.6932	0.06584	15.1893	0.08339	11.9923
0.01367	73.1390	0.03114	32.1181	0.04862	20.5691	0.06613	15.1222	0.08368	11.9504
0.01396	71.6151	0.03143	31.8205	0.04891	20.4465	0.06642	15.0557	0.08397	11.9087
0.01425	70.1533	0.03172	31.5284	0.04920	20.3253	0.06671	14.9898	0.08427	11.8673
0.01455	68.7501	0.03201	31.2416	0.04949	20.2056	0.06700	14.9244	0.08456	11.8262
0.01484	67.4019	0.03230	30.9599	0.04978	20.0872	0.06730	14.8596	0.08485	11.7853
0.01513	66.1055	0.03259	30.6833	0.05007	19.9702	0.06759	14.7954	0.08514	11.7448
0.01542	64.8580	0.03288	30.4116	0.05037	19.8546	0.06788	14.7317	0.08544	11.7045
0.01571	63.6567	0.03317	30.1446	0.05066	19.7403	0.06817	14.6685	0.08573	11.6645
0.01600	62.4992	0.03346	29.8823	0.05095	19.6273	0.06847	14.6059	0.08602	11.6248
0.01629	61.3829	0.03376	29.6245	0.05124	19.5156	0.06876	14.5438	0.08632	11.5853
0.01658	60.3058	0.03405	29.3711	0.05153	19.4051	0.06905	14.4823	0.08661	11.5461
0.01687	59.2659	0.03434	29.1220	0.05182	19.2959	0.06934	14.4212	0.08690	11.5072
0.01716	58.2612	0.03463	28.8771	0.05212	19.1879	0.06963	14.3607	0.08720	11.4685
0.01746	57.2900	0.03492	28.6363	0.05241	19.0811	0.06993	14.3007	0.08749	11.4301
COT	TAN	COT	TAN	COT	TAN	COT	TAN	COT	TAN
89°		88°		87°		86°		85°	

MIN	5°		6°		7°		8°		TA
	TAN	COT	TAN	COT	TAN	COT	TAN	COT	
0	0.08749	11.4301	0.10510	9.51436	0.12278	8.14435	0.14054	7.11537	0.158
1	0.08778	11.3919	0.10540	9.48781	0.12308	8.12481	0.14084	7.10038	0.158
2	0.08807	11.3540	0.10569	9.46141	0.12338	8.10536	0.14113	7.08546	0.158
3	0.08837	11.3163	0.10599	9.43515	0.12367	8.08600	0.14143	7.07059	0.159
4	0.08866	11.2789	0.10628	9.40904	0.12397	8.06674	0.14173	7.05579	0.159
5	0.08895	11.2417	0.10657	9.38307	0.12426	8.04756	0.14202	7.04105	0.159
6	0.08925	11.2048	0.10687	9.35724	0.12456	8.02848	0.14232	7.02637	0.160
7	0.08954	11.1681	0.10716	9.33155	0.12485	8.00948	0.14262	7.01174	0.160
8	0.08983	11.1316	0.10746	9.30599	0.12515	7.99058	0.14291	6.99718	0.160
9	0.09013	11.0954	0.10775	9.28058	0.12544	7.97176	0.14321	6.98268	0.161
10	0.09042	11.0594	0.10805	9.25530	0.12574	7.95302	0.14351	6.96823	0.161
11	0.09071	11.0237	0.10834	9.23016	0.12603	7.93438	0.14381	6.95385	0.161
12	0.09101	10.9882	0.10863	9.20516	0.12633	7.91582	0.14410	6.93952	0.161
13	0.09130	10.9529	0.10893	9.18028	0.12662	7.89734	0.14440	6.92525	0.162
14	0.09159	10.9178	0.10922	9.15554	0.12692	7.87895	0.14470	6.91104	0.162
15	0.09189	10.8829	0.10952	9.13093	0.12722	7.86064	0.14499	6.89688	0.162
16	0.09218	10.8483	0.10981	9.10646	0.12751	7.84242	0.14529	6.88278	0.163
17	0.09247	10.8139	0.11011	9.08211	0.12781	7.82428	0.14559	6.86874	0.163
18	0.09277	10.7797	0.11040	9.05789	0.12810	7.80622	0.14588	6.85475	0.163
19	0.09306	10.7457	0.11070	9.03379	0.12840	7.78825	0.14618	6.84082	0.164
20	0.09335	10.7119	0.11099	9.00983	0.12869	7.77035	0.14648	6.82694	0.164
21	0.09365	10.6783	0.11128	8.98598	0.12899	7.75254	0.14678	6.81312	0.164
22	0.09394	10.6450	0.11158	8.96227	0.12929	7.73480	0.14707	6.79936	0.164
23	0.09423	10.6118	0.11187	8.93867	0.12958	7.71715	0.14737	6.78564	0.165
24	0.09453	10.5789	0.11217	8.91520	0.12988	7.69957	0.14767	6.77199	0.165
25	0.09482	10.5462	0.11246	8.89185	0.13017	7.68208	0.14796	6.75838	0.165
26	0.09511	10.5136	0.11276	8.86862	0.13047	7.66466	0.14826	6.74483	0.166
27	0.09541	10.4813	0.11305	8.84551	0.13076	7.64732	0.14856	6.73133	0.166
28	0.09570	10.4491	0.11335	8.82252	0.13106	7.63005	0.14886	6.71789	0.166
29	0.09600	10.4172	0.11364	8.79964	0.13136	7.61287	0.14915	6.70450	0.167
30	0.09629	10.3854	0.11394	8.77689	0.13165	7.59575	0.14945	6.69116	0.167
31	0.09658	10.3538	0.11423	8.75425	0.13195	7.57872	0.14975	6.67787	0.167
32	0.09688	10.3224	0.11452	8.73172	0.13224	7.56176	0.15005	6.66463	0.167
33	0.09717	10.2913	0.11482	8.70931	0.13254	7.54487	0.15034	6.65144	0.168
34	0.09746	10.2602	0.11511	8.68701	0.13284	7.52806	0.15064	6.63831	0.168
35	0.09776	10.2294	0.11541	8.66482	0.13313	7.51132	0.15094	6.62523	0.168
36	0.09805	10.1988	0.11570	8.64275	0.13343	7.49465	0.15124	6.61219	0.169
37	0.09834	10.1683	0.11600	8.62078	0.13372	7.47806	0.15153	6.59921	0.169
38	0.09864	10.1381	0.11629	8.59893	0.13402	7.46154	0.15183	6.58627	0.169
39	0.09893	10.1080	0.11659	8.57718	0.13432	7.44509	0.15213	6.57339	0.170
40	0.09923	10.0780	0.11688	8.55555	0.13461	7.42871	0.15243	6.56055	0.170
41	0.09952	10.0483	0.11718	8.53402	0.13491	7.41240	0.15272	6.54777	0.170
42	0.09981	10.0187	0.11747	8.51259	0.13521	7.39616	0.15302	6.53503	0.170
43	0.10011	9.98931	0.11777	8.49128	0.13550	7.37999	0.15332	6.52234	0.171
44	0.10040	9.96007	0.11806	8.47007	0.13580	7.36389	0.15362	6.50970	0.171
45	0.10069	9.93101	0.11836	8.44896	0.13609	7.34786	0.15391	6.49710	0.171
46	0.10099	9.90211	0.11865	8.42795	0.13639	7.33190	0.15421	6.48456	0.172
47	0.10128	9.87338	0.11895	8.40705	0.13669	7.31600	0.15451	6.47206	0.172
48	0.10158	9.84482	0.11924	8.38625	0.13698	7.30018	0.15481	6.45961	0.172
49	0.10187	9.81641	0.11954	8.36555	0.13728	7.28442	0.15511	6.44720	0.173
50	0.10216	9.78817	0.11983	8.34496	0.13758	7.26873	0.15540	6.43484	0.173
51	0.10246	9.76009	0.12013	8.32446	0.13787	7.25310	0.15570	6.42253	0.173
52	0.10275	9.73217	0.12042	8.30406	0.13817	7.23754	0.15600	6.41026	0.173
53	0.10305	9.70441	0.12072	8.28376	0.13846	7.22204	0.15630	6.39804	0.174
54	0.10334	9.67680	0.12101	8.26355	0.13876	7.20661	0.15660	6.38587	0.174
55	0.10363	9.64935	0.12131	8.24345	0.13906	7.19125	0.15689	6.37374	0.174
56	0.10393	9.62205	0.12160	8.22344	0.13935	7.17594	0.15719	6.36165	0.175
57	0.10422	9.59490	0.12190	8.20352	0.13965	7.16071	0.15749	6.34961	0.175
58	0.10452	9.56791	0.12219	8.18370	0.13995	7.14553	0.15779	6.33761	0.175
59	0.10481	9.54106	0.12249	8.16398	0.14024	7.13042	0.15809	6.32566	0.176
60	0.10510	9.51436	0.12278	8.14435	0.14054	7.11537	0.15838	6.31375	0.176
	COT	TAN	COT	TAN	COT	TAN	COT	TAN	CO
	84°		83°		82°		81°		

10°		11°		12°		13°		14°	
TAN	COT	TAN	COT	TAN	COT	TAN	COT	TAN	COT
0.17633	5.67128	0.19438	5.14455	0.21256	4.70463	0.23087	4.33148	0.24933	4.01078
0.17663	5.66165	0.19468	5.13658	0.21286	4.69791	0.23117	4.32573	0.24964	4.00582
0.17693	5.65205	0.19498	5.12862	0.21316	4.69121	0.23148	4.32001	0.24995	4.00086
0.17723	5.64248	0.19529	5.12069	0.21347	4.68452	0.23179	4.31430	0.25026	3.99592
0.17753	5.63295	0.19559	5.11279	0.21377	4.67786	0.23209	4.30860	0.25056	3.99099
0.17783	5.62344	0.19589	5.10490	0.21408	4.67121	0.23240	4.30291	0.25087	3.98607
0.17813	5.61397	0.19619	5.09704	0.21438	4.66458	0.23271	4.29724	0.25118	3.98117
0.17843	5.60452	0.19649	5.08921	0.21469	4.65797	0.23301	4.29159	0.25149	3.97627
0.17873	5.59511	0.19680	5.08139	0.21499	4.65138	0.23332	4.28595	0.25180	3.97139
0.17903	5.58573	0.19710	5.07360	0.21529	4.64480	0.23363	4.28032	0.25211	3.96651
0.17933	5.57638	0.19740	5.06584	0.21560	4.63825	0.23393	4.27471	0.25242	3.96165
0.17963	5.56706	0.19770	5.05809	0.21590	4.63171	0.23424	4.26911	0.25273	3.95680
0.17993	5.55777	0.19801	5.05037	0.21621	4.62518	0.23455	4.26352	0.25304	3.95196
0.18023	5.54851	0.19831	5.04267	0.21651	4.61868	0.23485	4.25795	0.25335	3.94713
0.18053	5.53927	0.19861	5.03499	0.21682	4.61219	0.23516	4.25239	0.25366	3.94232
0.18083	5.53007	0.19891	5.02734	0.21712	4.60572	0.23547	4.24685	0.25397	3.93751
0.18113	5.52090	0.19921	5.01971	0.21743	4.59927	0.23578	4.24132	0.25428	3.93271
0.18143	5.51176	0.19952	5.01210	0.21773	4.59283	0.23608	4.23580	0.25459	3.92793
0.18173	5.50264	0.19982	5.00451	0.21804	4.58641	0.23639	4.23030	0.25490	3.92316
0.18203	5.49356	0.20012	4.99695	0.21834	4.58001	0.23670	4.22481	0.25521	3.91839
0.18233	5.48451	0.20042	4.98940	0.21864	4.57363	0.23700	4.21933	0.25552	3.91364
0.18263	5.47548	0.20073	4.98188	0.21895	4.56726	0.23731	4.21387	0.25583	3.90890
0.18293	5.46648	0.20103	4.97438	0.21925	4.56091	0.23762	4.20842	0.25614	3.90417
0.18323	5.45751	0.20133	4.96690	0.21956	4.55458	0.23793	4.20298	0.25645	3.89945
0.18353	5.44857	0.20164	4.95945	0.21986	4.54826	0.23823	4.19756	0.25676	3.89474
0.18384	5.43966	0.20194	4.95201	0.22017	4.54196	0.23854	4.19215	0.25707	3.89004
0.18414	5.43077	0.20224	4.94460	0.22047	4.53568	0.23885	4.18675	0.25738	3.88536
0.18444	5.42192	0.20254	4.93721	0.22078	4.52941	0.23916	4.18137	0.25769	3.88068
0.18474	5.41309	0.20285	4.92984	0.22108	4.52316	0.23946	4.17600	0.25800	3.87601
0.18504	5.40429	0.20315	4.92249	0.22139	4.51693	0.23977	4.17064	0.25831	3.87136
0.18534	5.39552	0.20345	4.91516	0.22169	4.51071	0.24008	4.16530	0.25862	3.86671
0.18564	5.38677	0.20376	4.90785	0.22200	4.50451	0.24039	4.15997	0.25893	3.86208
0.18594	5.37805	0.20406	4.90056	0.22231	4.49832	0.24069	4.15465	0.25924	3.85745
0.18624	5.36936	0.20436	4.89330	0.22261	4.49215	0.24100	4.14934	0.25955	3.85284
0.18654	5.36070	0.20466	4.88605	0.22292	4.48600	0.24131	4.14405	0.25986	3.84824
0.18684	5.35206	0.20497	4.87882	0.22322	4.47986	0.24162	4.13877	0.26017	3.84364
0.18714	5.34345	0.20527	4.87162	0.22353	4.47374	0.24193	4.13350	0.26048	3.83906
0.18745	5.33487	0.20557	4.86444	0.22383	4.46764	0.24223	4.12825	0.26079	3.83449
0.18775	5.32631	0.20588	4.85727	0.22414	4.46155	0.24254	4.12301	0.26110	3.82992
0.18805	5.31778	0.20618	4.85013	0.22444	4.45548	0.24285	4.11778	0.26141	3.82537
0.18835	5.30928	0.20648	4.84300	0.22475	4.44942	0.24316	4.11256	0.26172	3.82083
0.18865	5.30080	0.20679	4.83590	0.22505	4.44338	0.24347	4.10736	0.26203	3.81630
0.18895	5.29235	0.20709	4.82882	0.22536	4.43735	0.24377	4.10216	0.26235	3.81177
0.18925	5.28393	0.20739	4.82175	0.22567	4.43134	0.24408	4.09699	0.26266	3.80726
0.18955	5.27553	0.20770	4.81471	0.22597	4.42534	0.24439	4.09182	0.26297	3.80276
0.18986	5.26715	0.20800	4.80769	0.22628	4.41936	0.24470	4.08666	0.26328	3.79827
0.19016	5.25880	0.20830	4.80068	0.22658	4.41340	0.24501	4.08152	0.26359	3.79378
0.19046	5.25048	0.20861	4.79370	0.22689	4.40745	0.24532	4.07639	0.26390	3.78931
0.19076	5.24218	0.20891	4.78673	0.22719	4.40152	0.24562	4.07127	0.26421	3.78485
0.19106	5.23391	0.20921	4.77978	0.22750	4.39560	0.24593	4.06616	0.26452	3.78040
0.19136	5.22566	0.20952	4.77286	0.22781	4.38969	0.24624	4.06107	0.26483	3.77595
0.19166	5.21744	0.20982	4.76595	0.22811	4.38381	0.24655	4.05599	0.26515	3.77152
0.19197	5.20925	0.21013	4.75906	0.22842	4.37793	0.24686	4.05092	0.26546	3.76709
0.19227	5.20107	0.21043	4.75219	0.22872	4.37207	0.24717	4.04586	0.26577	3.76268
0.19257	5.19293	0.21073	4.74534	0.22903	4.36623	0.24747	4.04081	0.26608	3.75828
0.19287	5.18480	0.21104	4.73851	0.22934	4.36040	0.24778	4.03578	0.26639	3.75388
0.19317	5.17671	0.21134	4.73170	0.22964	4.35459	0.24809	4.03076	0.26670	3.74950
0.19347	5.16863	0.21164	4.72490	0.22995	4.34879	0.24840	4.02574	0.26701	3.74512
0.19378	5.16058	0.21195	4.71813	0.23026	4.34300	0.24871	4.02074	0.26733	3.74075
0.19408	5.15256	0.21225	4.71137	0.23056	4.33723	0.24902	4.01576	0.26764	3.73640
0.19438	5.14455	0.21256	4.70463	0.23087	4.33148	0.24933	4.01078	0.26795	3.73205
COT	TAN	COT	TAN	COT	TAN	COT	TAN	COT	TAN
79°		78°		77°		76°		75°	

MIN	15°		16°		17°		18°		
	TAN	COT	TAN	COT	TAN	COT	TAN	COT	TAN
0	0.26795	3.73205	0.28675	3.48741	0.30573	3.27085	0.32492	3.07768	0.3443
1	0.26826	3.72771	0.28706	3.48359	0.30605	3.26745	0.32524	3.07464	0.3446
2	0.26857	3.72338	0.28738	3.47977	0.30637	3.26406	0.32556	3.07160	0.3449
3	0.26888	3.71907	0.28769	3.47596	0.30669	3.26067	0.32588	3.06857	0.3453
4	0.26920	3.71476	0.28801	3.47216	0.30700	3.25729	0.32621	3.06554	0.3456
5	0.26951	3.71046	0.28832	3.46837	0.30732	3.25392	0.32653	3.06252	0.3459
6	0.26982	3.70616	0.28864	3.46458	0.30764	3.25055	0.32685	3.05950	0.3462
7	0.27013	3.70188	0.28895	3.46080	0.30796	3.24719	0.32717	3.05649	0.3466
8	0.27044	3.69761	0.28927	3.45703	0.30828	3.24383	0.32749	3.05349	0.3469
9	0.27076	3.69335	0.28958	3.45327	0.30860	3.24049	0.32782	3.05049	0.3472
10	0.27107	3.68909	0.28990	3.44951	0.30891	3.23714	0.32814	3.04749	0.3475
11	0.27138	3.68485	0.29021	3.44576	0.30923	3.23381	0.32846	3.04450	0.3479
12	0.27169	3.68061	0.29053	3.44202	0.30955	3.23048	0.32878	3.04152	0.3482
13	0.27201	3.67638	0.29084	3.43829	0.30987	3.22715	0.32911	3.03854	0.3485
14	0.27232	3.67217	0.29116	3.43456	0.31019	3.22384	0.32943	3.03556	0.3488
15	0.27263	3.66796	0.29147	3.43084	0.31051	3.22053	0.32975	3.03260	0.3492
16	0.27294	3.66376	0.29179	3.42713	0.31083	3.21722	0.33007	3.02963	0.3495
17	0.27326	3.65957	0.29210	3.42343	0.31115	3.21392	0.33040	3.02667	0.3498
18	0.27357	3.65538	0.29242	3.41973	0.31147	3.21063	0.33072	3.02372	0.3502
19	0.27388	3.65121	0.29274	3.41604	0.31178	3.20734	0.33104	3.02077	0.3505
20	0.27419	3.64705	0.29305	3.41236	0.31210	3.20406	0.33136	3.01783	0.3508
21	0.27451	3.64289	0.29337	3.40869	0.31242	3.20079	0.33169	3.01489	0.3511
22	0.27482	3.63874	0.29368	3.40502	0.31274	3.19752	0.33201	3.01196	0.3515
23	0.27513	3.63461	0.29400	3.40136	0.31306	3.19426	0.33233	3.00903	0.3518
24	0.27545	3.63048	0.29432	3.39771	0.31338	3.19100	0.33266	3.00611	0.3521
25	0.27576	3.62636	0.29463	3.39406	0.31370	3.18775	0.33298	3.00319	0.3524
26	0.27607	3.62224	0.29495	3.39042	0.31402	3.18451	0.33330	3.00028	0.3528
27	0.27638	3.61814	0.29526	3.38679	0.31434	3.18127	0.33363	2.99738	0.3531
28	0.27670	3.61405	0.29558	3.38317	0.31466	3.17804	0.33395	2.99447	0.3534
29	0.27701	3.60996	0.29590	3.37955	0.31498	3.17481	0.33427	2.99158	0.3537
30	0.27732	3.60588	0.29621	3.37594	0.31530	3.17159	0.33460	2.98868	0.3541
31	0.27764	3.60181	0.29653	3.37234	0.31562	3.16838	0.33492	2.98580	0.3544
32	0.27795	3.59775	0.29685	3.36875	0.31594	3.16517	0.33524	2.98292	0.3547
33	0.27826	3.59370	0.29716	3.36516	0.31626	3.16197	0.33557	2.98004	0.3551
34	0.27858	3.58966	0.29748	3.36158	0.31658	3.15877	0.33589	2.97717	0.3554
35	0.27889	3.58562	0.29780	3.35800	0.31690	3.15558	0.33621	2.97430	0.3557
36	0.27921	3.58160	0.29811	3.35443	0.31722	3.15240	0.33654	2.97144	0.3560
37	0.27952	3.57758	0.29843	3.35087	0.31754	3.14922	0.33686	2.96858	0.3564
38	0.27983	3.57357	0.29875	3.34732	0.31786	3.14605	0.33718	2.96573	0.3567
39	0.28015	3.56957	0.29906	3.34377	0.31818	3.14288	0.33751	2.96288	0.3570
40	0.28046	3.56557	0.29938	3.34023	0.31850	3.13972	0.33783	2.96004	0.3574
41	0.28077	3.56159	0.29970	3.33670	0.31882	3.13656	0.33816	2.95721	0.3577
42	0.28109	3.55761	0.30001	3.33317	0.31914	3.13341	0.33848	2.95437	0.3580
43	0.28140	3.55364	0.30033	3.32965	0.31946	3.13027	0.33881	2.95155	0.3583
44	0.28172	3.54968	0.30065	3.32614	0.31978	3.12713	0.33913	2.94872	0.3587
45	0.28203	3.54573	0.30097	3.32264	0.32010	3.12400	0.33945	2.94591	0.3590
46	0.28234	3.54179	0.30128	3.31914	0.32042	3.12087	0.33978	2.94309	0.3593
47	0.28266	3.53785	0.30160	3.31565	0.32074	3.11775	0.34010	2.94028	0.3596
48	0.28297	3.53393	0.30192	3.31216	0.32106	3.11464	0.34043	2.93748	0.3600
49	0.28329	3.53001	0.30224	3.30868	0.32139	3.11153	0.34075	2.93468	0.3603
50	0.28360	3.52609	0.30255	3.30521	0.32171	3.10842	0.34108	2.93189	0.3606
51	0.28391	3.52219	0.30287	3.30174	0.32203	3.10532	0.34140	2.92910	0.3610
52	0.28423	3.51829	0.30319	3.29829	0.32235	3.10223	0.34173	2.92632	0.3613
53	0.28454	3.51441	0.30351	3.29483	0.32267	3.09914	0.34205	2.92354	0.3616
54	0.28486	3.51053	0.30382	3.29139	0.32299	3.09606	0.34238	2.92076	0.3619
55	0.28517	3.50666	0.30414	3.28795	0.32331	3.09298	0.34270	2.91799	0.3623
56	0.28549	3.50279	0.30446	3.28452	0.32363	3.08991	0.34303	2.91523	0.3626
57	0.28580	3.49894	0.30478	3.28109	0.32396	3.08685	0.34335	2.91246	0.3629
58	0.28612	3.49509	0.30509	3.27767	0.32428	3.08379	0.34368	2.90971	0.3633
59	0.28643	3.49125	0.30541	3.27426	0.32460	3.08073	0.34400	2.90696	0.3636
60	0.28675	3.48741	0.30573	3.27085	0.32492	3.07768	0.34433	2.90421	0.3639
	COT	TAN	COT	TAN	COT	TAN	COT	TAN	CO
	74°		73°		72°		71°		

20°		21°		22°		23°		24°	
TAN	COT	TAN	COT	TAN	COT	TAN	COT	TAN	COT
0.36397	2.74748	0.38386	2.60509	0.40403	2.47509	0.42447	2.35585	0.44523	2.24604
0.36430	2.74499	0.38420	2.60283	0.40436	2.47302	0.42482	2.35395	0.44558	2.24428
0.36463	2.74251	0.38453	2.60057	0.40470	2.47095	0.42516	2.35205	0.44593	2.24252
0.36496	2.74004	0.38487	2.59831	0.40504	2.46888	0.42551	2.35015	0.44627	2.24077
0.36529	2.73756	0.38520	2.59606	0.40538	2.46682	0.42585	2.34825	0.44662	2.23902
0.36562	2.73509	0.38553	2.59381	0.40572	2.46476	0.42619	2.34636	0.44697	2.23727
0.36595	2.73263	0.38587	2.59156	0.40606	2.46270	0.42654	2.34447	0.44732	2.23553
0.36628	2.73017	0.38620	2.58932	0.40640	2.46065	0.42688	2.34258	0.44767	2.23378
0.36661	2.72771	0.38654	2.58708	0.40674	2.45860	0.42722	2.34069	0.44802	2.23204
0.36694	2.72526	0.38687	2.58484	0.40707	2.45655	0.42757	2.33881	0.44837	2.23030
0.36727	2.72281	0.38721	2.58261	0.40741	2.45451	0.42791	2.33693	0.44872	2.22857
0.36760	2.72036	0.38754	2.58038	0.40775	2.45246	0.42826	2.33505	0.44907	2.22683
0.36793	2.71792	0.38787	2.57815	0.40809	2.45043	0.42860	2.33317	0.44942	2.22510
0.36826	2.71548	0.38821	2.57593	0.40843	2.44839	0.42894	2.33130	0.44977	2.22337
0.36859	2.71305	0.38854	2.57371	0.40877	2.44636	0.42929	2.32943	0.45012	2.22164
0.36892	2.71062	0.38888	2.57150	0.40911	2.44433	0.42963	2.32756	0.45047	2.21992
0.36925	2.70819	0.38921	2.56928	0.40945	2.44230	0.42998	2.32570	0.45082	2.21819
0.36958	2.70577	0.38955	2.56707	0.40979	2.44027	0.43032	2.32383	0.45117	2.21647
0.36991	2.70335	0.38988	2.56487	0.41013	2.43825	0.43067	2.32197	0.45152	2.21475
0.37024	2.70094	0.39022	2.56266	0.41047	2.43623	0.43101	2.32012	0.45187	2.21304
0.37057	2.69853	0.39055	2.56046	0.41081	2.43422	0.43136	2.31826	0.45222	2.21132
0.37090	2.69612	0.39089	2.55827	0.41115	2.43220	0.43170	2.31641	0.45257	2.20961
0.37123	2.69371	0.39122	2.55608	0.41149	2.43019	0.43205	2.31456	0.45292	2.20790
0.37157	2.69131	0.39156	2.55389	0.41183	2.42819	0.43239	2.31271	0.45327	2.20619
0.37190	2.68892	0.39190	2.55170	0.41217	2.42618	0.43274	2.31086	0.45362	2.20449
0.37223	2.68653	0.39223	2.54952	0.41251	2.42418	0.43308	2.30902	0.45397	2.20278
0.37256	2.68414	0.39257	2.54734	0.41285	2.42218	0.43343	2.30718	0.45432	2.20108
0.37289	2.68175	0.39290	2.54516	0.41319	2.42019	0.43378	2.30534	0.45467	2.19938
0.37322	2.67937	0.39324	2.54299	0.41353	2.41819	0.43412	2.30351	0.45502	2.19769
0.37355	2.67700	0.39357	2.54082	0.41387	2.41620	0.43447	2.30167	0.45538	2.19599
0.37388	2.67462	0.39391	2.53865	0.41421	2.41421	0.43481	2.29984	0.45573	2.19430
0.37422	2.67225	0.39425	2.53648	0.41455	2.41223	0.43516	2.29801	0.45608	2.19261
0.37455	2.66989	0.39458	2.53432	0.41490	2.41025	0.43550	2.29619	0.45643	2.19092
0.37488	2.66752	0.39492	2.53217	0.41524	2.40827	0.43585	2.29437	0.45678	2.18923
0.37521	2.66516	0.39526	2.53001	0.41558	2.40629	0.43620	2.29254	0.45713	2.18755
0.37554	2.66281	0.39559	2.52786	0.41592	2.40432	0.43654	2.29073	0.45748	2.18587
0.37588	2.66046	0.39593	2.52571	0.41626	2.40235	0.43689	2.28891	0.45784	2.18419
0.37621	2.65811	0.39626	2.52357	0.41660	2.40038	0.43724	2.28710	0.45819	2.18251
0.37654	2.65576	0.39660	2.52142	0.41694	2.39841	0.43758	2.28528	0.45854	2.18084
0.37687	2.65342	0.39694	2.51929	0.41728	2.39645	0.43793	2.28348	0.45889	2.17916
0.37720	2.65109	0.39727	2.51715	0.41763	2.39449	0.43828	2.28167	0.45924	2.17749
0.37754	2.64875	0.39761	2.51502	0.41797	2.39253	0.43862	2.27987	0.45960	2.17582
0.37787	2.64642	0.39795	2.51289	0.41831	2.39058	0.43897	2.27806	0.45995	2.17416
0.37820	2.64410	0.39829	2.51076	0.41865	2.38863	0.43932	2.27626	0.46030	2.17249
0.37853	2.64177	0.39862	2.50864	0.41899	2.38668	0.43966	2.27447	0.46065	2.17083
0.37887	2.63945	0.39896	2.50652	0.41933	2.38473	0.44001	2.27267	0.46101	2.16917
0.37920	2.63714	0.39930	2.50440	0.41968	2.38279	0.44036	2.27088	0.46136	2.16751
0.37953	2.63483	0.39963	2.50229	0.42002	2.38084	0.44071	2.26909	0.46171	2.16585
0.37986	2.63252	0.39997	2.50018	0.42036	2.37891	0.44105	2.26730	0.46206	2.16420
0.38020	2.63021	0.40031	2.49807	0.42070	2.37697	0.44140	2.26552	0.46242	2.16255
0.38053	2.62791	0.40065	2.49597	0.42105	2.37504	0.44175	2.26374	0.46277	2.16090
0.38086	2.62561	0.40098	2.49386	0.42139	2.37311	0.44210	2.26196	0.46312	2.15925
0.38120	2.62332	0.40132	2.49177	0.42173	2.37118	0.44244	2.26018	0.46348	2.15760
0.38153	2.62103	0.40166	2.48967	0.42207	2.36925	0.44279	2.25840	0.46383	2.15596
0.38186	2.61874	0.40200	2.48758	0.42242	2.36733	0.44314	2.25663	0.46418	2.15432
0.38220	2.61646	0.40234	2.48549	0.42276	2.36541	0.44349	2.25486	0.46454	2.15268
0.38253	2.61418	0.40267	2.48340	0.42310	2.36349	0.44384	2.25309	0.46489	2.15104
0.38286	2.61190	0.40301	2.48132	0.42345	2.36158	0.44418	2.25132	0.46525	2.14940
0.38320	2.60963	0.40335	2.47924	0.42379	2.35967	0.44453	2.24956	0.46560	2.14777
0.38353	2.60736	0.40369	2.47716	0.42413	2.35776	0.44488	2.24780	0.46595	2.14614
0.38386	2.60509	0.40403	2.47509	0.42447	2.35585	0.44523	2.24604	0.46631	2.14451

COT	TAN	COT	TAN	COT	TAN	COT	TAN	COT	TAN
69°		68°		67°		66°		65°	

MIN	25°		26°		27°		28°		2
	TAN	COT	TAN	COT	TAN	COT	TAN	COT	TAN
0	0.46631	2.14451	0.48773	2.05030	0.50953	1.96261	0.53171	1.88073	0.55431
1	0.46666	2.14288	0.48809	2.04879	0.50989	1.96120	0.53208	1.87941	0.55469
2	0.46702	2.14125	0.48845	2.04728	0.51026	1.95979	0.53246	1.87809	0.55507
3	0.46737	2.13963	0.48881	2.04577	0.51063	1.95838	0.53283	1.87677	0.55545
4	0.46772	2.13801	0.48917	2.04426	0.51099	1.95698	0.53320	1.87546	0.55583
5	0.46808	2.13639	0.48953	2.04276	0.51136	1.95557	0.53358	1.87415	0.55621
6	0.46843	2.13477	0.48989	2.04125	0.51173	1.95417	0.53395	1.87283	0.55659
7	0.46879	2.13316	0.49026	2.03975	0.51209	1.95277	0.53432	1.87152	0.55697
8	0.46914	2.13154	0.49062	2.03825	0.51246	1.95137	0.53470	1.87021	0.55736
9	0.46950	2.12993	0.49098	2.03675	0.51283	1.94997	0.53507	1.86891	0.55774
10	0.46985	2.12832	0.49134	2.03526	0.51319	1.94858	0.53545	1.86760	0.55812
11	0.47021	2.12671	0.49170	2.03376	0.51356	1.94718	0.53582	1.86630	0.55850
12	0.47056	2.12511	0.49206	2.03227	0.51393	1.94579	0.53620	1.86499	0.55888
13	0.47092	2.12350	0.49242	2.03078	0.51430	1.94440	0.53657	1.86369	0.55926
14	0.47128	2.12190	0.49278	2.02929	0.51467	1.94301	0.53694	1.86239	0.55964
15	0.47163	2.12030	0.49315	2.02780	0.51503	1.94162	0.53732	1.86109	0.56003
16	0.47199	2.11871	0.49351	2.02631	0.51540	1.94023	0.53769	1.85979	0.56041
17	0.47234	2.11711	0.49387	2.02483	0.51577	1.93885	0.53807	1.85850	0.56079
18	0.47270	2.11552	0.49423	2.02335	0.51614	1.93746	0.53844	1.85720	0.56117
19	0.47305	2.11392	0.49459	2.02187	0.51651	1.93608	0.53882	1.85591	0.56156
20	0.47341	2.11233	0.49495	2.02039	0.51688	1.93470	0.53920	1.85462	0.56194
21	0.47377	2.11075	0.49532	2.01891	0.51724	1.93332	0.53957	1.85333	0.56232
22	0.47412	2.10916	0.49568	2.01743	0.51761	1.93195	0.53995	1.85204	0.56270
23	0.47448	2.10758	0.49604	2.01596	0.51798	1.93057	0.54032	1.85075	0.56309
24	0.47483	2.10600	0.49640	2.01449	0.51835	1.92920	0.54070	1.84946	0.56347
25	0.47519	2.10442	0.49677	2.01302	0.51872	1.92782	0.54107	1.84818	0.56385
26	0.47555	2.10284	0.49713	2.01155	0.51909	1.92645	0.54145	1.84689	0.56424
27	0.47590	2.10126	0.49749	2.01008	0.51946	1.92508	0.54183	1.84561	0.56462
28	0.47626	2.09969	0.49786	2.00862	0.51983	1.92371	0.54220	1.84433	0.5650
29	0.47662	2.09811	0.49822	2.00715	0.52020	1.92235	0.54258	1.84305	0.56539
30	0.47698	2.09654	0.49858	2.00569	0.52057	1.92098	0.54296	1.84177	0.56577
31	0.47733	2.09498	0.49894	2.00423	0.52094	1.91962	0.54333	1.84049	0.56616
32	0.47769	2.09341	0.49931	2.00277	0.52131	1.91826	0.54371	1.83922	0.56654
33	0.47805	2.09184	0.49967	2.00131	0.52168	1.91690	0.54409	1.83794	0.56693
34	0.47840	2.09028	0.50004	1.99986	0.52205	1.91554	0.54446	1.83667	0.5673
35	0.47876	2.08872	0.50040	1.99841	0.52242	1.91418	0.54484	1.83540	0.5676
36	0.47912	2.08716	0.50076	1.99695	0.52279	1.91282	0.54522	1.83413	0.56808
37	0.47948	2.08560	0.50113	1.99550	0.52316	1.91147	0.54560	1.83286	0.56846
38	0.47984	2.08405	0.50149	1.99406	0.52353	1.91012	0.54597	1.83159	0.56885
39	0.48019	2.08250	0.50185	1.99261	0.52390	1.90876	0.54635	1.83033	0.5692
40	0.48055	2.08094	0.50222	1.99116	0.52427	1.90741	0.54673	1.82906	0.5696
41	0.48091	2.07939	0.50258	1.98972	0.52464	1.90607	0.54711	1.82780	0.5700
42	0.48127	2.07785	0.50295	1.98828	0.52501	1.90472	0.54748	1.82654	0.5703
43	0.48163	2.07630	0.50331	1.98684	0.52538	1.90337	0.54786	1.82528	0.5707
44	0.48198	2.07476	0.50368	1.98540	0.52575	1.90203	0.54824	1.82402	0.5711
45	0.48234	2.07321	0.50404	1.98396	0.52613	1.90069	0.54862	1.82276	0.5715
46	0.48270	2.07167	0.50441	1.98253	0.52650	1.89935	0.54900	1.82150	0.5719
47	0.48306	2.07014	0.50477	1.98110	0.52687	1.89801	0.54938	1.82025	0.5723
48	0.48342	2.06860	0.50514	1.97966	0.52724	1.89667	0.54975	1.81899	0.5727
49	0.48378	2.06706	0.50550	1.97823	0.52761	1.89533	0.55013	1.81774	0.5730
50	0.48414	2.06553	0.50587	1.97681	0.52798	1.89400	0.55051	1.81649	0.5734
51	0.48450	2.06400	0.50623	1.97538	0.52836	1.89266	0.55089	1.81524	0.5738
52	0.48486	2.06247	0.50660	1.97395	0.52873	1.89133	0.55127	1.81399	0.5742
53	0.48521	2.06094	0.50696	1.97253	0.52910	1.89000	0.55165	1.81274	0.5746
54	0.48557	2.05942	0.50733	1.97111	0.52947	1.88867	0.55203	1.81150	0.5750
55	0.48593	2.05790	0.50769	1.96969	0.52985	1.88734	0.55241	1.81025	0.5754
56	0.48629	2.05637	0.50806	1.96827	0.53022	1.88602	0.55279	1.80901	0.5758
57	0.48665	2.05485	0.50843	1.96685	0.53059	1.88469	0.55317	1.80777	0.5761
58	0.48701	2.05333	0.50879	1.96544	0.53096	1.88337	0.55355	1.80653	0.5765
59	0.48737	2.05182	0.50916	1.96402	0.53134	1.88205	0.55393	1.80529	0.5769
60	0.48773	2.05030	0.50953	1.96261	0.53171	1.88073	0.55431	1.80405	0.5773

	COT	TAN	COT	TAN	COT	TAN	COT	TAN	CO
	64°		63°		62°		61°		

30°		31°		32°		33°		34°		
TAN	COT	TAN	COT	TAN	COT	TAN	COT	TAN	COT	
.57735	1.73205	0.60086	1.66428	0.62487	1.60033	0.64941	1.53986	0.67451	1.48256	60
.57774	1.73089	0.60126	1.66318	0.62527	1.59930	0.64982	1.53888	0.67493	1.48163	59
.57813	1.72973	0.60165	1.66209	0.62568	1.59826	0.65024	1.53791	0.67536	1.48070	58
.57851	1.72857	0.60205	1.66099	0.62608	1.59723	0.65065	1.53693	0.67578	1.47977	57
.57890	1.72741	0.60245	1.65990	0.62649	1.59620	0.65106	1.53595	0.67620	1.47885	56
.57929	1.72625	0.60284	1.65881	0.62689	1.59517	0.65148	1.53497	0.67663	1.47792	55
.57968	1.72509	0.60324	1.65772	0.62730	1.59414	0.65189	1.53400	0.67705	1.47699	54
.58007	1.72393	0.60364	1.65663	0.62770	1.59311	0.65231	1.53302	0.67748	1.47607	53
.58046	1.72278	0.60403	1.65554	0.62811	1.59208	0.65272	1.53205	0.67790	1.47514	52
.58085	1.72163	0.60443	1.65445	0.62852	1.59105	0.65314	1.53107	0.67832	1.47422	51
.58124	1.72047	0.60483	1.65337	0.62892	1.59002	0.65355	1.53010	0.67875	1.47330	50
.58162	1.71932	0.60522	1.65228	0.62933	1.58900	0.65397	1.52913	0.67917	1.47238	49
.58201	1.71817	0.60562	1.65120	0.62973	1.58797	0.65438	1.52816	0.67960	1.47146	48
.58240	1.71702	0.60602	1.65011	0.63014	1.58695	0.65480	1.52719	0.68002	1.47053	47
.58279	1.71588	0.60642	1.64903	0.63055	1.58593	0.65521	1.52622	0.68045	1.46962	46
.58318	1.71473	0.60681	1.64795	0.63095	1.58490	0.65563	1.52525	0.68088	1.46870	45
.58357	1.71358	0.60721	1.64687	0.63136	1.58388	0.65604	1.52429	0.68130	1.46778	44
.58396	1.71244	0.60761	1.64579	0.63177	1.58286	0.65646	1.52332	0.68173	1.46686	43
.58435	1.71129	0.60801	1.64471	0.63217	1.58184	0.65688	1.52235	0.68215	1.46595	42
.58474	1.71015	0.60841	1.64363	0.63258	1.58083	0.65729	1.52139	0.68258	1.46503	41
.58513	1.70901	0.60881	1.64256	0.63299	1.57981	0.65771	1.52043	0.68301	1.46411	40
.58552	1.70787	0.60921	1.64148	0.63340	1.57879	0.65813	1.51946	0.68343	1.46320	39
.58591	1.70673	0.60960	1.64041	0.63380	1.57778	0.65854	1.51850	0.68386	1.46229	38
.58631	1.70560	0.61000	1.63934	0.63421	1.57676	0.65896	1.51754	0.68429	1.46137	37
.58670	1.70446	0.61040	1.63826	0.63462	1.57575	0.65938	1.51658	0.68471	1.46046	36
.58709	1.70332	0.61080	1.63719	0.63503	1.57474	0.65980	1.51562	0.68514	1.45955	35
.58748	1.70219	0.61120	1.63612	0.63544	1.57372	0.66021	1.51466	0.68557	1.45864	34
.58787	1.70106	0.61160	1.63505	0.63584	1.57271	0.66063	1.51370	0.68600	1.45773	33
.58826	1.69992	0.61200	1.63398	0.63625	1.57170	0.66105	1.51275	0.68642	1.45682	32
.58865	1.69879	0.61240	1.63292	0.63666	1.57069	0.66147	1.51179	0.68685	1.45592	31
.58905	1.69766	0.61280	1.63185	0.63707	1.56969	0.66189	1.51084	0.68728	1.45501	30
.58944	1.69653	0.61320	1.63079	0.63748	1.56868	0.66230	1.50988	0.68771	1.45410	29
.58983	1.69541	0.61360	1.62972	0.63789	1.56767	0.66272	1.50893	0.68814	1.45320	28
.59022	1.69428	0.61400	1.62866	0.63830	1.56667	0.66314	1.50797	0.68857	1.45229	27
.59061	1.69316	0.61440	1.62760	0.63871	1.56566	0.66356	1.50702	0.68900	1.45139	26
.59101	1.69203	0.61480	1.62654	0.63912	1.56466	0.66398	1.50607	0.68942	1.45049	25
.59140	1.69091	0.61520	1.62548	0.63953	1.56366	0.66440	1.50512	0.68985	1.44958	24
.59179	1.68979	0.61561	1.62442	0.63994	1.56265	0.66482	1.50417	0.69028	1.44868	23
.59218	1.68866	0.61601	1.62336	0.64035	1.56165	0.66524	1.50322	0.69071	1.44778	22
.59258	1.68754	0.61641	1.62230	0.64076	1.56065	0.66566	1.50228	0.69114	1.44688	21
.59297	1.68643	0.61681	1.62125	0.64117	1.55966	0.66608	1.50133	0.69157	1.44598	20
.59336	1.68531	0.61721	1.62019	0.64158	1.55866	0.66650	1.50038	0.69200	1.44508	19
.59376	1.68419	0.61761	1.61914	0.64199	1.55766	0.66692	1.49944	0.69243	1.44418	18
.59415	1.68308	0.61801	1.61808	0.64240	1.55666	0.66734	1.49849	0.69286	1.44329	17
.59454	1.68196	0.61842	1.61703	0.64281	1.55567	0.66776	1.49755	0.69329	1.44239	16
.59494	1.68085	0.61882	1.61598	0.64322	1.55467	0.66818	1.49661	0.69372	1.44149	15
.59533	1.67974	0.61922	1.61493	0.64363	1.55368	0.66860	1.49566	0.69416	1.44060	14
.59573	1.67863	0.61962	1.61388	0.64404	1.55269	0.66902	1.49472	0.69459	1.43970	13
.59612	1.67752	0.62003	1.61283	0.64446	1.55170	0.66944	1.49378	0.69502	1.43881	12
.59651	1.67641	0.62043	1.61179	0.64487	1.55071	0.66986	1.49284	0.69545	1.43792	11
.59691	1.67530	0.62083	1.61074	0.64528	1.54972	0.67028	1.49190	0.69588	1.43703	10
.59730	1.67419	0.62124	1.60970	0.64569	1.54873	0.67071	1.49097	0.69631	1.43614	9
.59770	1.67309	0.62164	1.60865	0.64610	1.54774	0.67113	1.49003	0.69675	1.43525	8
.59809	1.67198	0.62204	1.60761	0.64652	1.54675	0.67155	1.48909	0.69718	1.43436	7
.59849	1.67088	0.62245	1.60657	0.64693	1.54576	0.67197	1.48816	0.69761	1.43347	6
.59888	1.66978	0.62285	1.60553	0.64734	1.54478	0.67239	1.48722	0.69804	1.43258	5
.59928	1.66867	0.62325	1.60449	0.64775	1.54379	0.67282	1.48629	0.69847	1.43169	4
.59967	1.66757	0.62366	1.60345	0.64817	1.54281	0.67324	1.48536	0.69891	1.43080	3
.60007	1.66647	0.62406	1.60241	0.64858	1.54183	0.67366	1.48442	0.69934	1.42992	2
.60046	1.66538	0.62446	1.60137	0.64899	1.54085	0.67409	1.48349	0.69977	1.42903	1
.60086	1.66428	0.62487	1.60033	0.64941	1.53986	0.67451	1.48256	0.70021	1.42815	0
COT	TAN	COT	TAN	COT	TAN	COT	TAN	COT	TAN	M I N
59°		58°		57°		56°		55°		

MIN	35°		36°		37°		38°		TA
	TAN	COT	TAN	COT	TAN	COT	TAN	COT	
0	0.70021	1.42815	0.72654	1.37638	0.75355	1.32704	0.78129	1.27994	0.809
1	0.70064	1.42726	0.72699	1.37554	0.75401	1.32624	0.78175	1.27917	0.810
2	0.70107	1.42638	0.72743	1.37470	0.75447	1.32544	0.78222	1.27841	0.810
3	0.70151	1.42550	0.72788	1.37386	0.75492	1.32464	0.78269	1.27764	0.811
4	0.70194	1.42462	0.72832	1.37302	0.75538	1.32384	0.78316	1.27688	0.811
5	0.70238	1.42374	0.72877	1.37218	0.75584	1.32304	0.78363	1.27611	0.812
6	0.70281	1.42286	0.72921	1.37134	0.75629	1.32224	0.78410	1.27535	0.812
7	0.70325	1.42198	0.72966	1.37050	0.75675	1.32144	0.78457	1.27458	0.813
8	0.70368	1.42110	0.73010	1.36967	0.75721	1.32064	0.78504	1.27382	0.813
9	0.70412	1.42022	0.73055	1.36883	0.75767	1.31984	0.78551	1.27306	0.814
10	0.70455	1.41934	0.73100	1.36800	0.75812	1.31904	0.78598	1.27230	0.814
11	0.70499	1.41847	0.73144	1.36716	0.75858	1.31825	0.78645	1.27153	0.815
12	0.70542	1.41759	0.73189	1.36633	0.75904	1.31745	0.78692	1.27077	0.815
13	0.70586	1.41672	0.73234	1.36549	0.75950	1.31666	0.78739	1.27001	0.816
14	0.70629	1.41584	0.73278	1.36466	0.75996	1.31586	0.78786	1.26925	0.816
15	0.70673	1.41497	0.73323	1.36383	0.76042	1.31507	0.78834	1.26849	0.817
16	0.70717	1.41409	0.73368	1.36300	0.76088	1.31427	0.78881	1.26774	0.817
17	0.70760	1.41322	0.73413	1.36217	0.76134	1.31348	0.78928	1.26698	0.818
18	0.70804	1.41235	0.73457	1.36134	0.76180	1.31269	0.78975	1.26622	0.818
19	0.70848	1.41148	0.73502	1.36051	0.76226	1.31190	0.79022	1.26546	0.818
20	0.70891	1.41061	0.73547	1.35968	0.76272	1.31110	0.79070	1.26471	0.819
21	0.70935	1.40974	0.73592	1.35885	0.76318	1.31031	0.79117	1.26395	0.819
22	0.70979	1.40887	0.73637	1.35802	0.76364	1.30952	0.79164	1.26319	0.820
23	0.71023	1.40800	0.73681	1.35719	0.76410	1.30873	0.79212	1.26244	0.820
24	0.71066	1.40714	0.73726	1.35637	0.76456	1.30795	0.79259	1.26169	0.821
25	0.71110	1.40627	0.73771	1.35554	0.76502	1.30716	0.79306	1.26093	0.821
26	0.71154	1.40540	0.73816	1.35472	0.76548	1.30637	0.79354	1.26018	0.822
27	0.71198	1.40454	0.73861	1.35389	0.76594	1.30558	0.79401	1.25943	0.822
28	0.71242	1.40367	0.73906	1.35307	0.76640	1.30480	0.79449	1.25867	0.823
29	0.71285	1.40281	0.73951	1.35224	0.76686	1.30401	0.79496	1.25792	0.823
30	0.71329	1.40195	0.73996	1.35142	0.76733	1.30323	0.79544	1.25717	0.824
31	0.71373	1.40109	0.74041	1.35060	0.76779	1.30244	0.79591	1.25642	0.824
32	0.71417	1.40022	0.74086	1.34978	0.76825	1.30166	0.79639	1.25567	0.825
33	0.71461	1.39936	0.74131	1.34896	0.76871	1.30087	0.79686	1.25492	0.825
34	0.71505	1.39850	0.74176	1.34814	0.76918	1.30009	0.79734	1.25417	0.826
35	0.71549	1.39764	0.74221	1.34732	0.76964	1.29931	0.79781	1.25343	0.826
36	0.71593	1.39679	0.74267	1.34650	0.77010	1.29853	0.79829	1.25268	0.827
37	0.71637	1.39593	0.74312	1.34568	0.77057	1.29775	0.79877	1.25193	0.827
38	0.71681	1.39507	0.74357	1.34487	0.77103	1.29696	0.79924	1.25118	0.828
39	0.71725	1.39421	0.74402	1.34405	0.77149	1.29618	0.79972	1.25044	0.828
40	0.71769	1.39336	0.74447	1.34323	0.77196	1.29541	0.80020	1.24969	0.829
41	0.71813	1.39250	0.74492	1.34242	0.77242	1.29463	0.80067	1.24895	0.829
42	0.71857	1.39165	0.74538	1.34160	0.77289	1.29385	0.80115	1.24820	0.830
43	0.71901	1.39079	0.74583	1.34079	0.77335	1.29307	0.80163	1.24746	0.830
44	0.71946	1.38994	0.74628	1.33998	0.77382	1.29229	0.80211	1.24672	0.83
45	0.71990	1.38909	0.74674	1.33916	0.77428	1.29152	0.80258	1.24597	0.83
46	0.72034	1.38824	0.74719	1.33835	0.77475	1.29074	0.80306	1.24523	0.83
47	0.72078	1.38738	0.74764	1.33754	0.77521	1.28997	0.80354	1.24449	0.83
48	0.72122	1.38653	0.74810	1.33673	0.77568	1.28919	0.80402	1.24375	0.83
49	0.72167	1.38568	0.74855	1.33592	0.77615	1.28842	0.80450	1.24301	0.83
50	0.72211	1.38484	0.74900	1.33511	0.77661	1.28764	0.80498	1.24227	0.83
51	0.72255	1.38399	0.74946	1.33430	0.77708	1.28687	0.80546	1.24153	0.83
52	0.72299	1.38314	0.74991	1.33349	0.77754	1.28610	0.80594	1.24079	0.83
53	0.72344	1.38229	0.75037	1.33268	0.77801	1.28533	0.80642	1.24005	0.83
54	0.72388	1.38145	0.75082	1.33187	0.77848	1.28456	0.80690	1.23931	0.83
55	0.72432	1.38060	0.75128	1.33107	0.77895	1.28379	0.80738	1.23858	0.83
56	0.72477	1.37976	0.75173	1.33026	0.77941	1.28302	0.80786	1.23784	0.83
57	0.72521	1.37891	0.75219	1.32946	0.77988	1.28225	0.80834	1.23710	0.83
58	0.72565	1.37807	0.75264	1.32865	0.78035	1.28148	0.80882	1.23637	0.83
59	0.72610	1.37722	0.75310	1.32785	0.78082	1.28071	0.80930	1.23563	0.83
60	0.72654	1.37638	0.75355	1.32704	0.78129	1.27994	0.80978	1.23490	0.83
	COT	TAN	COT	TAN	COT	TAN	COT	TAN	CO
	54°		53°		52°		51°		

M I N	40°		41°		42°		43°		44°	
	TAN	COT	TAN	COT	TAN	COT	TAN	COT	TAN	COT
0	0.83910	1.19175	0.86929	1.15037	0.90040	1.11061	0.93252	1.07237	0.96569	1.03553
1	0.83960	1.19105	0.86980	1.14969	0.90093	1.10996	0.93306	1.07174	0.96625	1.03493
2	0.84009	1.19035	0.87031	1.14902	0.90146	1.10931	0.93360	1.07112	0.96681	1.03433
3	0.84059	1.18964	0.87082	1.14834	0.90199	1.10867	0.93415	1.07049	0.96738	1.03372
4	0.84108	1.18894	0.87133	1.14767	0.90251	1.10802	0.93469	1.06987	0.96794	1.03312
5	0.84158	1.18824	0.87184	1.14699	0.90304	1.10737	0.93524	1.06925	0.96850	1.03252
6	0.84208	1.18754	0.87236	1.14632	0.90357	1.10672	0.93578	1.06862	0.96907	1.03192
7	0.84258	1.18684	0.87287	1.14565	0.90410	1.10607	0.93633	1.06800	0.96963	1.03132
8	0.84307	1.18614	0.87338	1.14498	0.90463	1.10543	0.93688	1.06738	0.97020	1.03072
9	0.84357	1.18544	0.87389	1.14430	0.90516	1.10478	0.93742	1.06676	0.97076	1.03012
10	0.84407	1.18474	0.87441	1.14363	0.90569	1.10414	0.93797	1.06613	0.97133	1.02952
11	0.84457	1.18404	0.87492	1.14296	0.90621	1.10349	0.93852	1.06551	0.97189	1.02892
12	0.84507	1.18334	0.87543	1.14229	0.90674	1.10285	0.93906	1.06489	0.97246	1.02832
13	0.84556	1.18264	0.87595	1.14162	0.90727	1.10220	0.93961	1.06427	0.97302	1.02772
14	0.84606	1.18194	0.87646	1.14095	0.90781	1.10156	0.94016	1.06365	0.97359	1.02713
15	0.84656	1.18125	0.87698	1.14028	0.90834	1.10091	0.94071	1.06303	0.97416	1.02653
16	0.84706	1.18055	0.87749	1.13961	0.90887	1.10027	0.94125	1.06241	0.97472	1.02593
17	0.84756	1.17986	0.87801	1.13894	0.90940	1.09963	0.94180	1.06179	0.97529	1.02533
18	0.84806	1.17916	0.87852	1.13828	0.90993	1.09899	0.94235	1.06117	0.97586	1.02474
19	0.84856	1.17846	0.87904	1.13761	0.91046	1.09834	0.94290	1.06056	0.97643	1.02414
20	0.84906	1.17777	0.87955	1.13694	0.91099	1.09770	0.94345	1.05994	0.97700	1.02355
21	0.84956	1.17708	0.88007	1.13627	0.91153	1.09706	0.94400	1.05932	0.97756	1.02295
22	0.85006	1.17638	0.88059	1.13561	0.91206	1.09642	0.94455	1.05870	0.97813	1.02236
23	0.85057	1.17569	0.88110	1.13494	0.91259	1.09578	0.94510	1.05809	0.97870	1.02176
24	0.85107	1.17500	0.88162	1.13428	0.91313	1.09514	0.94565	1.05747	0.97927	1.02117
25	0.85157	1.17430	0.88214	1.13361	0.91366	1.09450	0.94620	1.05685	0.97984	1.02057
26	0.85207	1.17361	0.88265	1.13295	0.91419	1.09386	0.94676	1.05624	0.98041	1.01998
27	0.85257	1.17292	0.88317	1.13228	0.91473	1.09322	0.94731	1.05562	0.98098	1.01939
28	0.85308	1.17223	0.88369	1.13162	0.91526	1.09258	0.94786	1.05501	0.98155	1.01879
29	0.85358	1.17154	0.88421	1.13096	0.91580	1.09195	0.94841	1.05439	0.98213	1.01820
30	0.85408	1.17085	0.88473	1.13029	0.91633	1.09131	0.94896	1.05378	0.98270	1.01761
31	0.85458	1.17016	0.88524	1.12963	0.91687	1.09067	0.94952	1.05317	0.98327	1.01702
32	0.85509	1.16947	0.88576	1.12897	0.91740	1.09003	0.95007	1.05255	0.98384	1.01642
33	0.85559	1.16878	0.88628	1.12831	0.91794	1.08940	0.95062	1.05194	0.98441	1.01583
34	0.85609	1.16809	0.88680	1.12765	0.91847	1.08876	0.95118	1.05133	0.98499	1.01524
35	0.85660	1.16741	0.88732	1.12699	0.91901	1.08813	0.95173	1.05072	0.98556	1.01465
36	0.85710	1.16672	0.88784	1.12633	0.91955	1.08749	0.95229	1.05010	0.98613	1.01406
37	0.85761	1.16603	0.88836	1.12567	0.92008	1.08686	0.95284	1.04949	0.98671	1.01347
38	0.85811	1.16535	0.88888	1.12501	0.92062	1.08622	0.95340	1.04888	0.98728	1.01288
39	0.85862	1.16466	0.88940	1.12435	0.92116	1.08559	0.95395	1.04827	0.98786	1.01229
40	0.85912	1.16398	0.88992	1.12369	0.92170	1.08496	0.95451	1.04766	0.98843	1.01170
41	0.85963	1.16329	0.89045	1.12303	0.92224	1.08432	0.95506	1.04705	0.98901	1.01112
42	0.86014	1.16261	0.89097	1.12238	0.92277	1.08369	0.95562	1.04644	0.98958	1.01053
43	0.86064	1.16192	0.89149	1.12172	0.92331	1.08306	0.95618	1.04583	0.99016	1.00994
44	0.86115	1.16124	0.89201	1.12106	0.92385	1.08243	0.95673	1.04522	0.99073	1.00935
45	0.86166	1.16056	0.89253	1.12041	0.92439	1.08179	0.95729	1.04461	0.99131	1.00876
46	0.86216	1.15987	0.89306	1.11975	0.92493	1.08116	0.95785	1.04401	0.99189	1.00818
47	0.86267	1.15919	0.89358	1.11909	0.92547	1.08053	0.95841	1.04340	0.99247	1.00759
48	0.86318	1.15851	0.89410	1.11844	0.92601	1.07990	0.95897	1.04279	0.99304	1.00701
49	0.86368	1.15783	0.89463	1.11778	0.92655	1.07927	0.95952	1.04218	0.99362	1.00642
50	0.86419	1.15715	0.89515	1.11713	0.92709	1.07864	0.96008	1.04158	0.99420	1.00583
51	0.86470	1.15647	0.89567	1.11648	0.92763	1.07801	0.96064	1.04097	0.99478	1.00525
52	0.86521	1.15579	0.89620	1.11582	0.92817	1.07738	0.96120	1.04036	0.99536	1.00466
53	0.86572	1.15511	0.89672	1.11517	0.92872	1.07676	0.96176	1.03976	0.99594	1.00408
54	0.86623	1.15443	0.89725	1.11452	0.92926	1.07613	0.96232	1.03915	0.99652	1.00350
55	0.86674	1.15375	0.89777	1.11387	0.92980	1.07550	0.96288	1.03855	0.99710	1.00291
56	0.86725	1.15308	0.89830	1.11321	0.93034	1.07487	0.96344	1.03794	0.99768	1.00233
57	0.86776	1.15240	0.89883	1.11256	0.93088	1.07425	0.96400	1.03734	0.99826	1.00174
58	0.86827	1.15172	0.89935	1.11191	0.93143	1.07362	0.96457	1.03674	0.99884	1.00116
59	0.86878	1.15104	0.89988	1.11126	0.93197	1.07299	0.96513	1.03613	0.99942	1.00057
60	0.86929	1.15037	0.90040	1.11061	0.93252	1.07237	0.96569	1.03553	1.00000	1.00000

	COT	TAN	COT	TAN	COT	TAN	COT	TAN	COT	TAN
	49°		48°		47°		46°		45°	

INDEX

A

Abscissa, 4-2
Acute angles, 5-1, 5-20
Addition, vector, 7-2 to 7-3, 7-6 to 7-9, 7-23, 7-24
Angles,
  Acute, 5-1, 5-20
  Complementary, 4-18 to 4-20, 4-33
  Coterminal, 4-3 to 4-4, 4-16 to 4-17, 4-30, 4-33
  Negative, 3-2, 3-31, 4-16, 4-32, 6-5, 6-34
  Obtuse, 5-1, 5-20
  Positive, 3-2, 3-31
  Quadrantal, 4-3, 4-23 to 4-24, 4-33
  Reference, 4-11 to 4-12, 4-31
Angles in standard position, 4-2 to 4-4, 4-30
Angles, measurements of, 3-1 to 3-19, 3-31 to 3-33
  Degrees, 3-2 to 3-3, 3-31
  Mils, 3-14 to 3-19, 3-33
  Radians, 3-4 to 3-13, 3-31 to 3-33
    Angular velocity, 3-8 to 3-11, 3-32
    Area of a sector, 3-12 to 3-13, 3-33
Angles, special, 4-21 to 4-24, 4-33
  Frequently used angles, 4-21 to 4-23, 4-33
    30°-60° angles, 4-21 to 4-22, 4-33
    45° angles, 4-22 to 4-23, 4-33
  Quadrantal angles, 4-23 to 4-24, 4-33
Angular velocity, 3-8 to 3-11, 3-32
Antilogarithm, 1-16 to 1-18, 1-24
Area formulas, 5-17 to 5-19, 5-21
Area of a sector, 3-12 to 3-13, 3-33

C

Characteristic, 1-9 to 1-10, 1-23
  Negative, 1-10, 1-23
  Positive, 1-9 to 1-10, 1-23
Cofunctions, 4-18 to 4-20, 4-33
Common logarithms, 1-8 to 1-10, 1-22 1-23
  Negative characteristics, 1-10, 1-2
  Positive characteristics, 1-9 to 1-10, 1-23
Complementary angles, 4-18 to 4-20, 4-33
Components of vectors, 7-4 to 7-6, 7-2 to 7-24
Computations with logarithms, 2-1 to 2-17
  Algebraic operations, 2-9 to 2-11,
  Applications, 2-12 to 2-13, 2-15
  Division, 2-5 to 2-6, 2-15
  Introduction, 2-1 to 2-2
  Multiplication, 2-2 to 2-4, 2-15
  Powers, 2-7, 2-15
  Roots, 2-7 to 2-8, 2-15
  Summary, 2-15
Coordinates, 4-2
Coterminal angles, 4-3 to 4-4, 4-30
Coterminal angles, functions of, 4-16 t 4-33

D

Degrees, 3-2 to 3-3, 3-31
Direction of a vector, 7-5, 7-24
Division with logarithms, 1-4 to 1-5, 1-22, 2-5 to 2-6, 2-15
Double-angle formulas, 6-16 to 6-18, 6-36

E

Equilibriant force, 7-13, 7-25
Equilibrium, 7-13 to 7-21, 7-25
  Rotational, 7-19 to 7-21, 7-25
  Translational, 7-15 to 7-19, 7-25
Exponential form, 1-2

F

Forces, 7-11 to 7-12, 7-24 to 7-25
Fundamental identities, 6-1 to 6-7, 6-34
  Negative angles, 6-5, 6-34
  Pythagorean, 6-3 to 6-5, 6-34
  Quotient, 6-3, 6-34
  Reciprocal, 6-2, 6-34
  Verifying trigonometric identities, 6-6 to 6-7

G

Graph of the cosine function, 4-27 to 4-28
Graph of the sine function, 4-27
Graph of the tangent function, 4-28

H

Half-angle formulas, 6-18 to 6-20, 6-36
Horizontal component, 7-4, 7-23

I

Identities, formulas for, 6-8 to 6-20, 6-35 to 6-36
  Double-angle formulas, 6-16 to 6-18, 6-36
  Half-angle formulas, 6-18 to 6-20, 6-36
  Sum and difference formulas, 6-8 to 6-15, 6-35
Identities, fundamental, 6-1 to 6-7, 6-34
  Negative angles, 6-5, 6-34
  Pythagorean, 6-3 to 6-5, 6-34
  Quotient, 6-3, 6-34
  Reciprocal, 6-2, 6-34
  Verifying trigonometric identities, 6-6 to 6-7
Identity, 6-1
Initial position, 3-2, 3-31
Interpolation, 1-13 to 1-15, 1-23 to 1-24
Inverse trigonometric functions, 6-21 to 6-27, 6-36
  Notation, 6-22, 6-36
  Principal values, 6-22 to 6-27, 6-36

L

Law of cosines, 5-11 to 5-16, 5-
  Case 3. Two sides and the i 5-13 to 5-15, 5-20
  Case 4. All three sides, 5-15
Law of sines, 5-2 to 5-10, 5-20
  Case 1. One side and two a 5-5, 5-20
  Case 2. Two sides and an a one of them, 5-5 to 5-10,
Length of arc, 3-7, 3-32
Linear velocity, 3-9, 3-33
Logarithm tables, 1-11 to 1-12,
Logarithmic form, 1-2
Logarithms, 1-1 to 1-27
  Antilogarithm, 1-16 to 1-18
  Common logarithms, 1-8 to 1-23
    Negative characteristics
    Positive characteristics, 1-23
  Interpolation, 1-13 to 1-15,
  Introduction, 1-1
  Laws for calculation, 1-3 to
    Division, 1-4 to 1-5, 1-
    Multiplication, 1-3 to
    Powers, 1-6, 1-22
    Roots, 1-6 to 1-7, 1-22
  Logarithm tables, 1-11 to 1
  Natural logarithms, 1-19 to 1-25
  Review of definitions, 1-1
  Summary, 1-22 to 1-25
Logarithms, computations with
  Algebraic operations, 2-9 t
  Applications, 2-12 to 2-13,
  Division, 2-5 to 2-6, 2-15
  Multiplication, 2-2 to 2-4,
  Powers, 2-7, 2-15
  Roots, 2-7 to 2-8, 2-15

M

Magnitude of a vector, 7-5, 7-2
Mantissa, 1-9, 1-23
Mils, 3-14 to 3-19, 3-33
Multiplication with logarithms, to 1-4, 1-22, 2-2 to 2-4, 2-15

N

Natural logarithms, 1-19 to 1-21, 1-24 to 1-25
Negative angle, 3-2, 3-31
Negative angles, functions of, 4-16, 4-32
Negative angles, identities for, 6-5, 6-34

O

Oblique triangles, 5-1 to 5-23
  Area formulas, 5-17 to 5-19, 5-21
  Introduction, 5-1 to 5-2
  Methods of solving oblique triangles, 5-2 to 5-16, 5-20
    Law of cosines, 5-11 to 5-16, 5-20
      Case 3. Two sides and the included angle, 5-13 to 5-15, 5-20
      Case 4. All three sides, 5-15 to 5-16, 5-20
    Law of sines, 5-2 to 5-10, 5-20
      Case 1. One side and two angles, 5-4 to 5-5, 5-20
      Case 2. Two sides and an angle opposite one of them, 5-5 to 5-10, 5-20
  Summary, 5-20 to 5-21
Obtuse angles, 5-1, 5-20
Ordinate, 4-2

P

Periodic, 4-27, 4-33
Periods of the trigonometric functions, 4-27 to 4-28, 4-33
  Graph of the cosine function, 4-27 to 4-28
  Graph of the sine function, 4-27
  Graph of the tangent function, 4-28
Positive angle, 3-2, 3-31
Powers, logarithms of, 1-6, 1-22, 2-7, 2-15
Principal values, 6-22 to 6-27, 6-36
Pythagorean identities, 6-3 to 6-5, 6-34
Pythagorean theorem, 3-20 to 3-21, 3-34

Q

Quadrant, functions of angles in any, 4-12 to 4-15, 4-31 to 4-32
  Quadrant I, 4-13, 4-31
  Quadrant II, 4-13 to 4-14, 4-31
  Quadrant III, 4-14 to 4-15, 4-32
  Quadrant IV, 4-15, 4-32
Quadrant system, 4-7 to 4-10, 4-30
Quadrantal angles, 4-3, 4-23 to 4-24,
Quadrants, 3-2, 3-31
Quotient identities, 6-3, 6-34

R

Radians, 3-4 to 3-13, 3-31 to 3-33
  Angular velocity, 3-8 to 3-11, 3-3
  Area of a sector, 3-12 to 3-13, 3-
Radius vector, 3-2, 3-31
Reciprocal identities, 6-2, 6-34
Rectangular coordinate system, 4-1 to
Reduction formulas, 4-12, 4-31
Reference angle, 4-11 to 4-12, 4-31
Reference triangle, 4-12, 4-31
Resolution of a force, 7-11, 7-25
Resultant vector, 7-2 to 7-3, 7-11, 7-2
Right triangles, properties of, 3-20 to 3-34
  Pythagorean theorem, 3-20 to 3-2
  Similar right triangles, 3-21 to 3-2
Roots, logarithms of, 1-6 to 1-7, 1-22, 2-8, 2-15
Rotational equilibrium, 7-19 to 7-21,

S

Scalar quantity, 7-1, 7-23
Scientific notation, 1-9 to 1-10, 1-23
Significant digits, 2-2, 2-15
Similar right triangles, 3-21 to 3-23, 3-
Similar triangles, 3-21, 3-34
Special angles, 4-21 to 4-24, 4-33
  Frequently used angles, 4-21 to 4-
    30°-60° angles, 4-21 to 4-22,
    45° angles, 4-22 to 4-23, 4-33
  Quadrantal angles, 4-23 to 4-24, 4
Subtraction, vector, 7-3 to 7-4, 7-23
Sum and difference formulas, 6-8 to 6-35

T

Terminal position, 3-2, 3-31
Torque, 7-19 to 7-21, 7-25
Translational equilibrium, 7-15 to 7-19
Trigonometric analysis, 4-1 to 4-35
  Angles in standard position, 4-2 t
    4-30
  Cofunctions and complementary a
    4-18 to 4-20, 4-33

Trigonometric analysis—Continued
Definition of the trigonometric functions, 4-4 to 4-6, 4-30
Functions of angles in any quadrant, 4-12 to 4-15, 4-31 to 4-32
Quadrant I, 4-13, 4-31
Quadrant II, 4-13 to 4-14, 4-31
Quadrant III, 4-14 to 4-15, 4-32
Quadrant IV, 4-15, 4-32
Functions of coterminal angles, 4-16 to 4-17, 4-33
Functions of negative angles, 4-16, 4-32
Introduction, 4-1
Periods of the trigonometric functions, 4-27 to 4-28, 4-33
Graph of the cosine function, 4-27 to 4-28
Graph of the sine function, 4-27
Graph of the tangent function, 4-28
Quadrant system, 4-7 to 4-10, 4-30
Rectangular coordinate system, 4-1 to 4-2
Reference angle, 4-11 to 4-12, 4-31
Special angles, 4-21 to 4-24, 4-33
Frequently used angles, 4-21 to 4-23, 4-33
30°-60° angles, 4-21 to 4-22, 4-33
45° angles, 4-22 to 4-23, 4-33
Quadrantal angles, 4-23 to 4-24, 4-33
Summary, 4-30 to 4-33
Trigonometric equations, 6-28 to 6-33, 6-36 to 6-37
Trigonometric functions, definition of, 4-4 to 4-6, 4-30
Trigonometric identities and equations, 6-1 to 6-39
Formulas for identities, 6-8 to 6-20, 6-35 to 6-36
Double-angle formulas, 6-16 to 6-18, 6-36
Half-angle formulas, 6-18 to 6-20, 6-36
Sum and difference formulas, 6-8 to 6-15, 6-35
Fundamental identities, 6-1 to 6-7, 6-34
Identities for negative angles, 6-5, 6-34
Pythagorean identities, 6-3 to 6-5, 6-34
Quotient identities, 6-3, 6-34
Reciprocal identities, 6-2, 6-34
Verifying trigonometric identities, 6-6 to 6-7
Trigonometric identities and eq
Continued
Introduction, 6-1
Inverse trigonometric funct
6-27, 6-36
Notation, 6-22, 6-36
Principal values, 6-22
Summary, 6-34 to 6-37
Trigonometric equations, 6
6-36 to 6-37
Trigonometric measurements, 3
Introduction, 3-1
Measuring angles, 3-1 to 3-
3-33
Degrees, 3-2 to 3-3, 3-
Mils, 3-14 to 3-19, 3-3
Radians, 3-4 to 3-13,
Angular velocity,
3-32
Area of a sector,
3-33
Properties of right triangle
3-34
Pythagorean theorem, 3
Similar right triangles,
3-34
Summary, 3-31 to 3-34
Trigonometric ratios, funct
tables, 3-24 to 3-29, 3-34
Tables of trigonometri
3-27 to 3-28, 3-34
Trigonometric ratios a
3-25 to 3-27, 3-34
Use of trigonometric r
functions, 3-29
Trigonometric ratios, functions
3-24 to 3-29, 3-34
Tables of trigonometric fu
3-28, 3-34
Trigonometric ratios and f
to 3-27, 3-34
Use of trigonometric ratios
functions, 3-29

V

Vector quantity, 7-1, 7-23
Vectors, 7-1 to 7-9, 7-23 to 7-2
Components of vectors, 7-
to 7-24
Vector addition, 7-2 to 7-3
Vector addition by compon
7-9, 7-24
Vector subtraction, 7-3 to

ectors and forces, 7-1 to 7-27
Equilibrium, 7-13 to 7-21, 7-25
Rotational equilibrium, 7-19 to 7-21, 7-25
Translational equilibrium, 7-15 to 7-19, 7-25
Forces, 7-11 to 7-12, 7-24 to 7-25
Introduction, 7-1
Summary, 7-23 to 7-25
Vectors and forces—Continued
Vectors, 7-1 to 7-9, 7-23 to 7-24
Components of vectors, 7-4 to 7-6, 7-23 to 7-24
Vector addition, 7-2 to 7-3, 7-23
Vector addition by components, 7-6 to 7-9, 7-24
Vector subtraction, 7-3 to 7-4, 7-23
Vertical component, 7-4, 7-23

☆ U.S. GOVERNMENT PRINTING OFFICE: 1989—631-071-00018

A FINAL QUESTION

What did you think of this text? Do you have any questions pertaining to this text? You and your command are urged to submit your constructive criticisms and your recommendations This tear-out form letter is provided for your convenience.

Date: _____

SSN: _____

From: _____
(Rank/Rate/Civilian Grade, Name)

_____

_____

To: Naval Education and Training Program
Management Support Activity (Code 3233)
Pensacola, Florida 32509-5000

Subj: TRAMAN MATHEMATICS, VOLUME 2-A, NAVEDTRA 10062

1. I have the following questions or comments pertaining the text:

(Fold along dotted line and staple or tape)

(Fold along dotted line and staple or tape)

DEPARTMENT OF THE NAVY

NAVAL EDUCATION AND TRAINING PROGRAM
MANAGEMENT SUPPORT ACTIVITY (Code 3233)
PENSACOLA, FL 32509-5000

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, $300

POSTAGE AND FE
NAVY DEPARTM
DoD-316

NAVAL EDUCATION AND TRAINING PROGRAM
MANAGEMENT SUPPORT ACTIVITY
BUILDING 2435 (Code 3233)
PENSACOLA, FL 32509-5000